

Chapter 28. Polar Regions**Coordinating Lead Authors**

Joan Nymand Larsen (Iceland), Oleg A. Anisimov (Russian Federation)

Lead Authors

Andrew Constable (Australia), Anne Hollowed (USA), Nancy Maynard (USA), Pål Prestrud (Norway), Terry Prowse (Canada), John Stone (Canada)

Contributing Authors

Terry Callaghan (UK), Mark Carey (USA), Peter Convey (UK), Andrew Derocher (Canada), Peter T. Fretwell (UK), Bruce C. Forbes (Finland), Solveig Glomsrød (Norway), Dominic Hodgson (UK), Eileen Hofmann (USA), Grete K. Hovelsrud (Norway), Gita L Ljubicic (Canada), Harald Loeng (Norway), Eugene Murphy (UK), Steve Nicol (Australia), Anders Oskal (Norway), James D. Reist (Canada), Phil Trathan (UK), Barbara Weinecke (Australia), Fred Wrona (Canada)

Review Editors

Maria Ananicheva (Russian Federation), F. Stuart Chapin III (USA)

Volunteer Chapter Scientist

Vasiliy Kokorev (Russian Federation)

Contents

Executive Summary

28.1. Introduction

28.2. Observed Changes and Vulnerability under Multiple Stressors

28.2.1. Hydrology and Freshwater Ecosystems

28.2.1.1. Arctic

28.2.1.2. Antarctica

28.2.2. Oceanography and Marine Ecosystems

28.2.2.1. Arctic

28.2.2.2. Antarctica

28.2.3. Terrestrial Ecosystems

28.2.3.1. Arctic

28.2.3.2. Antarctica

28.2.4. Health and Well-Being of Arctic Residents

28.2.4.1. Direct Impacts of a Changing Climate on the Health of Arctic Residents

28.2.4.2. Indirect Impacts of Climate Change on the Health of Arctic Residents

28.2.5. Indigenous Peoples and Traditional Knowledge

28.2.6. Economic Sectors

28.2.6.1. Arctic

28.2.6.2. Antarctica and the Southern Ocean

28.3. Key Projected Impacts and Vulnerabilities

28.3.1. Hydrology and Freshwater Ecosystems

28.3.1.1. Arctic

28.3.1.2. Antarctica

28.3.2. Oceanography and Marine Ecosystems

28.3.2.1. Ocean Acidification in the Arctic and Antarctic

28.3.2.2. Arctic

- 28.3.2.3. Antarctica and the Southern Ocean
- 28.3.3. Terrestrial Environment and Related Ecosystems
 - 28.3.3.1. Arctic
 - 28.3.3.2. Antarctica
- 28.3.4. Economic Sectors
 - 28.3.4.1. Fisheries
 - 28.3.4.2. Forestry and Farming
 - 28.3.4.3. Infrastructure, Transportation, and Terrestrial Resources
- 28.4. Human Adaptation
- 28.5. Research and Data Gaps

References

Frequently Asked Questions

- 28.1: What will be the net socio-economic impacts of change in the polar regions?
- 28.2: Why are changes in sea ice so important to the polar regions?

Executive Summary

Additional and stronger scientific evidence has accumulated since the AR4 that reinforces key findings made in the Fourth Assessment Report (AR4).

The impacts of climate change, and the adaptations to it, exhibit strong spatial heterogeneity in the Polar Regions because of the high diversity of social systems, bio-physical regions and associated drivers of change (*high confidence*). [28.2.2] For example, the tree line has moved northward and upward in many, but not all, Arctic areas (*high confidence*) and significant increases in tall shrubs and grasses have been observed in many places (*very high confidence*). [28.2.3.2]

Some marine species will shift their ranges in response to changing ocean and sea ice conditions in the Polar Regions (*medium confidence*). The response rate and the spatial extent of the shifts will differ by species based on their vulnerability to change and their life history. [28.2.2; 28.3.2] Loss of sea ice in summer and increased ocean temperature is expected to enhance secondary pelagic production in some but not all regions of the Arctic Ocean with associated changes in the energy pathways within the marine ecosystem (*medium confidence*). These changes are expected to alter the species composition and carrying capacity with associated impacts on marine fish and shellfish populations (*medium confidence*). [28.2.2.1] Also, changes in sea ice and the physical environment to the west of the Antarctic Peninsula are altering phytoplankton stocks and productivity, and krill (*high confidence*). [28.2.2.2]

Climate change is impacting terrestrial and freshwater ecosystems in some areas of Antarctica and the Arctic. This is due to ecological effects resulting from reductions in the duration and extent of ice and snow cover and enhanced permafrost thaw (*very high confidence*), and through changes in the precipitation-evaporation balance (*medium confidence*). [28.2]

The primary concern for polar bears over the foreseeable future is the recent and projected loss of annual ice over continental shelves, decreased ice duration, and decreased ice thickness (*high confidence*). Of the two subpopulations where data are adequate for assessing abundance effects, it is *very likely* that the recorded population declines are caused by reductions in sea ice extent. [28.2.2.1.2; 28.3.2.2.2]

Rising temperatures, leading to the further thawing of permafrost and changing precipitation patterns have the potential to affect infrastructure and related services in the Arctic (*high confidence*). Particular concerns

are associated with the damage of residential buildings due to thawing permafrost, including, Arctic cities; small, rural settlements; and storage facilities for hazardous materials. [28.2.4; 28.2.4.2; 28.2.5]

In addition, there is new scientific evidence that has emerged since the AR4.

The physical, biological and socio-economic impacts of climate change in the Arctic have to be seen in the context of often interconnected factors that include not only environmental changes caused by drivers other than climate change but also demography, culture and economic development. Climate change has compounded some of the existing vulnerabilities caused by these other factors (*high confidence*). [28.2.4; 28.2.5; 28.4] For example, food security for many indigenous and rural residents in the Arctic is being impacted by climate change and in combination with globalization and resource development projected to increase significantly in the future (*high confidence*). [28.2.4]

The rapid rate at which climate is changing in the Polar Regions will impact natural and social systems (*high confidence*) and may exceed the rate at which some of their components can successfully adapt (*low to medium confidence*). [28.2.4; 28.4] The decline of Arctic sea-ice in summer is occurring at a rate that exceeds most model projections (*high confidence*) and evidence of similarly rapid rates of change is emerging in some regions of Antarctica. [IPCC WGI Chapter 14] In the future, trends in Polar Regions of populations of marine mammals, fish and birds will be a complex response to multiple stressors and indirect effects (*high confidence*). [28.3.2] Already, accelerated rates of change in permafrost thaw, loss of coastal sea ice, sea level rise and increased weather intensity are forcing relocation of some indigenous communities in Alaska (*high confidence*). [28.2.4.2; 28.2.5; 28.3.4]

Shifts in the timing and magnitude of seasonal biomass production could disrupt matched phenologies in the food webs, leading to decreased survival of dependent species (*medium confidence*). If the timing of primary and secondary production is no longer matched to the timing of spawning or egg release, survival could be impacted with cascading implications to higher trophic levels. This impact would be exacerbated if shifts in timing occur rapidly (*medium confidence*). [28.2.2; 28.3.2] Climate change will increase the vulnerability of terrestrial ecosystems to invasions by non-indigenous species, the majority likely to arrive through direct human assistance (*high confidence*).

Ocean acidification has the potential to inhibit embryo development and shell formation of some zooplankton and krill in the Polar Regions with potentially far-reaching consequences to food webs in these regions (*medium confidence*). Embryos of Antarctic krill have been shown to be vulnerable to increased concentrations of CO₂ in the water (*high confidence*). As well, there is increasing evidence that pelagic mollusks (pteropods) are vulnerable to ocean acidification (*medium confidence*). [28.2.2; 28.3.2]

There is increased evidence that climate change will have large effects on Arctic communities, especially where narrowly based economies leave a smaller range of adaptive choices. [28.2.6.1; 28.4] Some commercial activities will become more profitable while others will face decline. Increased economic opportunities are expected with increased navigability in the Arctic Ocean and the expansion of some land- and freshwater-based transportation networks. [28.2.6.1.3; 28.3.4.3] The informal, subsistence-based economy will be impacted (*high confidence*). There is *high confidence* that changing sea-ice conditions will result in more difficult access for hunting marine mammals. [28.2.6.1.6] Although Arctic residents have a history of adapting to change, the complex inter-linkages between societal, economic, and political factors and climatic stresses represent unprecedented challenges for northern communities, particularly as the rate of change will be faster than the social systems can adapt (*high confidence*). [28.4; 28.4.1; 28.2.5]

Impacts on the health and well-being of Arctic residents from climate change are significant and projected to increase – especially for many indigenous peoples (*high confidence*). [28.2.4] These are expected to vary among the diverse settlements, which range from small, remote predominantly indigenous communities to large cities and industrial settlements (*high confidence*), especially those located in highly vulnerable locations along ocean and river shorelines. [28.2.4]

28.1. Introduction

Several recent climate impact assessments on Polar Regions have been undertaken, including the synthesis report on Snow, Water, Ice and Permafrost in the Arctic (AMAP, 2011), the State of the Arctic Coast 2010 (2011) reports, the Antarctic Climate and the Environment (Turner *et al.*, 2009, 2013), Arctic Resilience Interim Report 2013 (2013), and the findings of the International Polar Year (IPY) (Krupnick *et al.*, 2011). These reports draw a consistent pattern of climate-driven environmental, societal and economic changes in the Polar Regions in recent decades. In this chapter, we use the scientific literature, including these reports, to consolidate the assessment of the impacts of climate change on Polar Regions from 2007, advance new scientific evidence of impacts and identify key gaps in knowledge on current and future impacts.

Previous IPCC reports define the Arctic as the area within the Arctic Circle (66°N), and the Antarctic as the continent with surrounding Southern Ocean south of the polar front, which is generally close to 58°S (IPCC, 2007). For the purpose of this report we use the conventional IPCC definitions as a basis, while incorporating a degree of flexibility when describing the Polar Regions in relation to particular subjects.

[INSERT FIGURE 28-1 HERE]

Figure 28-1: Location maps of the north and south polar regions. Credit: P. Fretwell, British Antarctic Survey.]

Changes in the physical and chemical environments of the Polar Regions are detailed in IPCC WG-1 report. The Arctic has been warming since the 1980s at approximately twice the global rate demonstrating the strongest temperature changes (~1 °C per decade) in winter and spring, and smallest in autumn. Sea ice declined at an average rate of 13% per decade; the Arctic Ocean is projected to become nearly ice-free in summer within this century. The duration of snow cover extent and snow depth are decreasing in North-America while increasing in Eurasia. Since late 1970s permafrost temperatures have increased between 0.5 to 2 °C. In the Southern Hemisphere, the strongest rates of atmospheric warming are occurring in the western Antarctic Peninsula (WAP, between 0.2 and 0.3 °C per decade) and the islands of the Scotia Arc, where there have also been increases in oceanic temperatures and large regional decreases in winter sea ice extent and duration. Warming, although less than WAP, has also occurred in the continental margins near to Bellingshausen Sea, Prydz Bay and the Ross Sea, with areas of cooling in between. Land regions have experienced glacial recession and changes in the ice and permafrost habitats in the coastal margins. The Southern Ocean continues to warm, with increased freshening at the surface due to precipitation leading to increased stratification. In both Polar Regions, surface waters will become seasonally corrosive to aragonite within decades, with some regions being affected sooner than others (Box CC-OA, WGI AR5 Chapter 6). Observations and models indicate that the carbon cycle of the Arctic and Southern Oceans will be impacted by climate change and increased CO₂.

28.2. Observed Changes and Vulnerability under Multiple Stressors

28.2.1. Hydrology and Freshwater Ecosystems

28.2.1.1. Arctic

Arctic rivers and lakes continue to show pronounced changes to their hydrology and ecology. Previously noted increases in Eurasian Arctic river flow (1936-1999; Peterson *et al.*, 2002) could not, for a similar period (1951-2000), be attributed with certainty to precipitation changes (Milliman *et al.*, 2008) but has been, including more recent extreme increases (2007), to enhanced poleward atmospheric moisture transport (Zhang *et al.*, 2013). By contrast, decreased flow in high-latitude Canadian rivers (1964-2000; average -10%) does match that for precipitation (Déry and Wood, 2005). Recent data (1977-2007) for 19 circumpolar rivers also indicates an area-weighted average increase of +9.8% (-17.1 to 47.0%; Overeem and Syvitski, 2010) accompanied by shifts in flow timing, with May snowmelt increasing (avg. 66%) but flow in the subsequent month of peak discharge decreasing (~7%). Across the Russian Arctic, dates of spring maximum discharge have also become earlier, particularly in the most recent [1960-2001] period analyzed (average -5d; range for 4 regions +0.2 to -7.1 d), but no consistent trend exists for magnitude (average -1%; range +21 to -24%; Shiklomanov *et al.*, 2007). Earlier timing was most

pronounced in eastern, colder continental climates where increases in air temperature have been identified as the dominant control (Tan *et al.*, 2011).

Increases have also occurred in winter low flows for many Eurasian and North American rivers (primarily late 20th century; Smith *et al.*, 2007; St. Jacques and Sauchyn, 2009; Walvoord and Striegl, 2007; Ye *et al.*, 2009), the key exceptions being decreases in eastern North America and unchanged flow in small basins of eastern Eurasia (Rennermalm *et al.*, 2010). Most of such studies suggest permafrost thaw (WGI, Ch.4) has increased winter flow, while others suggest increases in net winter precipitation minus evapotranspiration (Landerer *et al.*, 2010; Rawlins *et al.*, 2009a, b). Insufficient precipitation stations preclude deciphering the relative importance of these factors (WGI, Ch. 2.5.1).

The surface-water temperatures of large water bodies has warmed (1985-2009; Schneider and Hook, 2010), particularly for mid- and high latitudes of the northern hemisphere with spatial patterns generally matching those for air temperature. Where water bodies warmed more rapidly than air temperature, decreasing ice cover was suggested as enhancing radiative warming. Paleolimnological evidence indicates that highest primary productivity was associated with warm, ice-free summer conditions and the lowest with periods of perennial ice (Melles *et al.*, 2007). Increasing water temperatures affect planktonic and benthic biomass and lead to changes in species composition (Christoffersen *et al.* 2008; Heino *et al.* 2009, Jansson *et al.* 2010). Reduced ice cover with higher air temperatures and evaporation are responsible for the late 20th c – early 21st c desiccation of some Arctic ponds (Smol and Douglas, 2007).

Changes have occurred in the size and number of permafrost lakes over the last half-century (Hinkel *et al.*, 2007; Marsh *et al.*, 2009), but their patterns and rates of change are not consistent because of differing thawing states, variations in warming and effects of human activities (Hinkel *et al.* 2007; Prowse and Brown, 2010a). Thawing permafrost affects the biogeochemistry of water entering lakes and rivers (Frey and McClelland, 2009; Kokelj *et al.*, 2009) and their ecological structure and function (Lantz and Kokelj, 2008; Mesquita *et al.*, 2010; Thompson *et al.*, 2008), such as enhancing eutrophication by a shift from pelagic to benthic-dominated production (Thompson *et al.*, 2012).

The aquatic ecosystem health and biodiversity of northern deltas is dependent on combined changes in the elevation of spring river ice-jam floods and sea level (Lesack and marsh, 2007, 2010). Diminishing ice shelves (last half-century) have also caused a decline in the number of freshwater epishelf lakes that develop behind them (Veillette *et al.*, 2008; Vincent *et al.*, 2009). While such biophysical dependencies have been established, temporal trends in such river-delta and epishelf lake impacts and their linkages to changing climate remain to be quantified precisely.

An interplay of freshwater-marine conditions also affect the timing, growth, run size and distribution of several Arctic freshwater and anadromous fish. Key examples include: the timing of marine exit of Yukon River Chinook salmon (*Oncorhynchus tshawytscha*; 1961-2009) varied with air and sea temperatures and sea ice cover (Mundy and Evenson, 2011); the growth of young-of-year Arctic cisco (*Coregonus autumnalis*; 1978 – 2004) varied in response to lagged sea-ice concentration and Mackenzie River discharge, also indicating that decreased sea-ice concentration and increased river discharge enhanced marine primary production leading to more favorable foraging conditions (Von Biela *et al.* (2011); and factors that influence the water level and freshening of rivers, as well as the strength, duration and directions of prevailing coastal winds, affect survival of anadromous fishes during coastal migration and their subsequent run size (Fechhelm *et al.*, 2007).

28.2.1.2. Antarctic

Biota of Antarctic freshwater systems (lakes, ponds, short streams and seasonally wetted areas) are dominated by benthic microbial communities of cyanobacteria and green algae in a simple food web. Mosses occur in some continental lakes with higher plants absent. Planktonic ecosystems are typically depauperate and include small algae, bacteria and colourless flagellates, with few metazoans and no fish (Quesada and Velázquez, 2012). Recent compilations of single-year datasets have reinforced previous conclusions on the changing freshwater habitats in Antarctica (Verleyen *et al.*, 2012). In regions where the climate has warmed the physical impacts on aquatic

ecosystems include loss of ice and perennial snow cover, increasing periods of seasonal open water, increased water column temperatures and changes in water column stratification. In some areas a negative water balance has occurred due to increased temperature and changes in wind strength driving enhanced evaporation and sublimation and leading to increased salinity in lakes in recent decades (Hodgson *et al.*, 2006a). In other areas, especially glacial forelands, increased temperatures have led to greater volumes of seasonal meltwater in streams and lakes together with increased nutrient fluxes (*high confidence*). In both cases the balance between precipitation and evaporation can have detectable effects on lake ecosystems (*medium confidence*) through changes in water body volume and lake chemistry (Lyons *et al.*, 2006; Quesada *et al.*, 2006). Non-dilute lakes with a low lake depth to surface area ratio are most susceptible to inter-annual and inter-decadal variability in the water balance, as measured by changes in specific conductance (*high confidence*) (Verleyen *et al.*, 2012). Warming in the northwestern Antarctic Peninsula region has resulted in permafrost degradation in the last c. 50 yr impacting surface geomorphology and hydrology (Bockheim *et al.*, 2013) with the potential to increase soil biomass.

28.2.2. Oceanography and Marine Ecosystems

28.2.2.1. Arctic

28.2.2.1.1. Marine plankton, fish, and other invertebrates

Working Group I documents the expected physical and chemical changes that will occur in Arctic marine ecosystems (WGI AR5 Chapters 4, 6, and 11). In addition to climate change, naturally occurring interannual, decadal, and multi-decadal variations in climate will influence the Arctic Ocean and its neighboring high latitude seas (WGII Chapter 5). In recent years (2007-2012) ocean conditions in the Bering Sea have been cold (Stabeno *et al.*, 2012a), while the Barents Sea has been warm (Lind and Ingvaldsen, 2012).

In this section we build on previous reviews of observed species responses to climate (Wassman *et al.* 2011) to summarize the current evidence of the impact of physical and chemical changes in marine systems on the phenology, spatial distribution and production of Arctic marine species. For each type of response, the implications for phytoplankton, zooplankton, fish and shellfish are discussed. The implications of these changes on marine ecosystem structure and function will be the result of the synergistic effects of all three types of biological responses.

Phenological response

The timing of spring phytoplankton blooms is a function of seasonal light, hydrographic conditions, and the timing of sea ice breakup (Wassman, 2011). In addition to the open water phytoplankton bloom, potentially large ice algal blooms can form under the sea ice (Arrigo, 2012). During the period 1997-2009, a trend towards earlier phytoplankton blooms was detected in approximately 11% of the area of the Arctic Ocean (Kahru *et al.*, 2011). This advanced timing of annual phytoplankton blooms coincided with decreased sea ice concentration in early summer. Brown and Arrigo (2013) studied the timing and intensity of spring blooms in the Bering Sea from 1997-2010 and found that in northern regions, sea ice consistently retreated in late spring and was associated with ice-edge blooms, whereas, in the southern regions the timing of sea ice retreat varied, with ice-edge blooms associated with late ice retreat, and open water blooms associated with early ice retreat. Given the short time series and limited studies, there is *medium confidence* that climate variability and change has altered the timing and the duration of phytoplankton production.

The life cycles of calanoid copepods in the Arctic Ocean and Barents Sea are timed to utilize ice algal and phytoplankton blooms (Falk-Petersen *et al.* 2009; Søreide *et al.*, 2010; Darnis *et al.* 2012). Based on a synthesis of existing data, Hunt *et al.* (2011) hypothesized that in the southeastern Bering Sea, ocean conditions and the timing of sea ice retreat influences the species composition of dominant zooplankton, with lipid rich copepods being more abundant in cold years.

There is ample evidence that the timing of spawning and hatching of some fish and shellfish is aligned to match larval emergence with seasonal increases availability of prey (Gjosaeter *et al.*, 2009; Vikebø *et al.*, 2010; Bouchard and Fortier, 2011; Drinkwater *et al.*, 2011). These regional phenological adjustments to local conditions occurred over many generations (Ormseth and Norcross, 2009; Geffen *et al.*, 2011; Kristiansen *et al.*, 2011). There is *medium to high confidence* that climate induced disruptions in this synchrony can result in increased larval or juvenile mortality or changes in the condition factor of fish and shellfish species in the Arctic marine ecosystems.

Observed spatial shifts

Spatial heterogeneity in primary production has been observed (Lee *et al.*, 2010; Grebmeier, 2012). Simulation modeling studies show that spatial differences in the abundance of four species of copepod can be explained by regional differences in the duration of the growing season and temperature (Ji *et al.*, 2012). Retrospective studies based on surveys from 1952–2005 in the Barents Sea revealed that changes in the species composition, abundance and distribution of euphausiids were related to climate-related changes in oceanographic conditions (Zhukova *et al.*, 2009).

Retrospective analysis of observed shifts in the spatial distribution of fish and shellfish species along latitudinal and depth gradients showed observed spatial shifts were consistent with expected responses of species to climate change (Simpson *et al.*, 2011; Poloczanska *et al.* 2013; Box CC-MB, Chapter 30). Retrospective studies from the Bering Sea, Barents Sea, and the northeast Atlantic Ocean and Icelandic waters, showed that fish shift their spatial distribution in response to climate variability (i.e. interannual, decadal or multi-decadal changes in ocean temperature; Mueter and Litzow, 2008; Sundby and Nakken, 2008; Hátún *et al.*, 2009; Valdimarsson *et al.*, 2012; Kotwicki and Lauth, 2013). There are limits to the movement potential of some species. Vulnerability assessments indicate that the movement of some sub-arctic fish and shellfish species into the Arctic Ocean may be impeded by the presence of water temperatures on the shelves that fall below their thermal tolerances (Hunt *et al.*, 2013; Hollowed *et al.*, 2013). Coupled bio-physical models have reproduced the observed spatial dynamics of some the species in the Bering and Barents Seas, and are being used to explain the role of climate variability and change on the distribution and abundance of some species (Huse and Ellingsen, 2008; Parada *et al.*, 2010). In summary, there is *medium to high confidence* based on observations and modeling that some fish and shellfish have shifted their distribution in response to climate impacts on the spatial distribution and volume of suitable habitat.

Observed variations in production

Seasonal patterns in light, sea ice cover, freshwater input, stratification and nutrient exchange act in concert to produce temporal cycles of ice algal and phytoplankton production in Arctic marine ecosystems (Wassmann, 2011; Perrette *et al.*, 2011; Tremblay *et al.*, 2012). Satellite observations and model estimates for the period 1988–2007 showed that phytoplankton productivity increased in the Arctic Ocean in response to a downward trend in the extent of summer sea ice (Zhang *et al.*, 2010). Satellite data provided evidence of a 20% increase in annual net primary production in the Arctic Ocean between 1998 and 2009 in response to extended ice free periods (Arrigo and van Dijken, 2011). Regional trends in primary production will differ in response to the amount of open water area in summer (Arrigo and van Dijken, 2011). Other studies showed gross primary production increased with increasing air temperature in the Arctic Basin and Eurasian shelves (Slagstad *et al.*, 2011). A recent 5 year study (2004–2008) in the Canadian Basin showed that smaller phytoplankton densities were higher than larger phytoplankton densities in years when sea surface temperatures were warmer, the water column was more stratified, and nutrients were more depleted during the Arctic summer (Li *et al.*, 2009; Morán *et al.*, 2010). Additional observations will help to resolve observed differences between in-situ and satellite derived estimates of primary production (Matrai *et al.*, 2013). In conclusion, based on recent observations and modeling, there is *medium to high confidence* that primary production has increased in the Arctic Ocean in response to changes in climate and its impact on the duration and areal extent of ice free periods in summer.

Regional differences in zooplankton production have been observed. During a period of ocean warming (1984–2010), Dalpadado *et al.* (2012) observed an increase in the biomass of lipid rich euphausiids in the Barents Sea and

relatively stable levels of biomass and production of *Calanus finmarchicus*. In the Bering Sea, observations over the most recent decade in the southeast Bering Sea showed *C. marshallae* were more abundant in cold years than warm years (Coyle *et al.*, 2011).

There is strong evidence that climate variability impacts the year-class strength of Arctic marine fish and shellfish through its influence on: predation risk; the quality, quantity and availability of prey; and reproductive success (Mueter *et al.*, 2007; Bakun 2010; Drinkwater *et al.*, 2010). Regional differences in the species responses to climate change will be a function of the exposure of the species to changing environmental conditions, the sensitivity of the species to these changes (Beaugrand and Kirby, 2010) and the abilities of species to adapt to changing conditions (Pörtner and Peck, 2010; Donelson *et al.*, 2011). There is *high confidence* that shifts in ocean conditions have impacted the abundance of fish and shellfish in Arctic regions. Observed trends in the abundance of commercial fish and shellfish may also be influenced by historical patterns of exploitation (Vert-pre *et al.*, 2013).

28.2.2.1.2. Marine mammals, polar bear, and sea birds

Studies on responses of Arctic and subarctic marine mammals to climate change are limited and vary according to insight into their habitat requirements and trophic relationships (Laidre *et al.*, 2008). Many Arctic and subarctic marine mammals are highly specialized, have long-life spans, and are poorly adapted to rapid environmental change (Moore and Huntington, 2008), and changes may be delayed until significant sea ice loss has occurred (Freitas *et al.*, 2008; Laidre *et al.*, 2008).

Climate change effects on Arctic and subarctic marine mammal species will vary by life history, distribution, and habitat specificity (*high confidence*). Climate change will improve conditions for a few species, have minor negative effects for others, and some will suffer major negative effects (Ragen *et al.*, 2008; Laidre *et al.*, 2008). Climate change resilience will vary and some ice-obligate species should survive in regions with sufficient ice and some may adapt to ice-free conditions (Moore and Huntington, 2008). Less ice-dependent species may be more adaptable but an increase in seasonally migrant species could increase competition (Moore and Huntington, 2008).

Climate change vulnerability was associated with feeding specialization, ice dependence, and ice reliance for prey access and predator avoidance (Laidre *et al.*, 2008). There is *medium agreement* on which species' life histories are most vulnerable. Hooded seals (*Cystophora cristata*) and narwhal (*Monodon monoceros*) were identified as most at risk and ringed seals (*Pusa hispida*) and bearded seals (*Erignathus barbatus*) as least sensitive (Laidre *et al.*, 2008). Kovacs *et al.* (2010) shared concern for hooded seals and narwhal but had concerns for ringed seals and bearded seals. Narwhal may have limited ability to respond to habitat alteration (Williams *et al.* 2011). Species that spend only part of the year in the Arctic (e.g., gray whale (*Eschrichtius robustus*), killer whale (*Orcinus orca*)) may benefit from reduced ice (Moore, 2008; Laidre *et al.*, 2008; Higdson and Ferguson, 2009; Matthews *et al.*, 2011; Ferguson *et al.*, 2012). Killer whale expansion into the Arctic could cause a trophic cascade (Higdson and Ferguson, 2009) although there is *limited evidence* at this time.

There is *limited evidence* although *medium agreement* that generalists and pelagic feeding species may benefit from increased marine productivity from reduced ice while benthic feeding species near continental shelf habitats may do poorly (Bluhm and Gradinger, 2008). There is *limited evidence* but *high agreement* that dietary or habitat specialists will do poorly with reduced ice. Reduction of summer/autumn ice was the primary extrinsic factor affecting Pacific walrus (*Odobenus rosmarus*) with predictions of distribution changes, reduced calf recruitment, and longer-term predictions of high extinction probability (Cooper *et al.*, 2006; MacCracken, 2012). Summer ice retreat may make migration to such habitats energetically unprofitable for ringed seals (Freitas *et al.*, 2008). Ice loss threatens Baltic ringed seals (Kovacs and Lydersen, 2008). In Hudson Bay, earlier spring break-up and changes in snow cover over lairs have reduced ringed seal recruitment (Ferguson *et al.*, 2005). Changes in snowfall over the 21st century were projected to reduce ringed seal habitat for lairs by 70% (Hezel *et al.*, 2012). Similarly, harp seal (*Pagophilus groenlandicus*) breeding habitat was affected by changing ice conditions that could reduce pup survival (Bajzak *et al.*, 2011). While there is *limited evidence*, there are concerns that climate change may cause indirect effects on Arctic marine mammals' health (e.g., pathogen transmission, food web changes, toxic chemical exposure, shipping, and development) (Burek *et al.*, 2008).

Empirical studies provide direct insight into the mechanisms of climate change impact on polar bears (*Ursus maritimus*) but modelling allows predictive capacity (Hunter *et al.*, 2010; Amstrup *et al.*, 2010; Durner *et al.*, 2011; Castro de la Guardia *et al.*, 2013).

Polar bears are highly specialized and use annual ice over the continental shelves as their preferred habitat (Durner *et al.*, 2009; Miller *et al.*, 2012). The recent and projected loss of annual ice over continental shelves, decreased ice duration, decreased ice thickness, and habitat fragmentation is causing reduced food intake, increased energy expenditure, and increased fasting in polar bears (*high confidence*) (Stirling and Parkinson, 2006; Regehr *et al.*, 2007; Durner *et al.*, 2009; Amstrup *et al.*, 2010; Hunter *et al.*, 2010; Derocher *et al.*, 2011; Rode *et al.*, 2012; Sahanatien and Derocher, 2012; Castro de la Guardia *et al.*, 2013).

Subpopulation response varies geographically. Only 2 of the 19 subpopulations, Western Hudson Bay (Regehr *et al.*, 2007) and the Southern Beaufort Sea (Regehr *et al.*, 2010; Rode *et al.*, 2010a) have data series adequate for clear identification of abundance effects related to climate change. Many other subpopulations show characteristics associated with decline but some remain stable. Declining ice is causing lower body condition, reduced individual growth rates, lower fasting endurance, lower reproductive rates, and lower survival (*high confidence*) (Regehr *et al.*, 2007; Regehr *et al.*, 2010; Rode *et al.*, 2010a; Molnar *et al.*, 2011; Rode *et al.*, 2012). Condition is a precursor to demographic change (*very high confidence*) (Regehr *et al.*, 2010; Hunter *et al.*, 2010; Rode *et al.*, 2010a; Robinson *et al.*, 2011). The decline in the subpopulation in Western Hudson Bay by 21% between 1987 and 2004 was related to climate change (*medium confidence*) (Regehr *et al.*, 2007).

Replacement of multiyear ice by annual ice could increase polar bear habitat (*low confidence*) (Derocher *et al.*, 2004). Increasing the distance to multiyear ice and terrestrial refugia at maximal melt may result in drowning, cub mortality, and increased energetic costs (Monnett and Gleason, 2006; Durner *et al.*, 2011; Pagano *et al.*, 2012).

There is *robust evidence* of changes in sea ice conditions changing polar bear distribution including den areas (*high confidence*) (Fischbach *et al.*, 2007; Schliebe *et al.*, 2008; Gleason and Rode, 2009; Towns *et al.*, 2010; Derocher *et al.*, 2011). The number of human-bear interactions are projected to increase with warming (*high confidence*) (Stirling and Parkinson, 2006; Towns *et al.*, 2009).

Use of terrestrial resources by polar bears was suggested as adaptive (Dyck *et al.*, 2007; Dyck and Romberg, 2007; Armstrong *et al.*, 2008; Dyck *et al.*, 2008; Dyck and Kebreab, 2009; Rockwell and Gormezano, 2009; Smith *et al.*, 2010). Polar bears cannot adapt to terrestrial foods (Stirling *et al.*, 2008b; Amstrup *et al.*, 2009; Slater *et al.*, 2010; Rode *et al.*, 2010b), and will *most likely* not be able to adapt to climate change and reduced sea ice extent (*very high confidence*). Changing ice conditions are linked to cannibalism (Amstrup *et al.*, 2006), altered feeding (Cherry *et al.*, 2009), unusual hunting behaviour (Stirling *et al.*, 2008a), and diet change (Iverson *et al.*, 2006; Thiemann *et al.*, 2008) (*medium confidence*).

Upwelling or subsurface convergence areas found in frontal zones and eddies, and the marginal ice zone, are associated with high marine productivity important to Arctic seabirds (e.g. Irons *et al.*, 2008). Long-term or permanent shifts in convergence areas and the marginal ice-edge zone induced by climate change may cause mismatch between the timing of breeding and the peak in food availability and, thus, potentially have strong negative impacts on seabird populations (Gaston *et al.*, 2005; Moline *et al.*, 2008; Gaston *et al.*, 2009; Grémillet and Boulinier, 2009) (*medium confidence*).

The contrasting results from the relatively few studies of impacts of climate change on Arctic seabirds, demonstrate that future impacts will be highly variable between species and between populations of the same species (*medium confidence*). Retreating sea ice and increasing SSTs have favored some species and disadvantaged others (Gaston *et al.* 2005; Byrd *et al.* 2008; Irons *et al.* 2008; Karnovsky *et al.* 2010; Fredriksen *et al.* 2013). Some species of sea birds respond to a wide range of sea surface temperatures via plasticity of their foraging behaviour, allowing them to maintain their fitness levels (Grémillet *et al.* 2012). Phenological changes and changes in productivity of some breeding colonies have been observed (Byrd *et al.* 2008; Gaston and Woo 2008; Moe *et al.* 2009). Negative trends in population size, observed over the last few decades for several species of widespread Arctic seabirds, may be

related to over-harvesting and pollution as well as climate change effects (Gaston, 2011). For those species whose distribution is limited by sea ice and cold water, polar warming could be beneficial (Mehlum 2012).

A major ecosystem shift in the Northern Bering Sea starting in the mid 1990s caused by increased temperatures and reduced sea ice cover had a negative impact on benthic prey for diving birds and these populations have declined in the area (Grebmeier *et al.*, 2006). More recently, the Bering Sea has turned colder again.

28.2.2.2. Antarctica

Productivity and food web dynamics in the Southern Ocean are dominated by the extreme seasonal fluctuations of irradiance and the dynamics of sea ice, along with temperature, carbonate chemistry and vertical mixing (Massom and Stammerjohn, 2010; Boyd *et al.*, 2012; Murphy *et al.*, 2012a). Moreover, there is large-scale regional variability in habitats (Grant *et al.*, 2006) and their responses to climate change (WGI). Antarctic krill, *Euphausia superba* (hereafter, 'krill') is the dominant consumer, eating diatoms, and, in turn, is the main prey of fish, squid, marine mammals and seabirds. Krill is dominant from the Bellingshausen Sea east through to the Weddell Sea and the Atlantic sector of the Southern Ocean (Rogers *et al.*, 2012). In the East Indian and southwest Pacific sectors of the Southern Ocean, the krill-dominated system lies to the south of the Southern Boundary of the Antarctic Circumpolar Current (Nicol *et al.*, 2000a,b) while to the north copepods and myctophid fish are most important (Rogers *et al.*, 2012). Further west, where the Weddell Sea exerts an influence, krill are found as far north as the Subantarctic Circumpolar Current Front (Jarvis *et al.*, 2010). Where sea ice dominates for most of the year, ice-obligate species (e.g. *Euphausia crystallorophias* and *Peluragramma antarcticum*) are most important (Smith *et al.*, 2007).

Few studies were available in AR4 to document and validate the changes in these systems resulting from climate change. Those studies reported increasing abundance of benthic sponges and their predators, declining populations of krill, Adélie and emperor penguins, and Weddell seals, and a possible increase in salps, noting some regional differences in these trends. The importance of climate processes in generating these changes could not be distinguished from the indirect consequences of the recovery of whale and seal populations from past over-exploitation (Trathan and Reid, 2009; Murphy *et al.*, 2012a,b).

28.2.2.2.1. Marine plankton, krill, fish, and other invertebrates

Distributions of phytoplankton and zooplankton have moved south with the frontal systems (Hinz *et al.*, 2012; Mackey *et al.*, 2012), including range expansion into the Southern Ocean from the north by the coccolithophorid, *Emiliania huxleyi* (Cubillos *et al.*, 2007), and the red-tide dinoflagellate *Noctiluca scintillans* (McLeod *et al.*, 2012) (*medium confidence*). There is *insufficient evidence* to determine whether other range shifts are occurring.

Collapsing ice shelves are altering the dynamics of benthic assemblages by exposing areas previously covered by ice shelves, allowing increased primary production and establishment of new assemblages (e.g. collapse of the Larson A/B ice shelves) (Peck *et al.*, 2009; Gutt *et al.*, 2011) (*medium confidence*). More icebergs are grounding, causing changes in local oceanography and declining productivity that consequently affects productivity of benthic assemblages (Thrush and Cummings 2011) (*low confidence*). Iceberg scour on shallow banks is also increasing, disrupting resident benthic assemblages (Barnes and Souster 2011; Gutt *et al.*, 2011) (*medium confidence*).

Primary production is changing regionally in response to changes in sea ice, glacial melt and oceanographic features (Arrigo *et al.*, 2008; Boyd *et al.*, 2012) (*medium confidence*). Off the west Antarctic Peninsula, phytoplankton stocks and productivity have decreased north of 63°S, but increased south of 63°S (Montes-Hugo *et al.*, 2009) (WGII Chapter 6) (*high confidence*). This study (based on time-series of satellite-derived and measured chlorophyll concentrations) also indicated a change from diatom-dominated assemblages to ones dominated by smaller phytoplankton (Montes-Hugo *et al.*, 2009). The reduced productivity in the north may be tempered by increased inputs of iron through changes to ocean processes in the region (Dinniman *et al.*, 2012) (*low confidence*).

Since the 1980s, Antarctic krill densities have declined in the Scotia Sea (Atkinson *et al.*, 2004), in parallel with regional declines in the extent and duration of winter sea ice (Flores *et al.*, 2012). Uncertainty remains over changes in the krill population because this decline was observed using net samples and is not reflected in acoustic abundance time series (Nicol and Brierley 2010); the observed changes in krill density may have been partly a result of changes in distribution (Murphy *et al.* 2007). Nevertheless, given its dependence on sea ice (Nicol *et al.*, 2008) the krill population may already have changed and will be subject to further alterations (*high confidence*).

The response of krill populations is probably a complex response to multiple stressors. Decreases in recruitment of post-larval krill across the Scotia Sea have been linked to declines in sea-ice extent in the Antarctic Peninsula region (Wiedenmann *et al.*, 2009) (*medium confidence*) but these declines may have been offset by increased growth arising from increased water temperature in that area (Wiedenmann *et al.*, 2008). However, near South Georgia krill productivity may have declined as a result of the increased metabolic costs of increasing temperatures (Hill *et al.*, 2013) (*low confidence*). The combined effects of changing sea ice, temperature and food have not been investigated.

28.2.2.2.2. *Marine mammals and sea birds*

In general, many Southern Ocean seals and seabirds exhibit strong relationships to a variety of climate indices, and many of these relationships are negative to warmer conditions (Trathan *et al.*, 2007; Barbraud *et al.*, 2012; Forcada *et al.*, 2012) (*low confidence*). Regional variations in climate change impacts on habitats and food will result in a mix of direct and indirect effects on these species. For example, Adélie penguin colonies are declining in recent decades throughout the Antarctic Peninsula while the reduction in chinstrap penguins is more regional (Lynch *et al.*, 2012) and related to reductions in krill availability (Lima and Estay, 2013). In contrast gentoo penguins are increasing in that region and expanding south (Lynch *et al.*, 2012) (*high confidence*). This may be explained by the reduced sea ice habitats and krill availability in the north resulting in a southward shift of krill predators, particularly those dependent on sea ice (Forcada *et al.*, 2012) and the replacement of these predators in the north by species that do not depend on sea ice, such as gentoo penguins and elephant seals (Costa *et al.*, 2010; Trivelpiece *et al.*, 2011; Ducklow *et al.*, 2012; Murphy *et al.*, 2013) (*low confidence*). A contrasting situation is in the Ross Sea where Adélie penguin populations have increased (Smith *et al.*, 2012). The mechanisms driving these changes are currently under review and may be more than simply sea ice (Lynch *et al.*, 2012; Melbourne-Thomas *et al.*, 2013). For example, too much or too little sea ice may have negative effects on the demography of Adélie and emperor penguins (see Barbraud *et al.*, 2012 for review). Also, increased snow precipitation which accumulates in breeding colonies can decrease survival of chicks of Adélie penguins when accompanied by reduced food supply (Chapman *et al.*, 2011).

Changes elsewhere are less well known. Some emperor penguin colonies have decreased in recent decades (Barbraud *et al.*, 2008; Jenouvrier *et al.*, 2009) (*low confidence*) and one breeding site has been recorded as having been vacated (Trathan *et al.*, 2011). However, there is *insufficient evidence* to make a global assessment of their current trend. In the subantarctic of the Indian sector, reductions in seal and seabird populations may indicate a region-wide shift to a system with lower productivity (Weimerskirch *et al.*, 2003; Jenouvrier *et al.*, 2005a, b) (*low confidence*) but commercial fishing activities may also play a role.

Where frontal systems are shifting south, productive foraging areas also move to higher latitudes. In the Indian sector, this is thought to be causing declines in king penguin colonies on subantarctic islands (Péron *et al.*, 2010) (*low confidence*), while the shift in wind patterns may be causing changes to the demography of albatross (Weimerskirch *et al.*, 2012) (*low confidence*).

As identified in the last assessment, some species' populations may suffer as a result of fisheries while others are recovering from past over-exploitation, either of which may confound interpretation of the response of these species and their food webs to climate change. The recovery of Antarctic fur seals on some subantarctic islands has been well documented and their populations may now be competing with krill-eating macaroni penguins (Trathan *et al.*, 2012). More recently, there has been confirmation that populations of some Antarctic whales are recovering, such as humpbacks (Nicol *et al.*, 2008; Zerbini *et al.*, 2010), suggesting that food is currently not limiting. In contrast, a number of albatross and petrel populations are declining as a result of incidental mortality in longline fisheries in southern and temperate waters where these birds forage (Croxall *et al.*, 2012).

28.2.3. Terrestrial Ecosystems

28.2.3.1. Arctic

Arctic terrestrial ecosystems have undergone dramatic changes throughout the late Pleistocene and Holocene (last 130 000 years) mainly driven by natural climate change. Significant altitudinal and latitudinal advances and retreats in tree line have been common, animal species have gone extinct, and animal populations have fluctuated significantly throughout this period e.g. (Lorenzen *et al.*, 2011; Salonen *et al.*, 2011; Mamet and Kershaw, 2012).

28.2.3.1.1. Phenology

Phenological responses attributable to warming are apparent in most Arctic terrestrial ecosystems (*medium confidence*). They vary from earlier onset and later end of season in western Arctic Russia (Zeng *et al.*, 2013), to little overall trend in plant phenology in the Swedish sub Arctic (Callaghan *et al.*, 2010), to dramatic earlier onset of phenophases in Greenland (Høye *et al.*, 2007; Post *et al.*, 2009a; Callaghan *et al.*, 2011a) (Figure 28-2).

[INSERT FIGURE 28-2 HERE

Figure 28-2: Temporal change in onset of flowering (plants), median date of emergence (arthropods) and clutch initiation dates (birds) in high-Arctic Greenland. Red dots are statistically significant, blue dots are not. Source: Høye *et al.*, 2007.]

28.2.3.1.2. Vegetation

The latest assessment of changes in NDVI (Normalized Difference Vegetation Index, a proxy for plant productivity) from satellite observations between 1982 and 2012 shows that about a third of the Pan-Arctic has substantially greened, <4% browned and > 57% did not change significantly (Xu *et al.*, 2013) (Figure 28-3). The greatest increases reported in recent years were in the North American high Arctic, along the Beaufort Sea and the east European Arctic (Zhang *et al.*, 2008; Pouliot *et al.*, 2009; Bhatt *et al.*, 2010; Forbes *et al.*, 2010; Walker *et al.*, 2011; Epstein *et al.*, 2012; Macias-Fauria *et al.*, 2012; Xu *et al.*, 2013).

[INSERT FIGURE 28-3 HERE

Figure 28-3: Significant changes ($p < 0.01$) in photosynthetically active period NDVI between 1982 and 2012. Source: Xu *et al.*, 2013.]

The positive trends in NDVI are associated with increases in the summer warmth index (sum of the monthly-mean temperatures above freezing expressed as °C per month) that have increased on average by 5°C per month for the Arctic as a whole (Xu *et al.*, 2013). However, the even greater 10 to 12°C per month increase for the land adjacent to the Chukchi and Bering Seas (Figure 28-3) was associated with decreases in NDVI. On the Yamal Peninsula in Russia the pattern of NDVI is partly due to surface disturbance, such as landslide activity (Walker *et al.*, 2009). Small rodent cycles reduce NDVI in sub Arctic Sweden, by decreasing biomass and changing plant species composition (Olofsson *et al.*, 2012). The changing NDVI signal should therefore generally be interpreted with care.

In common with treeline trees and herbs, the abundance and biomass of deciduous shrubs and graminoids (grasses and grass-like plants) have increased substantially in certain parts of the Arctic tundra in recent years, but remained stable or decreased in others (*very high confidence*). Attribution for the increases and decreases in deciduous shrubs and graminoids is heterogeneous with drivers varying among different regions (*very likely*), including Arctic warming, differences in herbivory, industrial development, legacies from past land use, and changes in moisture (Post and Pedersen 2008; Olofsson *et al.*, 2009; Forbes *et al.*, 2009 and 2010; Kitti *et al.*, 2009; Kumpula *et al.*, 2011 and 2012; Myers-Smith *et al.*, 2011; Elmendorf *et al.*, 2012b; Callaghan *et al.*, 2011b and 2013; Gamon *et al.*, 2013).

Shrubs have generally expanded their ranges and/or growth over the last 20 years (Danby and Hik, 2007; Hudson and Henry, 2009; Forbes *et al.*, 2010; Hallinger *et al.*, 2010; Rundqvist *et al.*, 2011; Hedenås *et al.*, 2011; Hill and Henry, 2011; Myers-Smith *et al.*, 2011a,b; Callaghan *et al.*, 2011b; Elmendorf *et al.*, 2012a,b; Macias-Fauria *et al.*, 2012), and have varied from dramatic, i.e. 200% area increase in study plots (Rundqvist *et al.*, 2011) in sub arctic Sweden to early invasion of a fell field community on west Greenland by low shrubs (Callaghan *et al.*, 2011a).

A synthesis (61 sites: (Elmendorf *et al.*, 2012a) of experimental warming studies of up to 20 years duration in tundra sites worldwide, showed, overall, increased growth of deciduous shrubs and graminoids, decreased cover of mosses and lichens, and decreased species diversity and evenness. Elmendorf *et al.*, (2012a) point out that the groups that increased most in abundance under simulated warming were graminoids in cold regions and primarily shrubs in warm regions of the tundra. However, strong heterogeneity in responses to the experimental warming suggested that other factors could moderate the effects of climate warming significantly like herbivory, differences in soil nutrients and pH, precipitation, winter temperatures and snow cover, and species composition and density.

Snow bed habitats have decreased in sub arctic Sweden (Björk and Molau, 2007; Hedenås *et al.*, 2011). In other plant communities, changes have been less dramatic, ranging from small increases in species richness in the south west Yukon of the Canadian sub Arctic (Danby *et al.*, 2011), through subtle changes in plant community composition in west and southeast Greenland (Daniëls and De Molenaar, 2011; Callaghan *et al.*, 2011a) to 70 year stability of a plant community on Svalbard (Prach *et al.*, 2010).

The responses to Arctic warming of lichen and bryophyte (mosses) diversity have been heterogenous, varying from consistent negative effects to significant increases in recent years (Hudson and Henry, 2009; Tømmervik *et al.*, 2009; Tømmervik *et al.*, 2012). Forbes and Kumpula (2009) recorded long-term and widespread lichen degradation in northern Finland attributed more to trampling of dry lichens by reindeer in summer than winter consumption as forage.

Palaeorecords of vegetation change indicate that the northern tree line should extend upwards and northwards during current climate warming (Callaghan *et al.*, 2005) because tree line is related to summer warmth (e.g. Harsch *et al.*, 2009). Although the tree line has moved northwards and upwards in many Arctic areas, it has not shown a general circumpolar expansion in recent decades (*high confidence*).

Model projections that suggest a displacement of between 11 and 50% of tundra by forest by 2100 (see references in Callaghan *et al.*, 2005) and shifts upslope by 2 to 6 m per year (Moen *et al.*, 2004) and northwards by 7.4–20 km per year (Kaplan and New, 2006) might be overestimating rate of tree line advance by a factor of up to 2000 (Van Bogaert *et al.*, 2011). The fastest upslope shifts of tree lines recorded during 20th century warming are 1 to 2 m per year (Shiyatov *et al.*, 2007; Kullman and Öberg, 2009) whereas the fastest so-far recorded northward-migrating tree line replaces tundra by taiga at a rate of 3–10 m per year (Kharuk *et al.*, 2006). In some areas, the location of the tree line has not changed or has changed very slowly (Payette, 2007; MacDonald *et al.*, 2008a). A global study by Harsch *et al.* (2009) showed that only 52% of 166 global tree line sites studied had advanced over the past 100 years. In many cases the tree line has even retreated (Cherosov *et al.*, 2010). At the small scale, the tree line has shown increase, decrease and stability in neighboring locations (Van Bogaert *et al.*, 2011; Lloyd *et al.*, 2011).

Evidence for densification of the forest at the sub Arctic tree line is robust and consistent within Fennoscandia (Tømmervik *et al.*, 2009; Rundqvist *et al.*, 2011; Hedenås *et al.*, 2011) and Canada (Danby and Hik, 2007). Dendroecological studies indicate enhanced conifer recruitment during the twentieth century in the northern Siberian taiga (Briffa *et al.*, 2008). Some of the changes are dramatic, such as an increase in area of mountain birch in study plots in northern Sweden by 600% between 1977/8 and 2009/10 (Rundqvist *et al.*, 2011) and a doubling of tree biomass in Finnmarksvidda in northern Norway since 1957 (Tømmervik *et al.*, 2009). However, model projections of displacement of deciduous forest by evergreen forest (Wolf *et al.*, 2008; Wramneby *et al.*, 2010) have not so far been validated.

Where the mountain birch tree line has increased in elevation and shrub (e.g. willow, dwarf birch) abundance has increased, the response can be an interaction between climate warming, herbivory pressure and earlier land use

(Olofsson *et al.*, 2009; Hofgaard *et al.*, 2010; Van Bogaert *et al.*, 2011). In Fennoscandia and Greenland, heavy grazing by large herbivores may significantly check deciduous low erect shrub (e.g. dwarf shrub and willow) growth (Post *et al.*, 2008; Kitti *et al.*, 2009; Olofsson *et al.*, 2009).

Less moisture from snow and more rain now favors broadleaf trees over conifers and mosses in some areas (Juday, 2009) while moisture deficits are reducing the growth of some northern forests (Goetz *et al.*, 2005; Verbyla, 2008; Yarie, 2008) and making them more susceptible to insect pest outbreaks (see references in (Callaghan *et al.*, 2011c). Death of trees through drought stress or insect pest activity will increase the probability of fire that will have positive feedbacks (increase warming) on the climate (Mack *et al.*, 2011).

28.2.3.1.3. *Changes in animal populations*

The documented collapse or dampening of population cycles of voles and lemmings over the last 20-30 years in parts of Fennoscandia and Greenland (Schmidt *et al.*, 2012), can be attributed with *high confidence* to climate change (Ims *et al.*, 2007; Gilg *et al.*, 2009; Ims *et al.*, 2011; Kausrud *et al.*, 2009). A shortening of the snow season and more thaw and/or rain events during the winter will influence on the subnivean space which provide thermal insulation, access to food, and protection from predators (Berg *et al.*, 2008; Johansson *et al.*, 2011; Kausrud *et al.*, 2009). However, the causes of the changes in the lemming and vole cycles are still being debated as factors other than climate change may also be of importance (Brommer *et al.*, 2010; Krebs, 2011).

Climate-mediated range expansion both in altitude and latitude of insect pests, and increased survival due to higher winter temperatures, has been documented for bark beetles in North America (Robertson *et al.*, 2009) and for geometrid moths in Fennoscandia (Jepsen *et al.*, 2008; Callaghan *et al.*, 2010; Jepsen *et al.*, 2011), causing more extensive forest damage than before. Outbreaks of insect pests like geometrid moths can even reduce the strengths of CO₂ sinks in some areas (Heliasz *et al.*, 2011).

The decline in wild reindeer and caribou (both *Rangifer tarandus*) populations in some regions of about 30 percent over the last 10-15 years has been linked both to climate warming and anthropogenic landscape changes (Post *et al.*, 2009a; Vors and Boyce 2009; Russell and Gunn 2010). Even though most of the Arctic has warmed, the decline in the populations has not been uniform. Some of the North American large, wild herds have for example declined by 75-90 percent, while other wild herds and semi-domestic herds in Fennoscandia and Russia have been stable or even increased (Gunn *et al.*, 2009; Vors and Boyce, 2009; Joly *et al.*, 2011; Forbes *et al.*, 2009; Forbes 2010; Kumpula *et al.*, 2012).

The expected and partially observed increased primary productivity of Arctic tundra may potentially increase the supply of food for Arctic ungulates. However, the overall quality of forage may decline during warming, for example if the nitrogen content of key fodder species for ungulates were to drop during warming (Turunen *et al.*, 2009; Heggberget *et al.*, 2010), while lichen biomass, an important winter fodder for reindeer, is decreasing over parts of the Arctic region. Herbivory also changes the vegetation itself in concert with the warming, further complicating the prediction of vegetation changes and their impacts on ungulate populations (van Der Wal *et al.*, 2007; Turunen *et al.*, 2009).

More frequent rain-on-snow icing events and thicker snow-packs caused by warmer winters and increased precipitation may restrict access to vegetation and may have profound negative influences on the population dynamics of Arctic ungulates (Berg *et al.*, 2008; Forchhammer *et al.*, 2008; Miller and Barry, 2009; Stien *et al.*, 2010; Hansen *et al.*, 2011; Stien *et al.*, 2012). Such events have caused heavy mortality in some semi-domestic reindeer herds and musk oxen in recent years (Grenfell and Putkonen, 2008; Forbes, 2009; Bartsch *et al.*, 2010), and have also been shown to synchronize the dynamics of a resident vertebrate community (small mammals, reindeer and Arctic fox) in Svalbard (Hansen *et al.*, 2013). In contrast, Tyler *et al.* (2008) and Tyler (2010) suggested that generally warmer winters enhance the abundance of reindeer populations.

It has been suggested that warming-induced trophic mismatches between forage availability and quality and timing of calving have a role in the decline of circumpolar reindeer and caribou populations (Post and Forchhammer, 2008; Post *et al.*, 2009a,b), although such trophic mismatch has been disputed (Griffith *et al.*, 2010).

Adjustment via phenotypic plasticity instead of adaptation by natural selection is expected to dominate vertebrate responses to rapid Arctic climate change, and many such adjustments have already been documented (Gilg *et al.*, 2012).

28.2.3.1.4. Long-term trends and event-driven changes

Long-term climate change impacts on vegetation and animal populations are accelerated when tipping points are triggered by events such as extreme weather, fire, insect pest and disease outbreaks. The impacts of winter thaw events on ecosystems are now well-documented e.g. (Bokhorst *et al.*, 2011) but studies of the severe impacts of tundra fires on vegetation and biospheric feedbacks are recent (Mack *et al.*, 2011). Results from experimental winter thaws were validated by a natural event in northern Norway and Sweden in 2007 that reduced NDVI by almost 30% over at least 1400 km² (Bokhorst *et al.*, 2009). Studies on relationships between climate change and plant disease are rare but Olofsson *et al.*, (2011) showed that increased snow accumulation led to a higher incidence of fungal growth on sub Arctic vegetation.

28.2.3.2. Antarctica

Antarctic terrestrial ecosystems occur in 15 biologically-distinct areas (Terauds *et al.*, 2012) with those in the maritime and sub-Antarctic islands experiencing the warmest temperatures, reduced extreme seasonality and greatest biodiversity (Convey, 2006). In the cooler conditions on the continent, species must be capable of exploiting the short periods where temperature and moisture availability are above physiological and biochemical thresholds. In many areas there is no visible vegetation, with life being limited, at the extreme, to endolithic (within rock) communities of algae, cyanobacteria, fungi, bacteria and lichens (Convey, 2006).

Few robust studies are available of biological responses to observed climatic changes in natural Antarctic terrestrial ecosystems. The rapid population expansion and local-scale colonisation by two native flowering plants (*Deschampsia antarctica* and *Colobanthus quitensis*) in maritime Antarctica (Parnikoza *et al.*, 2009) remains the only published repeat long-term monitoring study of any terrestrial vegetation or location in Antarctica. Radiocarbon dating of moss peat deposits has shown that growth rates and microbial productivity have risen rapidly on the Antarctic Peninsula since the 1960s, consistent with temperature changes, and are unprecedented in the last 150 years (Royles *et al.*, 2013). In east Antarctica moss growth rates over the last 50 yrs have been linked to changes in wind speed and temperature and their influence on water availability (Clarke *et al.*, 2012). A contributing factor is that air temperatures have increased past the critical temperature at which successful sexual reproduction (seed set) can now take place, changing the dominant mode of reproduction, and increasing the potential distance for dispersal (Convey, 2011) (*low confidence*). Similar changes in the local distribution and development of typical cryptogamic vegetation of this region have been reported (Convey, 2011), including the rapid colonisation of ice free ground made available through glacial retreat and reduction in extent of previously permanent snow cover (Olech and Chwedorzewska, 2011). As these vegetation changes create new habitat, there are concurrent changes in the local distribution and abundance of the invertebrate fauna that then colonise them (*low confidence*).

28.2.4. Health and Well-Being of Arctic Residents

The warming Arctic and major changes in the cryosphere are significantly impacting the health and well-being of Arctic residents and projected to increase – especially, for many indigenous peoples. While impacts are expected to vary among the diverse settlements that range from small, remote predominantly indigenous to large cities and industrial settlements, this section focuses more on health impacts of climate change on indigenous, isolated, and rural populations because they are especially vulnerable to climate change due to a strong dependence on the

environment for food, culture and way of life; their political and economic marginalization; existing social, health, poverty disparities; as well as their frequent close proximity to exposed locations along ocean, lake or river shorelines (Ford and Furgal, 2009; Galloway-McLean, 2010; Larsen *et al.*, 2010; Cochran *et al.*, 2013).

28.2.4.1. Direct Impacts of a Changing Climate on the Health of Arctic Residents

Direct impacts of climate changes on the health of Arctic residents include extreme weather events, rapidly changing weather conditions, and increasingly unsafe hunting conditions (physical/mental injuries, death, disease), temperature-related stress (limits of human survival in thermal environment, cold injuries, cold-related diseases), and UV-B radiation (immunosuppression, skin cancer, non-Hodgkin's lymphoma, cataracts) (Revich, 2008; AMAP, 2009; IPCC, 2012). Intense precipitation events and rapid snowmelt are expected to impact the magnitude and frequency of slumping and active layer detachment resulting in rock falls, debris flow, and avalanches (Kokelj *et al.*, 2009; Ford *et al.*, 2010). Other impacts from weather, extreme events, and natural disasters are the possibility of increasingly unpredictable, long duration and/or rapid onset of extreme weather events, storms, inundation by large storm surges, which, in turn, may create risks to safe travel or subsistence activities, loss of access to critical supplies and services to rural or isolated communities (e.g. food, fuel, telecommunications), and risk of being trapped outside one's own community (Laidre *et al.*, 2008; Parkinson, 2009; Brubaker *et al.*, 2011b). Changing river and sea ice conditions affect the safety of travel for indigenous populations especially, and inhibit access to critical hunting, herding and fishing areas (Andrachuk and Pearce, 2010; Derksen *et al.*, 2012; Huntington and Watson, 2012).

Cold exposure has been shown to increase the frequency of certain injuries (e.g. hypothermia, frostbite) or accidents, and diseases (respiratory, circulatory, cardiovascular, musculoskeletal skin) (Revich and Shaposhnikov, 2010). Studies in Northern Russia have indicated an association between low temperatures and social stress and cases of cardiomyopathy (Revich and Shaposhnikov, 2010). It is expected that winter warming in the Arctic will reduce winter mortality rates, primarily through a reduction in respiratory and cardiovascular deaths (Shaposhnikov *et al.*, 2010). Researchers project that a reduction in cold-related injuries may occur, assuming that the standard for protection against the cold is not reduced (including individual behavior-related factors) (Nayha, 2005). Conversely, studies are showing respiratory and cardio stress associated with extreme warm summer days and that rising temperatures are accompanied by increased air pollution and mortality, especially in Russian cities with large pollution sources (Revich, 2008; Revich and Shaposhnikov, 2012).

28.2.4.2. Indirect Impacts of Climate Change on the Health of Arctic Residents

Indirect effects of climate change on the health of Arctic residents include a complex set of impacts such as changes in animal and plant populations (species responses, infectious diseases), changes in the physical environment (ice and snow, permafrost), diet (food yields, availability of country food), the built environment (sanitation infrastructure, water supply system, waste systems, building structures), drinking water access, contaminants (local, long-range transported), and coastal issues (harmful algal blooms, erosion) (Maynard and Conway, 2007; Brubaker *et al.*, 2011a; Parkinson and Evengard, 2009; Chapter 11).

In addition to the climate change impacts and processes, are the complicated impacts from contaminants such as POPs (persistent organic pollutants), radioactivity, and heavy metals (e.g., mercury) which create additional and/or synergistic impacts on the overall health and well-being of all Arctic communities (Armitage *et al.*, 2011; UNEP/AMAP, 2011; Teran *et al.*, 2012). Ambient temperature variability and temperature gradients directly affect the volatilization, remobilization, and transport pathways of mercury and POPs in the atmosphere, ocean currents, sea ice and rivers. Transport pathways, intercompartmental distribution, bioaccumulation and transformation of environmental contaminants such as persistent organic pollutants (POPs), mercury (Hg) and radionuclides in the Arctic may consequently be affected by climate change (AMAP 2011; UNEP/AMAP 2011; Ma *et al.*, 2011; Teng *et al.*, 2012) (*high confidence*). Ma *et al.* (2011) and Hung *et al.* (2010) demonstrated that POPs are already being remobilized into the air from sinks in the Arctic region as a result of decreasing sea ice and increasing temperatures.

Contaminants and human health in the Arctic are tightly linked to the climate and Arctic ecosystems by factors such as contaminant cycling and climate (increased transport to and from the Arctic), and the related increased risks of transmission to residents through subsistence life ways (Maynard, 2006; AMAP, 2010; Armitage *et al.*, 2011; UNEP/AMAP 2011; Teran *et al.*, 2012). The consumption of traditional foods by indigenous peoples places these populations at the top of the Arctic food chain and through biomagnification, therefore, they may receive some of the highest exposures in the world to certain contaminants (Armitage *et al.*, 2011; UNEP/AMAP, 2011). Contaminants such as POPs are known for their adverse neurological and medical effects on humans, particularly, the developing fetus, children, women of reproductive age and the elderly, thus it is important to include contaminants as a significant part of any climate impact assessment (UNEP/AMAP, 2011).

Radioactivity in the Arctic is also a concern because there are many potential and existing radionuclide sources in some parts of the Arctic and contamination can remain for long periods of time in soils and some vegetation, creating potentially high exposures for people (AMAP, 2010). Climate changes can mobilize radionuclides throughout the Arctic environment, and also potentially impact infrastructure associated with nuclear activities by changes in permafrost, precipitation, erosion, and extreme weather events (AMAP, 2010).

Warming temperatures are enabling increased overwintering survival and distribution of new insects that sting and bite as well as many bird, animal, and insect species that can serve as disease vectors and, in turn, causing an increase in human exposure to new and emerging infectious diseases (Parkinson *et al.*, 2008; Epstein and Ferber, 2011). Examples of new and emerging diseases are tick-borne encephalitis (brain infection) in Russia and Canada (Ogden *et al.*, 2010; Tokarevich *et al.*, 2011) and Sweden (Lindgren and Gustafson, 2001), *Giardia* spp. and *Cryptosporidium* spp. infection of ringed seals (*Phoca hispida*) and bowhead whales (*Balaena mysticetus*) in the Arctic Ocean (Hughes-Hanks *et al.*, 2005). It is also expected that temperature increases will increase the incidence of zoonotic diseases as relocations of animal populations occur (Revich *et al.*, 2012; Hueffler *et al.*, 2013).

Harmful algal blooms (HABs), whose biotoxins can be a serious health hazard to humans or animals (paralysis, death), are increasing globally and expected to increase in the Arctic, and HABs are influenced directly by climate change related factors such as temperature, winds, currents, nutrients and runoff (Portier *et al.*, 2010; Epstein and Ferber, 2011; Walsh *et al.*, 2011; Chapters 6; 11). Increasing ocean temperatures have caused an outbreak of a cholera-like disease, *Vibrio parahaemolyticus*, in Alaskan oysters (McLaughlin *et al.*, 2005). In addition, warmer temperatures raise the possibility of anthrax exposure in Siberia from permafrost thawing of historic cattle burial grounds (Revich and Podolnaya, 2011).

The impacts of climate change on food security and basic nutrition are critical to human health because subsistence foods from the local environment provide Arctic residents, especially, indigenous peoples, with unique cultural and economic benefits necessary to well-being and contribute a significant proportion of daily requirements of nutrition, vitamins and essential elements to the diet (Ford and Berrang-Ford, 2009; Ford, 2009). However, climate change is already an important threat due to the decrease in predictability of weather patterns, low water levels and streams, timing of snow, ice extent and stability, impacting the opportunities for successful hunting, gathering, fishing and access to food sources and increasing the probability of accidents (Ford and Furgal, 2009; Ford *et al.*, 2010). Populations of marine and land mammals, fish and water fowl are also being reduced or displaced, thus, reducing the traditional food supply (Gearheard *et al.*, 2006; West and Hovelsrud, 2010; Lynn *et al.*, 2013).

Traditional food preservation methods such as drying of fish and meat, fermentation, and ice cellar storage are being compromised by warming temperatures, thus further reducing food available to the community (Brubaker *et al.*, 2011b). For example, food contamination caused by thawing of permafrost “ice cellars” is occurring and increasingly wet conditions make it harder to dry food for storage (Hovelsrud *et al.*, 2011). Indigenous people increasingly have to abandon their semi-nomadic lifestyles, limiting their overall flexibility to access traditional foods from more distant locations (www.arctichealth.yukon.ca). These reductions in the availability of traditional foods plus general globalization pressures are forcing indigenous communities to increasingly depend upon expensive, non-traditional and often less healthy western foods, increasing the rates of modern diseases associated with processed food and its packaging, such as cardiovascular diseases, diabetes, dental cavities, and obesity (Armitage *et al.*, 2011; Berrang-Ford *et al.*, 2011; Brubaker *et al.*, 2011b).

Climate change is beginning to threaten community and public health infrastructure, often in communities with no central water supply and treatment sources. This is especially serious in low-lying coastal Arctic communities (e.g., Shishmaref, Alaska, USA; Tuktoyaktuk, Northwest Territories, Canada) through increased river and coastal flooding and erosion, increased drought and thawing of permafrost, resulting in loss of reservoirs, damage to landfill sites, or sewage contamination (GAO, 2009; Bronen, 2011). Salt-water intrusion and bacterial contamination may also be threatening community water supplies (Parkinson *et al.*, 2008; Virginia and Yalowitz, 2012). Quantities of water available for drinking, basic hygiene and cooking are becoming limited due to damaged infrastructure, drought, and changes in hydrology (Virginia and Yalowitz, 2012). Disease incidence caused by contact with human waste may increase when flooding and damaged infrastructure spreads sewage in villages with no municipal water supply. This can result in higher rates of hospitalization for pneumonia, influenza, skin infections, and respiratory viral infections (Parkinson and Evengard, 2009; Virginia and Yalowitz, 2012). Compounding these impacts in rural areas as well as cities are respiratory and other illnesses caused by airborne pollutants (e.g., contaminants, microbes, dust, mold, pollen, smoke) (Revich, 2008; Rylander and Schilling, 2011; Revich and Shaposhnikov, 2012).

It is now well-documented that the many climate-related impacts on Arctic communities are causing significant psychological and mental distress and anxiety among residents (Portier *et al.*, 2010; Coyle and Susteren, 2012; AR5 Chapter 11; Levintova, 2010). For example, changes in the physical environment (e.g., through thawing permafrost and erosion) which may lead to forced or voluntary relocation of residents out of their villages or loss of traditional subsistence species are causing mental health impacts among indigenous and other vulnerable, isolated populations (Curtis *et al.*, 2005; Albrecht *et al.*, 2007; Coyle and Susteren, 2012; Maldonado *et al.*, 2013). Special concern has been expressed by many communities about the unusually high and increasing numbers of suicides in the Arctic especially among indigenous youth, and efforts are under way to try to develop a thorough assessment as well as establish effective intervention efforts (Albrecht *et al.*, 2007; Portier *et al.*, 2010; USARC, 2010).

28.2.5. Indigenous Peoples and Traditional Knowledge

Indigenous populations in the Arctic – the original Native inhabitants of the region – are considered especially vulnerable to climate change, due to their close relationship with the environment and its natural resources for physical, social, and cultural well-being (Nuttall *et al.*, 2005; Parkinson, 2009; Cochran *et al.*, 2013). Arctic indigenous peoples are estimated to number between 400,000 and 1.3 million (Bogoyavlensky and Siggner, 2004; Galloway-McLean, 2010). According to the 2010 census data, there are 68,3 thousand indigenous people living in the Russian Arctic. These Arctic residents depend heavily on the region's terrestrial, marine and freshwater renewable resources, including fish, mammals, birds, and plants; however, the ability of indigenous peoples to maintain traditional livelihoods such as hunting, harvesting, and herding is increasingly being threatened by the unprecedented rate of climate change (Nakashima *et al.*, 2012; Cochran *et al.*, 2013). In habitats across the Arctic, climate changes are affecting these livelihoods through decreased sea ice thickness and extent, less predictable weather, severe storms, sea level rise, changing seasonal melt/freeze-up of rivers and lakes, changes in snow type and timing, increasing shrub growth, permafrost thaw, and storm-related erosion which, in turn, are causing such severe loss of land in some regions that a number of Alaskan coastal villages are having to relocate entire communities (Oskal, 2008; Mahoney *et al.*, 2009; Forbes and Stammler, 2009; Bartsch *et al.*, 2010; Weatherhead *et al.*, 2010; Brubaker *et al.*, 2011b; Bronen, 2011; Bongo *et al.*, 2012; Eira *et al.*, 2012; McNeeley, 2012; Huntington and Watson, 2012; Maldonado *et al.*, 2013). In addressing these climate impacts, indigenous communities must at the same time consider multiple other stressors such as resource development (oil and gas, mining), pollution, changes in land use policies, changing forms of governance, and the prevalence in many indigenous communities of poverty, marginalization, and resulting health disparities (Abryutina, 2009; Reinert *et al.*, 2009; Magga *et al.*, 2011; Nakashima *et al.*, 2012; Vuojala-Magga *et al.*, 2011).

Traditional knowledge is the historical knowledge of indigenous peoples accumulated over many generations and it is increasingly emerging as an important knowledge base for more comprehensively addressing the impacts of environmental and other changes as well as development of appropriate adaptation strategies for indigenous communities (IPCC AR4; IPCC AR5, chp 15; Oskal, 2008; Reinert *et al.*, 2008; Wildcat, 2009; Nakashima *et al.*, 2012; Magga *et al.*, 2011; Vuojala-Magga *et al.*, 2011; Vogesser *et al.*, 2013) For example, Saami reindeer herders have specialized knowledge of dynamic snow conditions, which mediate access to forage on autumn, winter and

spring reindeer rangelands (Roturier and Roue, 2009; Eira *et al.*, 2012; Vikhamar-Schuler *et al.*, 2013) and traditional governance systems for relating to natural environments (Sara, 2013). Increasingly, traditional knowledge is being combined with western scientific knowledge to develop more sustainable adaptation strategies for all communities in the changing climate.

For example, at Clyde River, Nunavut, Canada, Inuit experts and scientists both note that wind speed has increased in recent years and that wind direction changes more often over shorter periods (within a day) than it did during the past few decades (Gearheard *et al.*, 2010; Overland *et al.*, 2012). In Norway, Sámi reindeer herders and scientists are both observing direct and indirect impacts to reindeer husbandry such as changes in snow and ice cover, forage availability and timing of river freeze-thaw patterns from increasing temperatures. (Eira *et al.*, 2012). On the Yamal Peninsula in western Siberia, detailed Nenets observations and recollections of iced-over autumn and winter pastures due to rain-on-snow events have proven suitable for calibrating the satellite-based microwave sensor SeaWinds (Bartsch *et al.*, 2010) and NASA's AMSR-E sensor (Bongo *et al.*, 2012).

28.2.6. Economic Sectors

28.2.6.1. Arctic

28.2.6.1.1. Agriculture and forestry

Climate change presents benefits and costs for forestry and agriculture (Aaheim *et al.*, 2009; Hovelsrud *et al.*, 2011). In Iceland for example tree limits are found at higher altitudes than before, and productivity of many plants has increased (Björnsson *et al.*, 2011). Grain production in Iceland, has increased in the last two decades, and work on soil conservation and forestry has benefited from warming (Sigurdsson *et al.*, 2007; Björnsson *et al.*, 2011), but also the number of new insect pests on trees and shrubs has increased in the past 20 years. A strong relationship between rate of new insect pest colonisation and outbreak intensity in forests exists with changes in annual temperature during the past century (Halldórsson *et al.*, 2013). Climate change impacts on species change and fire frequency have potential impact on commercial forest harvesting activity. Vulnerability of forestry to changes that affect road conditions and thus accessibility during thawing periods has been found in Sweden (Keskitalo, 2008). A case study on Greenland found challenges for plant diseases in potatoes and grass fields, with pathogens and pests present in agricultural cropping systems, e.g. black scurf (*Rhizoctonia*) and common scab (*Streptomyces scabies*) (Neergaard *et al.*, 2009).

28.2.6.1.2. Open and freshwater fisheries

Current commercial fisheries are sharply divided between regions of high-yield and value commercial fisheries in the southern Bering Sea, Baffin Bay, the east and west Greenland Seas, the Iceland Shelf Sea, the deep Norwegian/Greenland Sea, and the Barents Seas and subsistence fisheries in the coastal regions of the Arctic Ocean. The relative absence of commercial fishing activity in the Arctic Ocean results from a combination of fisheries policy, the abundance of the resource, the lack of infrastructure for capturing and processing fish, and the difficulties in accessing fishing grounds especially during winter. In most regions, fisheries management strategies have been developed to build sustainable fisheries and rebuild overfished stocks (Froese and Proelß, 2010; Livingston *et al.*, 2011). Recently observed changes in the spatial distribution and abundance of mackerel (*Scomber scombrus*) has challenged existing international agreements for shared resources in the North Atlantic (Arnason, 2012; Astthorsson *et al.*, 2012). Although loss of sea ice in summer is allowing greater access to fisheries resources in the Arctic Ocean, some nations have prohibited commercial fishing within their EEZ until there is sufficient understanding of stock status to ensure that proposed fisheries would be managed sustainably (Wilson and Ormseth, 2009; Stram and Evans, 2009).

Several Arctic coastal sea-run fishes are targeted for subsistence and commercial use in the Arctic. Commercial transactions from fishing are typically for local markets, however, the socioeconomic and cultural importance of these fishes to Indigenous Peoples far outweighs their monetary value. Reist *et al.* (2006) and Fehhelm *et al.* (2007)

found that climate related factors that influenced the water level and freshening of rivers were related to run size of arctic cisco (*Coregonus autumnalis*). Similarly, a recent study based on Chinook salmon (*Oncorhynchus tshawytscha*) run timing for the period (1961-2009) showed that success in the fishery was dependent on the timing of the marine exit, which was tightly coupled to environmental conditions that were linked to climate (Mundy and Evenson, 2011).

28.2.6.1.3. *Marine transportation*

Observations and climate models indicate that in the period between 1979-1988 and 1998-2007 the number of days with ice free conditions (less than 15% ice concentration) increased by 22 days along the Northern Sea Route (NSR) in the Russian Arctic, and by 19 days in the Northwest Passage (NWP) in the Canadian Arctic, while the average duration of the navigation season in the period 1980-1999 was 45 and 35 days, respectively (Mokhow and Khon, 2008). Increased shipping associated with the opening of the NSR will lead to increased resource extraction on land and in the sea, and with two-way commodity flows between the Atlantic and Pacific. The future status of marine, terrestrial and freshwater biota may be negatively affected due to substantial coastal infrastructure to facilitate offshore developments (Meschtyb *et al.*, 2010). Also, the frequency of marine transportation along the NSR is at its highest during the most productive and vulnerable season for fish and marine mammals, which is the late spring/summer, when these resources can be found throughout the NSR area (Østreng, 2006).

28.2.6.1.4. *Infrastructure*

Much of the physical infrastructure in the Arctic rely on and are adapted to local sea-ice conditions, permafrost, and snow (Huntington *et al.*, 2007; Sundby and Nakken, 2008; West and Hovelsrud, 2010; Forbes, 2011; Sherman *et al.*, 2009). Damage from ice action and flooding to installations such as bridges, pipelines, drilling platforms, and hydropower poses major economic costs and risks, which are more closely linked to the design of the structure than with thawing permafrost. Current engineering practices are designed to help minimize the impacts (Prowse *et al.*, 2009). Much of the infrastructure has been built with weather conditions in mind, but remains vulnerable and inadequate to respond to environmental emergencies, natural disasters, and non-environmental accidents (NRTEE, 2009). Northern safety, security, and environmental integrity are much dependent upon transportation infrastructure. Ice as a provisioning system provides a transportation corridor and a platform for a range of activities and access to food sources in the Arctic (Eicken *et al.*, 2009).

In Northern Canada climate warming presents an additional challenge for northern development and infrastructure design. While the impacts of climate change become increasingly significant over the longer time scales, in the short term of greater significance will be the impacts associated with ground disturbance and construction (Smith and Risebrough, 2010).

Climate change impacts have increased the demand for improved communication infrastructure and related services and community infrastructure for the safety and confidence in drinking water (NRTEE, 2009). The access, treatment and distribution of drinking water is generally dependent upon a stable platform of permafrost for pond or lake retention. Several communities have reported the need for more frequent water-quality testing both municipal systems and untreated water sources to ensure its suitability for drinking (Furgal, 2008).

28.2.6.1.5. *Resource exploration*

The Arctic has large reserves of minerals (Lindholt, 2006; Peters *et al.*, 2011; Harsem *et al.*, 2011) and potentially large reserves of undiscovered sources of raw minerals, and oil and gas. Predicted new access to offshore energy resources is hypothesized to be a significant share of the global supply of oil and gas (Gautier *et al.*, 2009; Berkman *et al.*, 2010). The socio-economic impacts of oil and gas exploration activity may be positive or negative (Duhaime *et al.*, 2004; Huntington *et al.*, 2007; Forbes, 2008; Kumpula *et al.*, 2011; Forbes *et al.*, 2009; Harsem *et al.*, 2011).

Climatic warming is accelerating access to northern lands for development (Forbes *et al.*, 2009). Yamal in Western Siberia has approximately 90 % of Russia's gas reserves, but at the same time represents the largest area of reindeer herding in the world (Jernsletten and Klokov, 2002; Stammler, 2005; Forbes and Kumpula, 2009). Development activities to obtain these resources would shrink the grazing lands, and have been characterized as one of the major human activities in the Arctic contributing to loss of "available room for adaptation" for reindeer husbandry (Oskal, 2008; Forbes *et al.*, 2009; Nuttall *et al.*, 2005). Sharp increases in future oil and gas and other resource development in the Russian North and other Arctic regions is anticipated - along with associated infrastructure, pollution, and other by products of development - which will reduce the availability of pasturelands for reindeer and indigenous communities (Derome and Lukina, 2011; Degteva and Nellermann, 2013).

28.2.6.1.6. *Informal, subsistence-based economy*

Hunting, gathering, herding, and fishing for subsistence, as well as commercial fishing, all play an important role in the mixed cash-subsistence economies (Nuttall *et al.*, 2005; Poppel and Kruse, 2009; Larsen and Huskey, 2010; Crate *et al.*, 2010). In the early 1990s - initially in western Canada, and later elsewhere - indigenous communities started reporting climate change impacts (Berkes and Armitage, 2010). According to some herders, whalers and walrus hunters non-predictable conditions resulting from more frequent occurrence of unusual weather events are the main effect of recent warming (Forbes and Stammler, 2009; Ignatowski and Rosales, 2013; Forbes *et al.*, 2009).

The Inuit and Saami have expressed strong concern about the effects of climate warming on their livelihoods (Forbes and Stammler, 2009; Magga *et al.*, 2011). For the Inuit, the issues revolve around sea ice conditions, such as later freeze-up in autumn, earlier melt-out and faster sea ice retreat in spring, and thinner, less predictable ice in general (Krupnik and Jolly, 2002; Cochran *et al.*, 2013). Diminished sea ice translates into more difficult access for hunting marine mammals, and greater risk for the long-term viability of subsistence species such as polar bear populations (Laidre *et al.*, 2008). Most Inuit communities depend to some extent on marine mammals for nutritional and cultural reasons, and many benefit economically from polar bear and narwhal hunting. A reduction in these resources represents a potentially significant economic loss (Hovelsrud *et al.*, 2008). Among Fennoscandian Saami, the economic viability of reindeer herding is threatened by competition with other land users coupled with strict agricultural norms (Forbes, 2006; Magga *et al.*, 2011). Reindeer herders are concerned that more extreme weather may exacerbate this situation (Oskal, 2008).

Climate change is affecting reindeer herding communities through greater variability in snow melt/freeze, ice, weather, winds, temperatures and precipitation, which, in turn are affecting snow quality and quantity - the most critical environmental variables for reindeer sustainability (Magga *et al.*, 2011; Eira *et al.*, 2012). Increasing temperature variations in wintertime, with temperatures rising above freezing with rain, followed by refreezing ("rain-on-snow" conditions), are becoming more frequent, forming ice layers in the snow which then block the animals' access to their forage and subsequent starvation (Maynard *et al.*, 2011; Eira *et al.*, 2012; Bongo *et al.*, 2012).

28.2.6.2. *Antarctica and the Southern Ocean*

Economic activities in the Antarctic have been limited to fishing and tourism (IPCC WG2, 2007). Ship-based tourism is a significant industry in Antarctica but does not involve permanent shore-based infrastructure. Over recent decades, the number of tourists landing in Antarctica has risen from 7322 in 1996/1997 to 32,637 in 2007/2008 (IAATO, 2012). Visits generally coincide with the times when wildlife are breeding and are often restricted because of the presence of fast ice, sea ice or icebergs. They are expected to continue to increase, with an increasing chance of terrestrial alien species being introduced from tourism and other vectors as ice-free areas increase from climate change (Chown *et al.*, 2012). Scientific activity by a number of nations is also taking place and has the potential to impact upon local ecologies. Mineral resource activity is prohibited south of 60°S under the Protocol on Environmental Protection to the Antarctic Treaty.

Fisheries in Antarctica, primarily through fisheries for Antarctic krill, could amount to approximately 6% of existing global marine capture fisheries (Nicol *et al.*, 2011). The pattern of the krill fishery has been affected by changes in the sea ice extent around the Antarctic Peninsula where the fishery has been taking advantage of the ice-free conditions and taking more of its catch during winter in that region (*high confidence*) (Kawaguchi *et al.*, 2009). Ecosystem-based management of krill fisheries by the Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR) is yet to include procedures to account for climate change impacts, although the need to do so has been identified (Trathan and Agnew, 2010; Constable, 2011).

28.3. Key Projected Impacts and Vulnerabilities

28.3.1. Hydrology and Freshwater Ecosystems

28.3.1.1. Arctic

Accompanying projected increases in high-latitude river flow (WGI Ch.12.4.5.4. and WGII Ch. 3.4.5.) are earlier spring runoff (Dankers and Middelkoop, 2008; Hay and McCabe, 2010; Pohl *et al.*, 2007), greater spring snowmelt (Adam *et al.*, 2009) and increases in spring sediment fluxes (Lewis and Lamoureaux, 2010). Enhanced permafrost thaw (WGI Ch. 12.4.6.2) will continue to affect the dynamics of thermokarst lakes and related ecological effects (Section 28.2.1.1). Thawing permafrost and changes in the hydrological regime of the Arctic rivers, particularly those traversing regions affected by industrial developments, will increase the contaminant flow (Nikanorov *et al.*, 2007). Loss of glacier ice masses will alter runoff hydrographs, sediment loads, water chemistry, thermal regimes, and related channel stability, habitat and biodiversity (Milner *et al.*, 2009; Moore *et al.* 2009).

Although snow, freshwater ice and permafrost affect the morphology of arctic alluvial channels, their future combined effects remain unclear (McNamara and Kane, 2009). For small permafrost streams, however, longer projected periods of flowing water will modify nutrient and organic matter processing (Greenwald *et al.*, 2008; Zarnetske *et al.*, 2008) but long-term negative impacts of increased sediment load on biological productivity could outweigh any positive effects from increased nutrient loading (Bowden *et al.*, 2008).

Changes to river-ice flooding are also projected to occur due to changes in i) hydraulic gradients for near-coastal locations because of sea-level rise, ii) streamwise air-temperature gradients, and iii) the timing and magnitude of spring snowmelt (Prowse *et al.* 2011). Synergistic/antagonistic effects among these factors, however, require detailed site-specific analyses for accurate projections of future conditions (Beltaos and Prowse, 2009). Reduced (increased) ice-jam flooding will have positive (negative) benefits for river-side northern communities/infrastructure but could also alter delta-riparian (Lesack and Marsh, 2010) and coastal-marine (Emmerton *et al.*, 2008) ecosystems. The quality of river water entering the marine environment will also be affected by the reduction or loss of stamukhi lakes that process river inputs (Dumas *et al.*, 2006; Galand *et al.*, 2008).

Future changes to lake-ice regimes will include: delayed freeze-up, advanced break-up, thinner ice and changes in cover composition (especially white ice in areas of enhanced winter precipitation), increased water temperature, and earlier and longer-lasting summer stratification (Dibike *et al.* 2011), all of which will affect a range of aquatic processes, including secondary productivity (Prowse and Brown, 2010b; Borgström and Museth, 2005; Prowse *et al.*, 2007). Patterns of species richness and diversity are also projected to change with alterations to ice duration - increased open-water periods favouring the development of new trophic levels, colonization of new aquatic species assemblages (Vincent *et al.*, 2009), greater atmosphere-water gas exchange and a decrease in winter kill of resident fish with cascading effects on lower trophic levels (Balayla *et al.*, 2010). The loss of ice, however, can also decrease key habitat availability and quality (Vincent *et al.*, 2008).

Geochemical responses of Arctic lakes will also be altered. As observed for thermokarst lakes, the loss of ice cover and associated warming can greatly increase methane production (Laurion *et al.*, 2010; Metje and Frenzel, 2007). Because temperature sensitivity has a stronger control over methane production than oxidation (Duc *et al.*, 2010), elevated water temperatures will enhance methanogenesis, causing increased methane release from sediments. The

net balance of these two processes operating under a broad range of future changing environmental factors, however, remains to be quantified (Laurion *et al.*, 2010; Walter *et al.*, 2007a, b; 2008).

As well as methane, increased water temperatures are projected to lead to reduced organic carbon (OC) burial. Projections, based on a range of six climate warming scenarios (Solomon *et al.*, 2007), indicates that there will be a 4-27% decrease (0.9-6.4 TgC yr⁻¹) in OC burial across lakes of the northern boreal zone by the end of the 21st Century as compared to rates for the approximately last half-century (Gudasz *et al.*, 2010). Although these estimates assume that future organic carbon delivery will be similar to present-day conditions, even with enhanced supply from thawing permafrost, higher water temperatures will increase organic carbon mineralization and thereby lower burial efficiency. The amount of burial also depends on lake depth and mixing regimes. For non-thermally stratified shallow lakes, there will be a greater opportunity for water-sediment mixing and hence, greater carbon recycling back into the water column. By contrast, for lakes that become increasingly thermally stratified, carbon sinking below the thermocline will tend not to return to the surface until an increasing later fall turnover, thereby decreasing the probability of sediment-stored carbon being returned to the water column (Flanagan *et al.*, 2006).

Changes in ice cover, thermal regimes and stratification patterns will also affect the fate of contaminants in northern lakes. Higher water temperatures can enhance the methylation of mercury and modify food-web and energy pathways, such as through enhanced algal scavenging (a major foodweb entry pathway for mercury) resulting in increased mercury bio-availability to higher trophic levels (Carrie *et al.*, 2010; Outridge *et al.*, 2007).

28.3.1.2. Antarctica

This assessment reinforces conclusions of AR4. Increased temperatures will impact aquatic ecosystems in Antarctica (*high confidence*) but the exact nature of these impacts will vary regionally. The most vulnerable freshwater systems are in the northern Antarctic Peninsula and maritime Antarctic islands, where a small increase in temperature can have widespread ecosystem impacts because the average temperature is within a few degrees of the melting point (Quesada and Velázquez 2012) (*high confidence*). Potential impacts are expected to range from immediate catastrophic impacts such as loss of bounding ice masses causing drainage of freshwater and epishelf lakes (Smith *et al.*, 2006; Hodgson, 2011), to more gradual impacts on changes in the amount and duration of catchment ice and snow cover, accelerated glacier melting, declining volumes of precipitation falling as snow, permafrost, active layer and hydrological changes, such as water retention times (e.g. Vieira *et al.*, 2010, Quesada and Velázquez 2012, Bockheim *et al.*, 2013) (*medium confidence*).

Changes in the thickness and duration of seasonal ice cover, longer melt seasons and larger volumes of water flowing into the lakes are expected in the future (Lyons *et al.*, 2006) (*medium confidence*) but the ecological effects will vary between lakes, depending on their depth to surface area ratio, with *insufficient evidence* to fully assess future changes in these systems. Longer ice free seasons may cause physical conditions to be more favorable for primary production (Hodgson and Smol 2008) but very high irradiances experienced during summer in some systems can substantially inhibit algal blooms under ice free conditions (Tanabe *et al.*, 2007), which would favor the growth of benthic cyanobacteria species (Hodgson *et al.*, 2005). In other lakes, increases in meltwater supply may increase suspended solids and reduce light penetration and may offset the increases in the underwater light regime predicted as a result of extended ice free periods (Quesada *et al.*, 2006).

Under a warming climate an increase in microbial biomass is likely because of the increased water supply from glacial melt and warmer temperatures, and could result in further development of soils and elevated nutrient and dissolved organic carbon delivery to lakes (Velázquez *et al.*, 2013). This organic supply will promote growth and reproduction in the benthos and plankton and imbalances in population dynamics (Quesada and Velázquez, 2013). Nutrient enrichment of some freshwater habitats in the vicinity of fur seal colonies will increase because of expanding fur seal populations (Quayle *et al.* 2013) (*high confidence*).

Away from glacial forelands, increasing aridity will occur in the long-term in some areas of the continent (Hodgson *et al.*, 2006b) and on subantarctic islands (Smith Jr *et al.*, 2012) (*medium confidence*). Closed basin lakes can dry up completely causing local extinctions or retreat into cryptic or resistant life-cycle stages, as experienced in Arctic

lakes (Smol and Douglas, 2007b). Other effects include dessication of moss banks due to increased evaporation and sublimation rates (Wasley *et al.*, 2006) (*medium confidence*). Studies have also shown that warming of once cold freshwater habitats in Antarctica will allow the sub- and maritime Antarctic species to re-invade and establish self-maintaining populations on the Antarctic continent, particularly where human vectors are involved (Barnes *et al.*, 2006, Hodgson *et al.*, 2006b) (*medium confidence*). For other organisms with lower dispersal capabilities there is increasing evidence of endemism, particularly in microbial groups (Vyverman *et al.*, 2010), with a possibility that surface Antarctic lakes contain endemic species that are relicts of Gondwana (cf. Convey and Stevens 2007) and that would become extinct should they be lost from these lakes as a result of climate change..

28.3.2. Oceanography and Marine Ecosystems

28.3.2.1. Ocean Acidification in the Arctic and Antarctic

Ocean acidification on polar marine food webs can have considerable implications (*medium certainty*). For example, if some regions in the Arctic become under-saturated with respect to aragonite (the primary structural component of the shells of some marine calcifiers such as mollusks and urchins) the growth and survival of these organisms will be impacted (WG I, Chapter 6 Figure 6.28; Chierici and Fransson, 2009; Fabry *et al.*, 2009; Yamamoto-Kawai *et al.*, 2009). In laboratory experiments, Arctic pteropods (*Limacina helicina*, a small planktonic mollusk) held under conditions consistent with projected ocean warming and acidification in the Arctic Ocean in early spring were able to extend their shells in corrosive waters but dissolution marks were observed (Comeau *et al.*, 2010; Comeau *et al.*, 2012). Additional studies are needed to scale up regional impacts to assess the population level impact of ocean acidification on *Limacina helicina* and other vulnerable species (Orr *et al.*, 2009). At the current time there is insufficient data to fully assess the ecosystem consequences of acidification on pteropods because it is unclear whether other species, with a similar nutritive value, will replace pteropods.

In the Southern Ocean, foraminifera have thinner shells than in the Holocene and there is evidence for shell thickness to be related to atmospheric CO₂, supporting the hypothesis that ocean acidification will affect this abundant protozoan in this region (Moy *et al.*, 2009). Similarly, shells are thinner from sediment traps in aragonite under-saturated water (below the aragonite saturation horizon - ASH) compared to those captured above the ASH in Subantarctic waters, but there is no time series of data related to change in the ASH (Roberts *et al.*, 2011). Shell dissolution has been observed in surface waters in the Atlantic sector as a result of both upwelling and atmospheric changes in CO₂ (Bednarsek *et al.*, 2012) (*medium confidence*). Other impacts of acidification on Southern Ocean organisms are currently uncertain, but short term negative impacts need to be considered together with an organism's capacity to adapt in the longer term (Watson *et al.*, 2012).

Only a few studies have been conducted on commercially exploited polar species on ocean acidification. Antarctic krill embryonic development (Kawaguchi *et al.*, 2011) and post-larval krill metabolic physiology (Saba *et al.*, 2012) may be impeded by elevated CO₂ concentrations, which may negatively impact the reproductive success of krill more generally under emission scenarios used in CMIP5 (Kawaguchi *et al.*, 2013) (*medium confidence*). Long *et al.* (2013) examined the effects of acidification on red king crab (*Paralithodes camtschaticus*) and found animals exposed to reduced pH exhibited increased hatch duration, decreased egg yolk, increased larval size, and decreased larval survival. In contrast, Hurst *et al.*, (2012) conducted laboratory experiments at levels of elevated CO₂ predicted to be present in the Gulf of Alaska and Bering Sea in the next century and found that juvenile walleye pollock exhibited a general resiliency of growth energetics to the direct effects of CO₂ changes.

28.3.2.2. Arctic

28.3.2.2.1. Marine plankton, fish, and other invertebrates

Phenological response

Projected changes in the timing, spatial distribution and intensity of spring blooms may result in mis-matches with the timing of the emergence of Arctic grazers (Søreide *et al.*, 2010). Based on past experience, some species will adapt to local conditions by shifting key life cycle events (hatch-date, maturity schedule and reproductive timing) or diet to accommodate differences in the regional timing and availability of prey and environmental conditions (Ormseth and Norcross, 2007; Sundby and Nakken, 2008; Vikebø *et al.*, 2010, Darnis *et al.*, 2012). For example, loss of sea ice cover in spring is expected to change fish behavior in ice bound areas (Mundy and Evenson, 2011). It is uncertain whether endemic animals will be able to alter key phenologies fast enough to keep pace with the projected rates of change in the Arctic Ocean.

Projected spatial shifts

Simulation studies revealed that a 2 week longer growing season and a 2 degree C increase in temperature would not be sufficient to allow expatriate species (*Calanus finmarchicus* or *C. marshallae*) to invade the Arctic Ocean (Ji *et al.*, 2012). Ellingsen *et al.*, (2008) projected future zooplankton distribution and abundance in the Barents Sea for the period 1995-2059 using a regional climate model which was forced with climate model output based on the IPCC-SRES B2 scenario. They projected that by 2059, Atlantic origin zooplankton will increase and Arctic origin zooplankton will decrease in the Barents Sea.

The literature is mixed with respect to the potential for future movement of fish and shellfish into the Arctic Ocean. Modeling studies project that marine fish stocks potentially will shift their distributions into the Arctic Ocean resulting in an increase in biodiversity in the region (Cheung *et al.*, 2009; Cheung *et al.*, 2011; Box CC-MB). However, other studies show the persistence of cold sea water temperatures on the shelf regions of the Arctic Ocean and Northern Bering Sea will restrict or retard the movement of several sub-arctic fish and shellfish species into the Arctic Ocean (Sigler *et al.*, 2011; Stabeno *et al.*, 2012b; Hunt *et al.*, 2013). In waters off the coasts of Europe there is a potential for increased fish production because of the combined effects of intrusion of Atlantic water over the relatively broader shelf regions and advective corridors for larval drift and range expansion of spawners. Huse and Ellingsen (2008) forced a spatially explicit coupled bio-physical model for the Barents Sea with future climate scenarios to project the implications of climate change on the spawning distribution of capelin (*Mallotus villosus*). Projections show that the spawning distribution of capelin will shift to the east and new spawning grounds will be colonized. A key factor governing this expansion will be the availability of pelagic prey. In the Bering Sea, there is evidence that planktivorous species like walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea will shift their distribution in response to shifts in ocean temperature (Kotwicki and Lauth, 2013). In summary, the spatial distribution of fish and shellfish in the Barents and Bering Seas will shift in response to climate change (*high confidence*)

Projected impacts on production

In the deep basins of the Arctic Ocean the number of ice free days in summer are expected to result longer productive seasons (Slagstad *et al.*, 2011, *high confidence*). Ellingsen *et al.* (2008) projected that annual primary production would increase by 2059 in the Barents Sea. Tremblay *et al.* (2012) hypothesized that longer ice free periods in summer in the Arctic Ocean could provide for more opportunities for episodic nutrient pulses that would enhance secondary production through the growing season. However, in the Arctic Ocean, these changes in primary production may be offset later in the year by increased zooplankton grazing (Olli *et al.*, 2007) or nutrient depletion due to stronger stratification and shifts in the mixed layer depth (Tremblay *et al.*, 2012; Wassmann, 2011). Therefore, there is *medium confidence* that annual phytoplankton production will increase the central Arctic Ocean.

In the few cases where future abundance of fish has been projected using climate change scenarios, species exhibited different trends related to their vulnerability. Forward extrapolation of observed responses suggests that increased summer sea surface temperatures in the Bering Sea and Barents Sea will cause a decrease in the abundance of energy rich copepods and euphausiids (Coyle *et al.*, 2011; Slagstad *et al.*, 2011). This change in prey quality is expected to lower survival of walleye pollock in the eastern Bering Sea by 2050 (Mueter *et al.*, 2011). Climate enhanced stock projection models showed time trends in cross-shelf transport of juvenile northern rock sole (*Lepidopsetta polyxystra*) to nursery areas will not be substantially altered by climate change (Wilderbuer *et al.*, 2012).

28.3.2.2.2. *Marine mammals, polar bear, and seabirds*

The effects of the projected reduction in sea ice extent in this century (Wang and Overland, 2011) on Arctic marine mammals and sea birds will vary spatially and temporally (Laidre *et al.*, 2008). Many ice-associated marine mammals and sea birds will be affected by ice loss with altered species distributions, migration patterns, behaviour, interspecific interactions, demography, population changes, and vulnerability to extinction but there is limited evidence of changes for most species (*high confidence*).

The polar bear population of the Southern Beaufort Sea is projected to decline by 99% by 2100 with a probability estimated at 0.80-0.94 under A1B (Hunter *et al.*, 2010). The Northern Beaufort Sea population is stable although decline is predicted with warming (Stirling *et al.*, 2011). Projected extirpation of approximately two-thirds of the world's polar bears was predicted for mid-century under A1B (Amstrup *et al.*, 2008). Aspects of this study were criticized (Armstrong *et al.*, 2008) but refuted (Amstrup *et al.*, 2009). The two-thirds decline is consistent with other studies and has robust evidence with *medium agreement*. Projected extinction of polar bears is *unlikely*. There is *very high confidence* of subpopulation extirpation.

Several factors other than climate influence sea bird population dynamics (Regular *et al.*, 2010), and projections of changes with a continued Arctic warming are therefore highly uncertain. Pattern of change will be non-uniform and highly complex (ACIA, 2005). At present, the resolution of AOGCMs are not detailed enough to project spatial changes in mesoscale oceanographic features like frontal zones and eddies of importance to Arctic sea birds.

It is likely that the high Arctic seabird species partly or completely dependent on the sympagic ecosystem or the cold Arctic waters close to the ice-edge will be negatively impacted if the projected changes in these physical parameters occur (*medium confidence*). A general increase in SSTs, retreat of the ice cover, and earlier break up of fast ice may improve the environmental conditions and food abundance for sea bird species that have their range in the southern part of the Arctic or south of the Arctic (*medium confidence*). A poleward expansion of the range of these species is expected during a continued warming (*medium confidence*).

Several factors other than climate influence sea bird population dynamics (Regular *et al.*, 2010), and projections of changes with a continued Arctic warming are therefore highly uncertain. Pattern of change will be non-uniform and highly complex (ACIA 2005). At present, the resolution of AOGCMs are not detailed enough to project spatial changes in mesoscale oceanographic features like frontal zones and eddies of importance to Arctic sea birds.

28.3.2.3. *Antarctica and the Southern Ocean*

Continued rising temperatures in the Southern Ocean will result in increased metabolic costs in many ectothermic pelagic species, southward movement of temperate species and contraction of the range of polar species (*medium confidence*). Southward movement of ocean fronts and associated biota that are prey of subantarctic island-based predators, will result in energetic inefficiencies for some of those predators (Péron *et al.*, 2012; Weimerskirch *et al.*, 2012) (*low confidence*).

For Antarctic krill, *insufficient evidence* is available to predict what will happen to the circumpolar productivity of krill because of regional variability of the effects of climate change on the different factors (positive and negative)

that affect krill, directly and indirectly. For example, increased metabolic and growth rates from warming may be countered by a reduced food supply and the effects of ocean acidification (28.2.2.2, 28.3.2.1). Also, areas that are already warm may result in slower growth with further warming, such as could happen in the northern Scotia Arc (Wiedenmann *et al.*, 2008; Hill *et al.*, 2013). Models of recruitment and population dynamics indicate that the biomass of krill will decline if surface warming continues, but preliminary projections incorporating a range of factors are uncertain (*low confidence*) (Murphy *et al.*, 2007, 2012b). Physiological and behavioural responses might also ameliorate impacts. For example, krill are now known to exploit the full depth of the ocean, which could provide escapes from further warming (Schmidt *et al.*, 2011) as well as refuge from air-breathing predators.

The strong dependence of species in more southern regions (e.g. southern west Antarctic Peninsula, WAP, and Ross Sea region) on sea-ice means that changes in sea-ice distribution will cause spatial shifts in the structure of ice-obligate food webs (Murphy *et al.*, 2012b) (*low confidence*). Projections show that the loss of summer sea ice from the west Antarctic Peninsula is expected to result in ice-dependent seals declining and being replaced by other seal species that are not dependent on sea ice (Siniff *et al.*, 2008, Costa *et al.*, 2010) (*low confidence*). There is *insufficient evidence* to determine whether there will be a mismatch in phenologies of different species as a result of changes in the winter sea ice season (timing and winter extent), such as might occur if the timing of sea ice melt was not at a time of optimal growing conditions for phytoplankton (Trathan and Agnew, 2010).

Reductions in krill abundance in the marine food webs around the South Atlantic islands may result in a shift in their structure towards a more fish-centred ecosystem as observed in the Indian Sector (Trathan, *et al.*, 2007; Shreeve *et al.*, 2009; Waluda *et al.*, 2010; Trathan *et al.*, 2012; Murphy *et al.*, 2012a; Murphy *et al.*, 2012b) (*low confidence*). Also, salps have been postulated to be competitors with krill for phytoplankton around the Antarctic Peninsula when oceanic conditions displace shelf and near-shelf waters during times of low sea ice (Ducklow *et al.*, 2012). In the absence of krill, longer food chains have lower trophic efficiency (Murphy *et al.*, 2012; Muprhy *et al.*, 2013) and the long-term implications of this for higher trophic levels are unknown.

Coastal environments will be impacted by the dynamics of fast ice, ice shelves and glacier tongues. These factors will positively affect local primary production and food web dynamics (Peck *et al.*, 2009) but negatively affect benthic communities (Barnes and Souster, 2011) (*low confidence*). Projections of the response of emperor penguins and Southern Ocean seabirds based on AR4 model outputs for sea ice and temperature in east Antarctica indicate that general declines in these populations are to be expected if sea ice habitats decline in the future (Barbraud *et al.*, 2011; Jenouvrier *et al.*, 2012) (*low confidence*). However, these responses are also expected to be regionally specific because of the regional differences in expectations of change in the ice habitats (*high confidence*). Additional studies at other sites are needed to improve confidence levels of predictions.

28.3.3. Terrestrial Environment and Related Ecosystems

28.3.3.1. Arctic

The boreal forest is generally projected by models to move northward under a warming climate, that will displace between 11% and 50% of the tundra within 100 years (Callaghan *et al.*, 2005; Wolf *et al.*, 2008; Tchebakova *et al.*, 2009; Wramneby *et al.*, 2010 in a pattern similar to that which occurred during the early Holocene climatic warming (*high confidence*). Pearson *et al.* (2013) projected that at least half of vegetated Arctic areas will shift to a different physiognomic class, and woody cover will increase by as much as 52%, in line with what has been occurring in northwest Eurasia (Macias-Fauria *et al.* 2012).

Dynamic vegetation models applied to Europe and the Barents Region project a general increase in net annual primary production by climate warming and CO₂ fertilization (Wolf *et al.*, 2008; Wramneby *et al.*, 2010; Anisimov *et al.*, 2011). Boreal needle-leaved evergreen coniferous forest replaces tundra and expands into the mountain areas of Fennoscandia, but this advance may be delayed or prevented in regions already occupied by clonal deciduous shrubs whose *in situ* growth has increased significantly in recent decades (Macias-Fauria *et al.* 2012).

In contrast to these expected results, shrubs, currently expanding in area in many Arctic locations, were modelled to decrease in extent over the next 100 years after an initial increase (Wolf *et al.*, 2008). Also counter-intuitively, tundra areas increased in the projections. This was a result of changes at the highest latitudes that opened land for colonisation at a rate exceeding displacement of tundra by shrubs in the south.

Several studies have calculated the magnitude of the effects of vegetation change in the Arctic on negative feedbacks of CO₂ sequestration and increased evapo-transpiration and the positive feedback of decreased albedo (Swann *et al.*, 2010; Wramneby *et al.*, 2010; Wolf *et al.*, 2010; Pearson *et al.*, 2013). It is *likely* that vegetation changes will result in an overall positive feedback on the climate.

Recent changes and results of climate change simulation experiments in the field have shown that there are considerable uncertainties in the projected rates of change e.g. (Van Bogaert *et al.* 2010). Furthermore, the models do not yet include vertebrate and invertebrate herbivory, extreme events such as tundra fire and extreme winter warming damage or changes in land use that either reduce the rate of vegetation change or open up niches for rapid change. Projections suggest increases in the ranges of the autumn and winter moths that have outbreaks in populations resulting in the defoliation of birch forest (Jepsen *et al.*, 2008 and 2011), and a general increase in the “background” (non-outbreak) invertebrate herbivores (Wolf *et al.*, 2008).

Animal terrestrial biodiversity is generally projected to increase in the Arctic during warming by immigration of new species from the south, vegetation changes, and indirectly by introduction of invasive species caused by increased human activities and increased survival of such species (*high confidence*) (Post *et al.*, 2009; Gilg *et al.*, 2012; CAFF 2013). Many native Arctic species will *most likely* be increasingly threatened during this century.

28.3.3.2. Antarctica

Projected effects of climate change on Antarctic terrestrial species are limited to knowledge of their ecophysiological tolerances to changes in air temperature, wind speed, precipitation (rain and snowfall), permafrost thaw and exposure of new habitat through glacial/ice retreat. The climate is expected to become more tolerable to a number of species, leading to increases in biomass and extent of existing ecological communities.

The frequency with which new potential colonising plant and animal species arrive in Antarctica (particularly the Antarctic Peninsula region) from lower latitudes, and the subsequent probability of their successful establishment will increase with regional climate warming and associated environmental changes (Chown *et al.*, 2012) (*high confidence*). Human-assisted transfers of biota may be more important by two orders of magnitude than natural introductions (Frenot *et al.*, 2005) as the transfer is faster and avoids extreme environments such as altitude or oceans (Barnes *et al.*, 2006). The potential for anthropogenic introduction of non-indigenous species to Antarctic terrestrial areas, which could have devastating consequences to the local biodiversity, will increase (Convey *et al.*, 2009; Convey, 2011, Hughes and Convey, 2010; Braun *et al.*, 2012) (*high confidence*). At present, established non-indigenous species in the sub- and maritime Antarctic are very restricted in their distributions (Frenot *et al.*, 2005). Climate change could result in a greater rate of spread of invasive species through colonisation of areas exposed by glacial retreat, as has occurred at South Georgia (Cook *et al.*, 2010) and in the maritime Antarctic (Olech and Chwedorzewska 2011). Biosecurity measures may be needed to help control dispersal of established non-indigenous species to new locations, particularly given the expected increase in human activities in terrestrial areas (Hughes and Convey, 2010; Convey *et al.*, 2011). An important gap in understanding is the degree to which climate change may facilitate some established but localised alien species to become invasive and widespread (Frenot *et al.*, 2005; Convey 2010; Hughes and Convey 2010; Cowan *et al.*, 2011), which has been shown for the sub-Antarctic (Chown *et al.*, 2012).

Overall, the likely impacts of existing and new non-indigenous species on the native terrestrial ecosystems of Antarctica and the sub-Antarctic islands, along with the continued increased presence of Antarctic fur seals are likely to have far greater importance over the timescale under consideration than are those attributable to climate change itself (Turner *et al.*, 2009; Convey and Lebouvier, 2009; Convey, 2010).

28.3.4. Economic Sectors

Projections of economic costs of climate change impacts for different economic sectors in the Arctic are limited, but current assessments suggest that there will be both benefits and costs (Forbes, 2011; AMAP, 2011). Non-Arctic actors are likely to receive most of the benefits from increased shipping and commercial development of renewable and non-renewable resources, while indigenous peoples and local Arctic communities will have a harder time maintaining their way of life (Hovelsrud *et al.*, 2011).

Contributing to the complexity of measuring the future economic effects of climate change is the uncertainty in future predictions and the rapid speed of change, which are linked with the uncertainty of the technological and ecological effects of such change (NorAcia, 2010). Communities within the same eco-zone may experience different effects from identical climate-related events because of marked local variations in site, situation, culture and economy (Clark *et al.*, 2008).

Economic cost estimates have been made for the case of the Alaskan economy, for example, which suggest that a heavy reliance on climate-sensitive businesses such as tourism, forestry, and fisheries, renders the economy vulnerable to climate change, and that Alaska Native peoples, reliant on the biodiversity of the Alaskan ecosystem, are being affected disproportionately (Epstein and Ferber, 2011). Some Alaskan villages such as Shishmaref, Kivalina, and Newtok have already lost critical infrastructure and services and are becoming unlivable due to storm damage and coastal erosion but the high costs and limitations of government mechanisms are significant barriers to the actual relocation of these communities (Bronen, 2011; Cochran *et al.*, 2013; Maldonado *et al.*, 2013).

28.3.4.1. Fisheries

Climate change will impact the spatial distribution and catch of some open ocean fisheries in the Barents and Bering Seas (*high certainty*). The future of commercial fisheries in Arctic Ocean is uncertain. There is strong evidence and considerable data showing links between climate driven shifts in ocean conditions and shifts in the spatial distribution and abundance of commercial species in the Bering and Barents Seas (Section 28.3.2.2.1). In limited cases, coupled bio-physical models or climate enhanced stock projection models have been used to predict future commercial yield or shifts in fishing locations. However, these predictions are uncertain (Huse and Ellingsen, 2008; Ianelli *et al.*, 2011; Wilderbuer *et al.* 2012). Cheung *et al.* (2011) used projections from an earth system model to estimate shifts in bio-climatic windows that included climate change effects on biogeochemistry (oxygen and acidity) and primary production to project future catch potential of 120 demersal fish and invertebrates. Results from their model suggested that the catch potential will increase in the Barents and Greenland Seas and regions > 70° north latitude (Cheung *et al.*, 2011). In contrast, vulnerability analysis suggests that only a few species are expected to be abundant enough to support viable fisheries in the Arctic Ocean (Hollowed *et al.*, 2013). Potential fisheries for snow crab on shelf areas of the Arctic Ocean may be limited by the associated impacts of ocean acidification. If fisheries develop in the Arctic Ocean, adoption of sustainable strategies for management will be a high priority (Molenaar, 2009). The moratorium on fishing in the U.S. portion of the Chukchi and Beaufort Seas would prevent fishing until sufficient data become available to manage the stock sustainably (Wilson and Ormseth, 2009).

Predicting of how harvesters will respond to changing economic, institutional and environmental conditions under climate change is difficult. Current techniques, track fishers choices based on revenues and costs associated with targeting a species in a given time and area with a particular gear given projected changes in the abundance and spatial distribution of target species (Haynie and Pfeiffer, 2012). However, estimates of future revenues and costs will depend, in part, on future demand for fish, global fish markets and trends in aquaculture practices (Merino *et al.*, 2012; Rice and Garcia, 2011).

28.3.4.2. Forestry and Farming

Climate change is likely to have positive impacts for agriculture, including extended growing season (Grønlund, 2009; Falloon and Betts, 2009; Tholstrup and Rasmussen, 2009), although variations across regions are expected (Hovelsrud *et al.*, 2011), and the importance of impacts to the Arctic economy will likely remain minor (Eskeland and Flottorp, 2006). Potential positive effects of climatic warming for forestry include decreased risk of snow damage. Kilpelainen *et al.* (2010) estimate a 50% decrease in snow damage in Finland towards the end of the century. A warmer climate is likely to impact access conditions and plant diseases for forestry and farming. Grønlund (2009) found in the case of Northern Norway – where about half of the arable land area is covered by forest and 40% by marshland – that the potential harnessing of arable land for farming will be at the cost of forestry production, or dried-up marshlands, which may contribute to more greenhouse emissions. Larger field areas may contribute to land erosion through rainfall and predicted unstable winters, and may increase conditions for plant diseases and fungal infections (Grønlund, 2009). If the winter season continues to shorten due to climate change (Xu *et al.*, 2013), accessibility to logging sites will be negatively affected. Accessibility is higher when frozen ground makes transportation possible in sensitive locations or areas that lack road. If weather changes occur when logging has taken place, sanding of roads may be necessary which carries significant economic costs. Impact on carrying capacity of ground or road accessibility will thus affect forestry economically. Challenges may include limited storage space for wood (Keskitalo, 2008).

28.3.4.3. Infrastructure, Transportation, and Terrestrial Resources

Rising temperatures and changing precipitation patterns have the potential to affect all infrastructure types and related services, as much of the infrastructure in the North is dependent upon the cryosphere to, for example, provide stable surfaces for buildings and pipelines, contain waste, stabilize shorelines and provide access to remote communities in the winter (Furgal and Prowse, 2008; Huntington *et al.*, 2007; Sherman *et al.*, 2009; Sundby and Nakken, 2008; West and Hovelsrud, 2010; Forbes, 2011). In the long term marine and freshwater transportation will need to shift its reliance from ice routes to open-water or land-based transportation systems. Of appropriate community adaptations to the predicted changes relocation is one option to deal with persistent flooding and bank erosion (Furgal, 2008; NRTEE, 2009). Changing sea-ice (multiyear) conditions are suspected i.e. to have a regulating impact on marine shipping and coastal infrastructure through possible hazards on them (Eicken *et al.*, 2009).

By adapting transportation models to integrate monthly climate model (CCSM3) predictions of air temperature, combined with datasets on land cover, topography, hydrography, built infrastructure, and locations of human settlements, estimates have been made of changes to inland accessibility for northern landscapes northward of 40°N by mid-21st Century (Stephenson *et al.*, 2011). Milder air temperatures and/or increased snowfall reduce the possibilities for constructing inland winter-road networks, including ice roads, with the major seasonal reductions in road potential (based on a 2000 kg vehicle) being in the winter shoulder-season months of November and April. The average decline (compared to a baseline of 2000-2014) for eight circumpolar countries was projected to be -14%, varying from -11 to -82%. In absolute terms, Canada and Russia (both at -13%) account for the majority of declining winter-road potential with $\sim 1 \times 10^6$ km² being lost (See Table 28.1). The winter road season has decreased since the 1970s on the Alaska's North Slope, from as much as 200 to 100 days in some areas (Hinzman, *et al.*, 2005).

Climate change is expected to lead to a nearly ice free Arctic Ocean in late summer and increased navigability of Arctic marine waters. New possibilities for shipping routes and extended use of existing routes may result from increased melting of sea ice (Paxian *et al.*, 2010; Corbett *et al.*, 2010; Khon *et al.*, 2010; Peters *et al.*, 2011; Stephenson *et al.*, 2011).

Projections made by Stephenson *et al.* (2011) suggest that all five Arctic littoral states will gain increased maritime access to their current exclusive economic zones, especially Greenland (+28% relative to baseline), Canada (+19%), Russia (+16%) and United States (+15%). In contrast, Iceland, Norway, Sweden, and Finland display little or no increase in maritime accessibility, (Table 28.1) (Stephenson *et al.*, 2011).

GCMs generally underestimate the duration of the ice-free period in the Arctic Ocean and simulate slower changes than those observed in the past decades (Stroeve *et al.*, 2007). Mokhow and Khon (2008) used a sub-set of climate models that better than other GCMs reproduce the observed sea ice dynamics to project the duration of the navigation season along the NSR and through the NWP under the moderate SRES-A1B emission scenario. According to their results, by the end of the 21st century the NSR may be open for navigation 4.5±1.3 months per year, while the NWP may be open 2-4 months per year (Figure 28-4). The models did not predict any significant changes of the ice conditions in the NWP until the early 2030s.

[INSERT FIGURE 28-4 HERE

Figure 28-4: Projected duration of the navigation period (days) over the Northwest Passage and Northern Sea Route. Source: Mokhow and Khon, 2008.]

[INSERT TABLE 28-1 HERE

Table 28-1: Annually averaged changes in inland and maritime transportation accessibility by mid-century (2045–2059) versus baseline (2000–2014). Source: Stephenson *et al.*, 2011.]

An increase in the length of the summer shipping season, with sea-ice duration expected to be 10 days shorter by 2020 and 20-30 days shorter by 2080, is likely to be the most obvious impact of changing climate on Arctic marine transportation (Prowse *et al.*, 2009). Reduction in sea ice and increased marine traffic could offer opportunities for economic diversification in new service sectors supporting marine shipping. Loss of sea ice may open up waterways and opportunities for increased cruise traffic (e.g. Glomsrød and Aslaksen, 2009), and add to an already rapid increase in cruise tourism (Stewart *et al.*, 2010; Stewart *et al.*, 2007; Howell *et al.*, 2007). Climate change has increased the prevalence of cruise tourism throughout Greenland, Norway, Alaska and Canada because of decreasing sea ice extent.

Projected declines in sea-ice covers leading to development of integrated land and marine transportation networks in Northern Canada may stimulate further mine exploration and development (Prowse *et al.*, 2009). These possibilities however also come with challenges including their predicted contribution to the largest change in contaminant movement into or within the Arctic, as well as their significant negative impacts on the traditional ways of life of northern residents (Furgal and Prowse, 2008). Added shipping and economic activity will increase the amount of black carbon and reinforce warming trends in the region (Lack and Corbett, 2012), leading to additional economic activity.

Longer shipping season and improved access to ports may lead to increased petroleum activities, although possible increased wave activity and coastal erosion may increase costs related to infrastructure and technology. Peters *et al.* (2011) find by using a bottom-up shipping model and a detailed global energy market model to construct emission inventories of Arctic shipping and petroleum activities in 2030 and 2050 – and based on estimated sea-ice extent – that there will be rapid growth in transit shipping; oil and gas production will be moving into locations requiring more ship transport; and this will lead to rapid growth in emissions from oil and gas transport by ship.

The Arctic contains vast resources of oil, which is hard to replace as transportation fuel, and vast resources of gas, a more climate benign fuel than coal. Petroleum resources are unevenly distributed among Arctic regions and states. Arctic resources will play a growing role in the world economy, but increased accessibility is expected to create challenges for extraction, transport, engineering, search-and-rescue needs and responses to accidents (Hovelsrud *et al.*, 2011), and climatic change presents the oil and gas industry with challenges in terms of planning and predictions (Harsem *et al.*, 2011). Increased emissions due to rapid growth in Arctic Ocean transportation of oil and gas are projected (Peters *et al.*, 2011). Due to high costs and difficult access conditions the impact on future oil and gas production in the Arctic remains unclear (Peters *et al.*, 2011; Lindholdt and Glomsrød, 2012).

28.4. Human Adaptation

There is general agreement that both indigenous and non-indigenous people in the Arctic have a history of adapting to natural variability in the climate and natural resource base, as well as recent socio-economic, cultural and technological changes (Forbes and Stammer, 2009; Wenzel, 2009; West and Hovelsrud, 2010; Ford and Pearce, 2010; Bolton *et al.*, 2011; Cochran *et al.*, 2013). Climate change exacerbates the existing stresses faced by Arctic communities (Rybråten and Hovelsrud, 2010; Crute and Nuttall, 2009) and is only one of many important factors influencing adaptation (Berrang-Ford *et al.*, 2011). Climate adaptation needs to be seen in the context of these interconnected and mutually reinforcing factors (Tyler *et al.*, 2007; Hovelsrud and Smit, 2010). The challenges faced today by communities in the Arctic are complex and interlinked and are testing their traditional adaptive capacity.

Climatic and other large-scale changes have potentially large effects on Arctic communities, in particular where simple economies leave a narrower range of adaptive choices (Anisimov and Vaughan, 2007; Ford and Furgal, 2009; Andrachuk and Pearce, 2010; Ford *et al.*, 2010; Forbes, 2011; Berkes *et al.*, 2003). There is considerable evidence that changing weather patterns, declining sea-ice and river as well as lake ice, thawing permafrost, and plant and animal species' abundance and composition have consequences for communities in the Arctic (see 28.2.4, 28.2.5.2 and 28.3.4). Sea-ice is particularly important for coastal communities which rely upon it for transportation between communities and hunting areas (Krupnik *et al.*, 2010). Changes in the duration and condition of sea ice and the consequent changes to country food availability significantly impact the wellbeing of communities (Furgal and Seguin, 2006; Ford and Berrang-Ford, 2009; Ford *et al.*, 2010), outdoor tourism (Dawson *et al.*, 2010) and hunting and fishing (Wiig *et al.*, 2008; Brander, 2010).

Adaptation to climate change is taking place at the local and regional levels where impacts are often felt most acutely and the resources most readily available (Oskal, 2008; Hovelsrud and Smit, 2010). Current experiences and projections of future conditions often lead to technological adaptation responses such as flood and water management and snow avalanche protection (West and Hovelsrud, 2010; Hovelsrud and Smit, 2010) rather than policy responses (Hedensted Lund *et al.*, 2012; Rudberg *et al.*, 2012). Climate variability and extreme events are found to be salient drivers of adaptation (Berrang-Ford *et al.*, 2011; Dannevig *et al.*, 2012; Amundsen *et al.*, 2010).

The lack of local scale climate projections, combined with uncertainties in future economic, social and technological developments often act as barriers to adaptation. These barriers, together with other societal determinants such as ethics, cultures, and attitudes towards risk may cause inaction (Adger *et al.*, 2009; West and Hovelsrud, 2010). Resolving divergent values across and within different communities poses a challenge for governance regimes. A determining factor in building adaptive capacity is the flexibility of enabling institutions to develop robust options (Keskitalo *et al.*, 2009; Hovelsrud and Smit 2010; Forbes *et al.*, 2009; Ford and Goldhar, 2012; Whyte, 2013). In the North American and Scandinavian context, adaptive co-management responses have been developed through land claims settlements and/or multi-scale institutional cooperation to foster social learning (Berkes, 2009; Armitage *et al.*, 2008).

Indigenous Peoples

While Arctic indigenous peoples with traditional lifestyles are facing unprecedented impacts to their ways of life from climate change and resource development (oil and gas, mining, forestry, hydropower, tourism, etc.), they are already implementing creative ways of adapting (Cruikshank, 2001; Forbes *et al.*, 2006; Krupnik and Ray, 2007; Salick and Ross, 2009; Green and Raygorodetsky, 2010; Cullen-Unsworth *et al.*, 2011; Alexander *et al.*, 2011; Bongo *et al.*, 2012). Examples of indigenous adaptation strategies have included changing resource bases, shifting land use and/or settlement areas, combining technologies with traditional knowledge, changing timing and location of hunting, gathering, herding, and fishing areas, and improving communications and education (Galloway McLean, 2010; Bongo *et al.*, 2012). Protection of grazing land will be the most important adaptive strategy for reindeer herders under climate change (Forbes *et al.*, 2009; Magga *et al.*, 2011; Kumpula *et al.*, 2012; Degteva and Nellemann, 2013).

The adaptive capacity of Arctic indigenous peoples is largely due to an extensive traditional knowledge and cultural repertoire, and flexible social networks (see Chapter 12, section 12.3) (Williams and Hardison, 2013). The dynamic nature of traditional knowledge is valuable for adapting to current conditions (Kitti *et al.*, 2006; Tyler *et al.*, 2007; Eira *et al.*, 2012). The sharing of knowledge ensures rapid responses to crises (Ford *et al.*, 2007). In addition, cultural values such as sharing, patience, persistence, calmness, respect for elders and the environment are important. Some studies suggest that traditional knowledge may not always be sufficient to meet the rapid changes in climate (see also Chapter 12) and it may be perceived to be less reliable because the changing conditions are beyond the current knowledge range (Ingram *et al.*, 2002; Ford *et al.*, 2006; Valdivia *et al.*, 2010; Hovelsrud *et al.*, 2010).

Over the last half-century, the adaptive capacity in some indigenous communities has been challenged by the transition from semi-nomadic hunting groups to permanent settlements (Ford *et al.*, 2010). Forced or voluntary migration as an adaptation response can have deep cultural impacts (Shearer, 2011,2012; Maldonado *et al.*, 2013). The establishment of permanent communities, particularly those associated with new industrial development, can also lead to increasing employment opportunities and income diversification for indigenous peoples. The intergenerational transfers of knowledge and skills through school curricula, land camps, and involvement in community-based monitoring programmes may strengthen adaptive capacity (Bolton *et al.*, 2011; Hovelsrud and Smit, 2010; Ford *et al.*, 2007; Forbes 2007).

Renewable resource harvesting remains a significant component of Arctic livelihoods and with climate change hunting and fishing has become a riskier undertaking and many communities are already adapting (Gearheard *et al.*, 2011; Laidler *et al.*, 2011). Adaptation includes taking more supplies when hunting; constructing permanent shelters on land as refuges from storms; improved communications infrastructure; greater use of global positioning systems (GPS) for navigation; synthetic aperture radar (SAR) to provide estimates of sea-ice conditions (Laidler *et al.*, 2011) and the use of larger or faster vehicles (Ford *et al.*, 2010). Avoiding dangerous terrain can result in longer and time-consuming journeys which can be inconvenient to those with wage-earning employment (Ford *et al.*, 2007).

Reindeer herders have developed a wide range of adaptation strategies in response to changing pasture conditions. These include: moving herds to better pastures (Bartsch *et al.*, 2010); providing supplemental feeding (Helle and Jaakkola, 2008; Forbes and Kumpula, 2009); retaining a few castrated reindeer males to break through heavy ice-crust (Oskal, 2008; Reinert *et al.*, 2008); ensuring an optimal herd size (Forbes *et al.*, 2009; Tyler *et al.*, 2007); and creating multicultural initiatives combining traditional knowledge with scientific (Vuojala-Magga *et al.*, 2011; Bongo *et al.*, 2012). Coastal fishers have adapted to changing climate by targeting different species and diversifying income sources (Hovelsrud *et al.*, 2010).

In some Arctic countries indigenous peoples have successfully negotiated land claims rights and have become key players in addressing climate change (Abele *et al.*, 2009). In some instances this has given rise to tensions over land/water use between traditional livelihoods and new opportunities (e.g. tourism and natural resource development) (Forbes *et al.*, 2006; Hovelsrud and Smit, 2010). Some territorial governments in Northern Canada have promoted adaptation by providing hunter support programs (Ford *et al.*, 2006, 2010).

The health of many indigenous people is being affected by the interaction of changes in the climate with ongoing changes in human, economic and biophysical systems (Donaldson *et al.*, 2010). The distribution of traditional foods between communities and the use of community freezers in the Canadian Arctic has improved food security, an important factor for health (Ford *et al.*, 2010). While wage employment may enhance the possibilities for adaptive capacity, greater involvement in full time jobs can threaten social and cultural cohesion and mental well-being by disrupting the traditional cycle of land-based practices (Berner *et al.*, 2005; Furgal, 2008).

28.5. Research and Data Gaps

There remains a poor knowledge of coupling among, and thresholds within, bio-geophysical and socio-economic processes to fully assess the effects of a changing climate, and to separate them from those due to other environmental stressors:

- Existing integrative models are either lacking or insufficiently validated to project and to assess the cascading effects on, and feedbacks from the systems in the Polar Regions, in particular socio-economic systems.
- There is a need to enhance or establish a coordinated network of long-term representative sites for monitoring and assessment of climate change detection and attribution studies in Polar Regions. Regional differences and confounding variables will need to be considered in designing field and modelling studies. Standardised methods and approaches of biophysical and socio-economic analysis along with coordinated sampling in more regions will be necessary.

There are more specific research gaps, including:

- Many mechanisms of how climate change and ocean acidification may be affecting polar ecosystems have been proposed but few studies of physiological tolerances of species, long term field studies of ecosystem effects and ecosystem modelling studies are available to be able to attribute with high confidence current and future change in these ecosystems to climate change.
- More comprehensive studies including long-term monitoring on the increasing impacts from climate changes on Arctic communities (urban and rural) and their health, well-being, traditional livelihoods and life ways are needed. There is a need to assess more fully vulnerabilities and to develop response capacities at the local and regional level.

Frequently Asked Questions

FAQ 28.1: What will be the net socio-economic impacts of change in the polar regions?

[to remain at the end of the chapter]

Climate change will have costs and benefits for Polar Regions. Climate change, exacerbated by other large-scale changes, can have potentially large effects on Arctic communities, where relatively simple economies leave a narrower range of adaptive choices.

In the Arctic, positive impacts include new possibilities for economic diversification, marine shipping, agricultural production, forestry, and tourism. The Northern Sea Route is predicted to have up to 125 days per year suitable for navigation by 2050, while the heating energy demand in the populated Arctic areas is predicted to decline by 15%. In addition, there could be greater accessibility to offshore mineral and energy resources although challenges related to environmental impacts and traditional livelihoods are possible.

Changing sea ice condition and permafrost thawing may cause damage to bridges, pipelines, drilling platforms, hydropower and other infrastructure. This poses major economic costs and human risks, although these impacts are closely linked to the design of the structure. Furthermore, warmer winter temperatures will shorten the accessibility of ice roads that are critical for communications between settlements and economic development and have implications for increased costs.. Statistically, a long-term mean increase of 2 to 3°C in autumn and spring air temperature produces an approximate 10 to 15 day delay in freeze-up and advance in break-up, respectively.

Particular concerns are associated with projected increase in the frequency and severity of ice-jam floods on Siberian rivers. They may have potentially catastrophic consequences for the villages and cities located in the river plain, as exemplified by the 2001 Lena River flood, which demolished most of the buildings in the city of Lensk.

Changing sea ice conditions will impact indigenous livelihoods, and changes in resources, including marine mammals, could represent a significant economic loss for many local communities. Food security and health and well-being are expected to be impacted negatively.

In the Antarctic, tourism is expected to increase, and risks exist of accidental pollution from maritime accidents, along with an increasing likelihood of the introduction of alien species to terrestrial environments. Fishing for Antarctic krill near to the Antarctic continent is expected to become more common during winter months in areas where there is less winter sea ice.

FAQ 28.2: Why are changes in sea ice so important to the polar regions? *[to remain at the end of the chapter]*

Sea ice is a dominant feature of Polar Oceans. Shifts in the distribution and extent of sea ice during the growing season impacts the duration, magnitude and species composition of primary and secondary production in the Polar Regions. With less sea ice many marine ecosystems will experience more light, which can accelerate the growth of phytoplankton, and shift the balance between the primary production by ice algae and water-borne phytoplankton,

with implications for Arctic food webs. In contrast, sea ice is also an important habitat for juvenile Antarctic krill, providing food and protection from predators. Krill is a basic food source for many species in polar marine ecosystems.

Changes in sea ice will have other impacts, beyond these “bottom-up” consequences for marine foodwebs. Mammals and birds utilize sea ice as haul-outs during foraging trips (seals, walrus, and polar bears in the Arctic and seals and penguins in the Antarctic). Some seals (e.g. Bearded seals in the Arctic and crabeater and leopard seals in the Antarctic) give birth and nurse pups in pack ice. Shifts in the spatial distribution and extent of sea ice will alter the spatial overlap of predators and their prey. According to model projections, within 50-70 years loss of hunting habitats may lead to elimination of polar bears from seasonally ice-covered areas, where two thirds of their world population currently live. The vulnerability of marine species to changes in sea ice will depend on the exposure to change, which will vary by location, as well as the sensitivity of the species to changing environmental conditions and the adaptive capacity of each species. More open waters and longer ice-free period in the northern seas enhance the effect of wave action and coastal erosion with implications for coastal communities and infrastructure.

While the overall sea ice extent in the Southern Ocean has not changed markedly in recent decades, there have been increases in oceanic temperatures and large regional decreases in winter sea ice extent and duration in the western Antarctic Peninsula region of West Antarctica and the islands of the Scotia Arc.

References

- Aaheim, A., H. Dannevig, T. Ericsson, B. van Oort, L. Innbjør, T. Rauken, H. Vennemo, H. Johansen, M. Tofteng, C. Aall, K. Groven and E. Heiberg, 2009. Konsekvenser av klimaendringer, tilpasning og sårbarhet i Norge – Rapport til Klimatilpasningsutvalget. [in Norwegian]. CICERO Report. 4:238 pp.
- Abele, F., T. J. Courchene, F.L. Sidele, and F. St-Hilaire, 2009: Northern Exposure: Peoples, Powers and Prospects in Canada's North. Montreal: The Institute for Research on Public Policy. xii, 605 p. : ill., Montréal, Québec
- Abryutina, L.I., 2009: Indigenous peoples of the Russian North: Social and Climatic changes. In: *Climate Change and Arctic Sustainable Development: Scientific, Social Cultural, and educational challenges*. UNESCO, Paris, pp. 164-173.
- Adam, J.C., A.F. Hamlet, and D.P. Lettenmaier, 2009: Implications of global climate change for snowmelt hydrology in the twenty-first century. *Hydrological Processes*, **23(7)**, 962-972.
- Adger, W., S. Dessai, M. Goulden, M. Hulme, I. Lorenzoni, D. Nelson, L. Naess, J. Wolf, and A. Wreford, 2009: Are there social limits to adaptation to climate change? *Climatic Change*, **93(3)**, 335-354.
- Albrecht, G., G. Sartore, L. Connor, N. Higginbotham, S. Freeman, B. Kelly, H. Stain, A. Tonna, and G. Pollard, 2007: Solastalgia: the distress caused by environmental change. *Australasian Psychiatry*, **15**, S95-S98.
- Alexander, C., N. Bynum, E. Johnson, U. King, T. Mustonen, P. Neofotis, N. Oetttlé, C. Rosenzweig, C. Sakakibara, V. Shadrin, M. Vicarelli, Jon Waterhouse, and B. Weeks, 2011: Linking Indigenous and Scientific Knowledge of Climate Change. *Bioscience*, **61(6)**, pp. 477-484.
- AMAP, 2011. Arctic Pollution 2011. Arctic Monitoring and Assessment Programme (AMAP). Oslo, Norway. Vi + 38 pp.
- AMAP, 2011: *Snow, Water, Ice and Permafrost in the Arctic (SWIPA): Climate Change and the Cryosphere*. Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway. xii + 538 pp.
- AMAP. 2009. AMAP Assessment 2009: Human Health in the Arctic. Arctic Monitoring and Assessment Programme (AMAP). Oslo, Norway. Xii + 254 pp.
- AMAP. 2010. AMAP Assessment 2009: Radioactivity in the Arctic. Arctic Monitoring and Assessment Programme (AMAP). Oslo, Norway. Xii + 92.
- Amstrup, S.C., B.G. Marcot, and D.C. Douglas, 2008: A Bayesian network approach to forecasting the 21st century worldwide status of polar bears. In: *Arctic Sea Ice Decline: Observations, Projections, Mechanisms, and Implications*. [DeWeaver, E.T., C.M. Bitz, and L.B. Tremblay (eds.)]. AGU, Washington, D. C., pp. 213.
- Amstrup, S.C., E.T. DeWeaver, D.C. Douglas, B.G. Marcot, G.M. Durner, C.M. Bitz, and D.A. Bailey, 2010: Greenhouse gas mitigation can reduce sea-ice loss and increase polar bear persistence. *Nature*, **468(7326)**, 955-U351
- Amstrup, S.C., H. Caswell, E. DeWeaver, I. Stirling, D.C. Douglas, B.G. Marcot, and C.M. Hunter, 2009: Rebuttal of "Polar Bear Population Forecasts: A Public-Policy Forecasting Audit". *Interfaces*, doi:10.1287/inte.1090.0444.

- Amstrup, S.C., I. Stirling, T.S. Smith, C. Perham, and G.W. Thiemann, 2006: Recent observations of intraspecific predation and cannibalism among polar bears in the southern Beaufort Sea. *Polar Biology*, **29(11)**, 997-1002.
- Amundsen, Helene, Frode Viktor Berglund and Hege Westskog, 2010. Overcoming barriers to climate change adaptation – a question of multilevel governance?. *Environment and Planning C*, 28 (2): pp. 276-289.
- Andrachuk, M. and T. Pearce, 2010: Vulnerability and Adaptation in Two Communities in the Inuvialuit Settlement Region. In: *Community Adaptation and Vulnerability in Arctic Regions*. [Hovelsrud, G.K. and B. Smit(eds.)]. Springer Netherlands, pp. 63-81.
- Anisimov, O. A., Zhil'tsova, E. L. and Reneva, S. A. 2011. Estimation of critical levels of climate change influence on the natural terrestrial ecosystems on the territory of Russia. *Meteorology and Hydrology* 36, 723-730 DOI: 10.3103/S1068373911110033
- Anisimov, O. and D. Vaughan, 2007: Polar regions. In: *Climate Change 2007:Impacts, adaptation, and vulnerability. Contribution of Working group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, pp. 653-686.
- Arctic Council, 2013. *Arctic Resilience Interim Report 2013*. Stockholm Environment Institute and Stockholm Resilience Centre, Stockholm.
- Armitage D., 2008: Governance and the commons in a multi-level world. *International Journal of the Commons*, **2(1)**
- Armitage, J.M., C.L. Quinn, and F. Wania. 2011. Global Climate change and contaminants – an overview of opportunities and priorities for modeling the potential implications for long-term human exposure to organic compounds in the Arctic. *J. Environ. Monit.* 13, 1532. Pp. 1532-1546.
- Armstrong, J.S., K.C. Green, and W. Soon, 2008: Polar Bear Population Forecasts: A Public-Policy Forecasting Audit. *Interfaces*, **38(5)**, 382-395.
- Arnason, R., 2012: Global warming: New challenges for the common fisheries policy? *Ocean & Coastal Management*, **70**, 4-9.
- Arrigo, K., G. van Dijken, and S. Bushinsky, 2008: Primary production in the Southern Ocean, 1997-2006. *Journal of Geophysical Research-Oceans*, **113**, [C8].
- Arrigo, K.R., and G. L. van Dijken, 2011: Secular trends in Arctic Ocean net primary production. *Journal of Geophysical Research*, **116**, C09011.
- Arrigo, K.R., D.K. Perovich, R.S. Pickart, Z.W. Brown, G.L. van Dijken, K.E. Lowry, M.M. Mills, M.A. Palmer, W.M. Balch, F. Bahr, N.R. Bates, C. Benitez-Nelson, B. Bowler, E. Brownlee, J.K. Ehn, K.E. Frey, R. Garley, S.R. Laney, L. Lubelczyk, J. Mathis, A. Matsuoka, B.G. Mitchell, G.W.K. Moore, E. Ortega-Retuerta, S. Pal, C.M. Polashenski, R.A. Reynolds, B. Schieber, H.M. Sosik, M. Stephens, and J.H. Swift, 2012: Massive Phytoplankton Blooms Under Arctic Sea Ice. *Science*, **336(6087)**, 1408-1408.
- Astthorsson, O. S., H. Valdimarsson, A. Gudmundsdottir, G. J. Ó'skarsson. 2013: Climate-related variations in the occurrence and distribution of mackerel (*Scomber scombrus*) in Icelandic waters. *ICES Journal of Marine Science*, **69(7)**, 1289–1297.
- Atkinson, A., V. Siegel, E. Pakhomov, and P. Rothery, 2004: Long-term decline in krill stock and increase in salps within the Southern Ocean. *Nature*, 2996(Letters to Nature), 1-4.
- Bajzak, C.E., M.O. Hammill, G.B. Stenson, and S. Prinsenberg, 2011: Drifting away: Implications of changes in ice conditions for a pack-ice-breeding phocid, the harp seal (*Pagophilus groenlandicus*). *Canadian Journal of Zoology*, **89(11)**, 1050-1062.
- Bakun, A., 2010: Linking climate to population variability in marine ecosystems characterized by non-simple dynamics: Conceptual templates and schematic constructs. *Journal of Marine Systems*, **79(3–4)**, 361-373.
- Balayla, D., T. Lauridsen, M. S ndergaard, and E. Jeppesen, 2010: Larger zooplankton in Danish lakes after cold winters: are winter fish kills of importance? *Hydrobiologia*, **646(1)**, 159-172.
- Barber, D.G., J.V. Lukovich, J. Keogak, S. Baryluk, L. Fortier, and G.H.R. Henry, 2008: The Changing Climate of the Arctic. *Arctic*, **61**, 7-26.
- Barbraud, C., C. Marteau, V. Ridoux, K. Delord, and H. Weimerskirch, 2008: Demographic response of a population of white-chinned petrels *Procellaria aequinoctialis* to climate and longline fishery bycatch. *Journal of Applied Ecology*, **45(5)**, 1460-1467.
- Barbraud, C., P. Rivalan, P. Inchausti, M. Nevoux, V. Rolland, and H. Weimerskirch, 2011: Contrasted demographic responses facing future climate change in Southern Ocean seabirds. *Journal of Animal Ecology*, **80(1)**, 89-100.

- Barbraud, C., V. Rolland, S. Jenouvrier, M. Nevoux, K. Delord, and H. Weimerskirch, 2012: Effects of climate change and fisheries bycatch on Southern Ocean seabirds: a review. *Marine Ecology Progress Series*, **454**, 285-307.
- Barnes, D.K.A. and T. Souster, 2011: Reduced survival of Antarctic benthos linked to climate-induced iceberg scouring. *Nature Climate Change*, **1(7)**, 365-368.
- Barnes, D.K.A., D.A. Hodgson, P. Convey, C.S. Allen, and A. Clarke, 2006: Incursion and excursion of Antarctic biota: past, present and future. *Global Ecology and Biogeography*, **15(2)**, 121-142.
- Barnes, D.K.A., D.A. Hodgson, P. Convey, C.S. Allen, and A. Clarke, 2006: Incursion and excursion of Antarctic biota: past, present and future. *Global Ecology and Biogeography*, **15(2)**, 121-142
- Bartsch, A., T. Kumpula, B.C. Forbes, and F. Stammer, 2010: Detection of snow surface thawing and refreezing in the Eurasian Arctic with QuikSCAT: implications for reindeer herding. *Ecological Applications*, **20(8)**, 2346-2358.
- Beaugrand, G. and R.R. Kirby, 2010: Spatial changes in the sensitivity of Atlantic cod to climate-driven effects in the plankton. *Climate Research*, **41(1)**, 15-19.
- Bednarsek, N., G.A. Tarling, D.C.E. Bakker, S. Fielding, E.M. Jones, H.J. Venables, P. Ward, A. Kuzirian, B. Leze, R.A. Feely, and E.J. Murphy, 2012: Extensive dissolution of live pteropods in the Southern Ocean. *Nature Geoscience*, **5(12)**, 881-885.
- Beltaos, S. and T.D. Prowse, 2009. River ice hydrology in a shrinking cryosphere. *Hydrological Processes*, 23(1): 122-144.
- Berg, T.B., N.M. Schmidt, T.T. Høye, P.J. Aastrup, D.K. Hendrichsen, M.C. Forchhammer, and D.R. Klein, 2008: High-Arctic Plant-Herbivore Interactions under Climate Influence. *Advances in Ecological Research*, **40**, 275-298.
- Berg**, T.B., N.M. Schmidt., T.T. Høye., P.J. Aastrup., D.K. Hendrichsen., M.C. Forchhammer, and D.R. Klein, 2008: High-Arctic Plant-Herbivore Interactions Under Climate Influence. *Advances in Ecological Research*, **40**, 275-298.
- Berkes, F. and D. Armitage, 2010: Co-Management Institutions, Knowledge and Learning: Adapting to Change in the Arctic. *Etudes Inuit Studies*, **34(1)**, 109.
- Berkes, F., 2009: Evolution of co-management: Role of knowledge generation, bridging organizations and social learning. *Journal of Environmental Management*, **90(5)**, 1692-1702.
- Berkman, P.A. and Royal United Services Institute for Defence and Security Studies., 2010: *Environmental security in the Arctic Ocean : promoting co-operation and preventing conflict*. Published on behalf of The Royal United Services Institute for Defence and Security Studies by Routledge Journals, Abingdon, .
- Berrang-Ford, L., J.D. Ford, and J. Paterson, 2011: Are we adapting to climate change? *Global Environmental Change*, **21(1)**, 25-33.
- Berrang-Ford, L., J.D. Ford, and J. Paterson, 2011: Are we adapting to climate change? *Global Environmental Change*, **21(1)**, 25-33.
- Bhatt, U.S., D.A. Walker, M.K. Raynolds, J.C. Comiso, H.E. Epstein, G. Jia, R. Gens, J.E. Pinzon, C.J. Tucker, C.E. Tweedie, and P.J. Webber, 2010: Circumpolar Arctic tundra vegetation change is linked to sea ice decline. *Earth Interactions*, **14(8)**.
- Björk, R.G. and U. Molau, 2007: Ecology of alpine snowbeds and the impact of global change. *Arctic, Antarctic, and Alpine Research*, **39(1)**, 34-43.
- Björnsson H., T. Johannesson and A. Snorrason, 2011. Recent climate change, projected impacts and adaptation capacity in Iceland. Í: Linkov, I. & T. S. Bridges (Ed.) Global change and local adaptation. NATO Science for Peace and Security Series - C: Environmental Security. Springer, Dordrecht, s. 465-475
- Bluhm, B.A. and R. Gradinger, 2008: Regional Variability in Food Availability for Arctic Marine Mammals. *Ecological Applications*, **18(2, Supplement: Arctic Marine Mammals and Climate Change)**, pp. S77-S96.
- Bockheim, J., G. Vieira, M. Ramos, J. Lopez-Martinez, E. Serrano, M. Guglielmin, K. Wilhelm, A. Nieuwendam, 2013: Climate Warming and Permafrost Dynamics in the Antarctic Peninsula Region . *Global and Planetary Change*, **100: 215-223**
- Bogoyavlensky, D. and A. Siggner, 2004: *Arctic Demography*. *Arctic Human Development Report*. Stefansson Arctic Institute, Akureyri, Iceland, pp. 27.
- Bokhorst, S., J.W. Bjerke, L.E. Street, T.V. Callaghan, and G.K. Phoenix, 2011: Impacts of multiple extreme winter warming events on sub-Arctic heathland: Phenology, reproduction, growth, and CO2 flux responses. *Global Change Biology*, **17(9)**, 2817-2830.

- Bokhorst, S.F., J.W. Bjerke, H. Tømmervik, T.V. Callaghan, and G.K. Phoenix, 2009: Winter warming events damage sub-Arctic vegetation: Consistent evidence from an experimental manipulation and a natural event. *Journal of Ecology*, **97**(6), 1408-1415.
- Bolton, K., M. Lougheed, J. Ford, S. Nickels, C. Grable, and J. Shirley, 2011: *What we know, don't know, and need to know about climate change in Nunavut, Nunavik, and Nunatsiavut: A systematic literature review and gap analysis*. Inuit Tapiriit Kanatami, Ottawa, Canada, pp. 128.
- Bongo, M.P., A. Degteva, I.M.G. Eira, I. Hanssen-Bauer, A. Ivanoff, O.H. Magga, S.D. Mathiesen, N.G. Maynard, A. Oskal, M. Pogodaev, M.N. Sara, D.V. Schuler, E.I. Turi, and R.W. Corell, 2012: *Eurasian Reindeer Herding, Traditional Knowledge and Adaptation to Climate Change*. In: *UNESCO. Indigenous Peoples and Climate Change: Vulnerability, Adaptation and Traditional Knowledge. Report of workshop on Indigenous Peoples, Marginalized Populations and Climate Change: Vulnerability, Adaptation and Traditional Knowledge*. UNESCO, Paris.
- Bongo, M.P., A. Degteva, I.M.G. Eira, I. Hanssen-Bauer, A. Ivanoff, O.H. Magga, S.D. Mathiesen, N.G. Maynard, A. Oskal, M. Pogodaev, M.N. Sara, D.V. Schuler, E.I. Turi, and R.W. Corell, 2012: *Eurasian Reindeer Herding, Traditional Knowledge and Adaptation to Climate Change*. In: *UNESCO. Indigenous Peoples and Climate Change: Vulnerability, Adaptation and Traditional Knowledge. Report of workshop on Indigenous Peoples, Marginalized Populations and Climate Change: Vulnerability, Adaptation and Traditional Knowledge*. UNESCO, Paris.
- Bongo, M.P., A. Degteva, I.M.G. Eira, I. Hanssen-Bauer, A. Ivanoff, O.H. Magga, S.D. Mathiesen, N.G. Maynard, A. Oskal, M. Pogodaev, M.N. Sara, D.V. Schuler, E.I. Turi, and R.W. Corell, 2012: *Eurasian Reindeer Herding, Traditional Knowledge and Adaptation to Climate Change*. In: *UNESCO. Indigenous Peoples and Climate Change: Vulnerability, Adaptation and Traditional Knowledge. Report of workshop on Indigenous Peoples, Marginalized Populations and Climate Change: Vulnerability, Adaptation and Traditional Knowledge*. UNESCO, Paris.
- Borgström, R. and J. Museth, 2005: Accumulated snow and summer temperature ? critical factors for recruitment to high mountain populations of brown trout (*Salmo trutta* L.). *Ecology of Freshwater Fish*, **14**(4), 375-384.
- Bouchard, C. and L. Fortier, 2011: Circum-arctic comparison of the hatching season of polar cod *Boreogadus saida*: A test of the freshwater winter refuge hypothesis. *Progress in Oceanography*, **90**(1-4), 105-116.
- Bowden, W.B., M.N. Gooseff, A. Balsler, A. Green, B.J. Peterson, and J. Bradford, 2008: Sediment and nutrient delivery from thermokarst features in the foothills of the North Slope, Alaska: Potential impacts on headwater stream ecosystems. **113**, G02026.
- Boyd, P.W., K.R. Arrigo, R. Strzepek, and G.L. van Dijken, 2012: Mapping phytoplankton iron utilization: Insights into Southern Ocean supply mechanisms. *Journal of Geophysical Research: Oceans*, **117**(C6), - C06009.
- Brander, K.M., 2010: Cod *Gadus morhua* and climate change: processes, productivity and prediction. *Journal of Fish Biology*, **77**, 1899.
- Braun, C., O. Mustafa, A. Nordt, S. Pfeiffer, and H. Peter, 2012: Environmental monitoring and management proposals for the Fildes region, King George Island, Antarctic. *Polar Research*, **31**, <http://dx.doi.org/10.3402/polar.v31i0.18206>.
- Briffa, K.R., V.V. Shishov, T.M. Melvin, E.A. Vaganov, H. Grudd, R.M. Hantemirov, M. Eronen, and M.M. Naurzbaev, 2008: Trends in recent temperature and radial tree growth spanning 2000 years across northwest Eurasia. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **363**(1501), 2271-2284.
- Brommer, J.E., H. Pietiäinen, K. Ahola, P. Karell, T. Karstinenz, and H. Kolunen, 2010: The return of the vole cycle in southern Finland refutes the generality of the loss of cycles through 'climatic forcing'. *Global Change Biology*, **16**(2), 577-586.
- Bronen, R. 2009. Forced Migration of Alaskan Indigenous Communities Due to Climate Change: Creating a Human Rights Response. In *Linking Environmental Change, Migration & Social Vulnerability*. A. Oliver-Smith and X. Shen, eds. Pp. 68-73. UNU Institute for Environment and Human Security. Hohenkammer, Germany.
- Bronen, R. 2011. Climate-induced community relocations: creating an adaptive governance framework based in human rights doctrine. *NYU Review of Law and Social Change* 35:356-406
- Brown, Z. W., K. R. Arrigo, 2013: Sea ice impacts on spring bloom dynamics and net primary production in the Eastern Bering Sea. *Journal of Geophysical Research: Oceans*, **118**, 43-62.
- Brubaker, M., J. Berner, R. Chavan, and J. Warren, 2011a: Climate change and health effects in Northwest Alaska. *Global Health Action*, **4**, 10.3402/gha.v4i0.8445.

- Brubaker, M.Y., J.N. Bell, J.E. Berner, and J.A. Warren, 2011b: Climate change health assessment: a novel approach for Alaska Native communities. *International Journal of Circumpolar Health*, **70(3)**, 266-273.
- Brubaker, M.Y., J.N. Bell, J.E. Berner, and J.A. Warren, 2011b: Climate change health assessment: a novel approach for Alaska Native communities. *International Journal of Circumpolar Health*, **70(3)**, 266-273.
- Burek, K.A., F.M.D. Gulland, and T.M. O'Hara, 2008: effects of climate change on arctic marine mammal health. *Ecological Applications*, **18(2)**, S126-S134.
- Byrd, G.V., J.A. Schmutz, and H.M. Renner, 2008: Contrasting population trends of piscivorous seabirds in the Pribilof Islands: A 30-year perspective. *Deep-Sea Research Part II-Topical Studies in Oceanography*, **55(16-17)**, 1846-1855.
- CAFF 2013. Arctic Biodiversity Assessment. Status and trends in Arctic biodiversity. Conservation of Arctic Flora and Fauna, Akureyri. 557pp
- Callaghan, T.V., C.E. Tweedie, J. Åkerman, C. Andrews, J. Bergstedt, M.G. Butler, T.R. Christensen, D. Cooley, U. Dahlberg, R.K. Danby, F.J.A. Daniëls, J.G. De Molenaar, J. Dick, C.E. Mortensen, D. Ebert-May, U. Emanuelsson, H. Eriksson, H. Hedenås, G.H.R. Henry, D.S. Hik, J.E. Hobbie, E.J. Jantze, C. Jaspers, C. Johansson, M. Johansson, D.R. Johnson, J.F. Johnstone, C. Jonasson, C. Kennedy, A.J. Kenney, F. Keuper, S. Koh, C.J. Krebs, H. Lantuit, M.J. Lara, D. Lin, V.L. Lougheed, J. Madsen, N. Matveyeva, D.C. McEwen, I.H. Myers-Smith, Y.K. Narozhniy, H. Olsson, V.A. Pohjola, L.W. Price, F. Rigét, S. Rundqvist, A. Sandström, M. Tamstorf, R.V. Bogaert, S. Villarreal, P.J. Webber, and V.A. Zemtsov, 2011b: Multi-decadal changes in tundra environments and ecosystems: Synthesis of the international Polar year-back to the future project (IPY-BTF). *Ambio*, **40(6)**, 705-716.
- Callaghan, T.V., F. Bergholm, T.R. Christensen, C. Jonasson, U. Kokfelt, and M. Johansson, 2010: A new climate era in the sub-Arctic: Accelerating climate changes and multiple impacts. *Geophysical Research Letters*, **37(14)**, 6.
- Callaghan, T.V., L.O. Björn, Y. Chernov, F.S. Chapin, T.R. Christensen, B. Huntley, R. Ims, S. Jonasson, D. Jolly, N. Matveyeva, N. Panikov, W.C. Oechel, G.R. Shaver, J. Elster, H. Henttonen, I.S. Jónsdóttir, K. Laine, S. Schaphoff, S. Sitch, E. Taulavuori, K. Taulavuori, and C. Zöckler, 2005: Tundra and Polar Desert Ecosystems. In: *ACIA. Arctic Climate Impacts Assessment*. [Symon, C., L. Arris, and B. Heal(eds.)]. Cambridge University Press, Cambridge, UK, pp. 243-352.
- Callaghan, T.V., M. Johansson, R.D. Brown, P.Y. Groisman, N. Labba, and V. Radionov, 2011c: Changing snow cover and its impacts. In: *Snow, Water, Ice and Permafrost in the Arctic (SWIPA)*. AMAP, Oslo, Norway, pp. 4-1 to 4-58.
- Callaghan, T.V., T.R. Christensen, and E.J. Jantze, 2011a: Plant and vegetation dynamics on Disko island, west Greenland: Snapshots separated by over 40 years. *Ambio*, **40(6)**, 624-637.
- Callaghan, T.V., C. Jonasson, T. Thierfelder, Z. Yang, H. Hedenas, M. Johansson, U. Molau, R. Van Bogaert, A. Michelsen, J. Olofsson, D. Gwynn-Jones, S. Bokhorst, G. Phoenix, J. W. Bjerke, H. Tommervik, T.R. Christensen, E. Hanna, E. K. Koller, V.L. Sloan, 2013: Ecosystem change and stability over multiple decades in the Swedish subarctic: complex processes and multiple drivers. *Transactions of the Royal Society*, **368(1624)**, <http://dx.doi.org/10.1098/rstb.2012.0488>
- Canada NRTEE (National Round Table on the Environment and the Economy). 2009. True North: Adapting Infrastructure to Climate Change in Northern Canada. NRTEE, Ottawa, 146 p. (<http://www.nrtee-trnee.com/eng/publications/true-north/true-north-eng.pdf>).
- Carrie, J., F. Wang, H. Sanei, R.W. Macdonald, P.M. Outridge, and G.A. Stern, 2010: Increasing Contaminant Burdens in an Arctic Fish, Burbot (*Lota lota*), in a Warming Climate. *Environmental Science & Technology*, **44(1)**, 316-322.
- Castro de la Guardia, L., Derocher, A.E., Myers, P.G., Terwisscha van Scheltinga, A.D., and Lunn, N.J. 2013. Future sea ice conditions in western Hudson Bay and consequences for polar bears in the 21 century. *Global Change Biology*, **19**, 2675-2687.
- Chapman, E.W., E.E. Hofmann, D.L. Patterson, C.A. Ribic, and W.R. Fraser, 2011: Marine and terrestrial factors affecting Adélie penguin *Pygoscelis adeliae* chick growth and recruitment off the western Antarctic Peninsula. *Marine Ecology Progress Series*, **436**, 273-289.
- Cherosov, M.M., A.P. Isaev, V.I. Mironova, L.P. Lytkina, L.D. Gavriyeva, R.R. Sofronov, A.P. Arzhakova, N.V. Barashkova, I.A. Ivanov, I.F. Shurduk, A.P. Efimova, N.S. Karpov, P.A. Timofeyev, and L.V. Kuznetsova, 2010: Vegetation and human activity. In: *The Far North: plant biodiversity and ecology of Yakutia*. [Troeva, E.I., A.P. Isaev, M.M. Cherosov, and N.S. Karpov(eds.)]. Springer, Berlin, Germany, pp. 261-295.

- Cherry, S.G., A.E. Derocher, I. Stirling, and E.S. Richardson, 2009: Fasting physiology of polar bears in relation to environmental change and breeding behavior in the Beaufort Sea. *Polar Biology*, **32(3)**, 383-391.
- Cheung, W. W. L., J. Dunne, J. L. Sarmienot, and D. Pauly, 2011: Integrating ecophysiology and plankton dynamics into projected maximum fisheries catch potential under climate change in the Northeast Atlantic. *ICES Journal of Marine Fisheries*, **68(6)**:1008-1018.
- Cheung, W.W.L., V.W.Y. Lam, J.L. Sarmiento, K. Kearney, R. Watson, and D. Pauly, 2009: Projecting global marine biodiversity impacts under climate change scenarios. *Fish and Fisheries*, **10(3)**, 235-251.
- Chierici, M. and A. Fransson, 2009: Calcium carbonate saturation in the surface water of the Arctic Ocean: undersaturation in freshwater influenced shelves. *Biogeosciences*, **6(11)**, 2421-2431.
- Chown, S.L., J.E. Lee, K.A. Hughes, J. Barmes, P.J. Barrett, D.M. Bergstrom, P. Convey, D.A. Cowan, K. Crosbie, G. Dyer, Y. Frenot, S.M. Grant, D. Herr, M.C.I. Kennicutt, M. Lamers, A. Murray, H.P. Possingham, K. Reid, M.J. Riddle, P.G. Ryan, L. Sanson, J.D. Shaw, M.D. Sparrow, C. Summerhayes, A. Terauds, and D.H. Wall, 2012: Challenges to the future conservation of the Antarctic. *Science*, **337**, 158-159.
- Chown, S.L., J.E. Lee, K.A. Hughes, J. Barmes, P.J. Barrett, D.M. Bergstrom, P. Convey, D.A. Cowan, K. Crosbie, G. Dyer, Y. Frenot, S.M. Grant, D. Herr, M.C.I. Kennicutt, M. Lamers, A. Murray, H.P. Possingham, K. Reid, M.J. Riddle, P.G. Ryan, L. Sanson, J.D. Shaw, M.D. Sparrow, C. Summerhayes, A. Terauds, and D.H. Wall, 2012: Challenges to the future conservation of the Antarctic. *Science*, **337**, 158-159.
- Christoffersen, K.S., S.L. Amsinck, F. Landkildhus, T.L. Lauridsen, and E. Jeppesen, 2008: Lake Flora and Fauna in Relation to Ice-Melt, Water Temperature and Chemistry at Zackenberg. In: *Advances in Ecological Research*. [Hans Meltofte, Torben R. Christensen, Bo Elberling, Mads C. Forchhammer and Morten Rasch (ed.)]. Academic Press, pp. 371-389.
- Clark, D.A., D.S. Lee, M.M.R. Freeman, and S.G. Clark, 2008: Polar Bear Conservation in Canada: Defining the Policy Problems. *Arctic*, **61(4)**, 347-360.
- Clarke, L.J., S.A. Robinson, Q. Hua, D.J. Ayre, and D. Fink, 2012: Radiocarbon bomb spike reveals biological effects of Antarctic climate change. *Global Change Biology*, **18(1)**, 301-310.
- Cochran, P., O. H. Huntington, C. Pungowiyi, S. Tom, F S. Chapin III, H. P. Huntington, N. G. Maynard, S. F. Trainor. 2013. Indigenous Frameworks for Observing and Responding to Climate Change in Alaska. Special Issue Climatic Change. Vol. 12. Issue 3. 557-567. DOI:10.1007/s10584-013-0735-2
- Comeau, S., J.P. Gattuso, A.M. Nisumaa, and J. Orr, 2012: Impact of aragonite saturation state changes on migratory pteropods. *Proceedings of the Royal Society B-Biological Sciences*, 279(1729), 732-738.
- Comeau, S., R. Jeffree, J. Teyssié, and J. Gattuso, 2010: Response of the Arctic Pteropod *Limacina helicina* Projected Future Environmental Conditions. *PLoS ONE*, **5(6)**, e11362.
- Constable, A.J., 2011: Lessons from CCAMLR on the implementation of the ecosystem approach to managing fisheries. *Fish and Fisheries*, DOI: 10.1111/j.1467-2979.2011.00410.x.
- Convey, P. and M. Lebouvier, 2009: Environmental change and human impacts on terrestrial ecosystems of the sub-Antarctic islands between their discovery and the mid-twentieth century. *Society*, **143(1)**.
- Convey, P. and M.I. Stevens, 2007: Antarctic Biodiversity. *Science*, **317**, 1877-1878
- Convey, P., 2006: Roberto Bargagli, Ecological Studies 175: Antarctic Ecosystems – Environmental Contamination, Climate Change and Human Impact. *Journal of Paleolimnology*, **36(2)**, 223-224.
- Convey, P., 2010: Terrestrial biodiversity in Antarctica – Recent advances and future challenges. *Polar Science*, **4(2)**, 135-147.
- Convey, P., 2011: Antarctic terrestrial biodiversity in a changing world. *Polar Biology*, **34**, 1629-1641.
- Convey, P., R. Bindschadler, G. Di Prisco, E. Fahrbach, J. Gutt, D.A. Hodgson, P.A. Mayewski, C.P. Summerhayes, J. Turner, and A. Consortium, 2009: Antarctic climate change and the environment. *International Journal of Climatology*, **21(6)**, 541-563.
- Convey, P., R. Key, M. Belchier, and C. Waller, 2011: Recent range expansions in non-native predatory beetles on sub-Antarctic South Georgia. *Polar Biology*, **34(4)**, 597-602.
- Cook, A.J., S. Poncet, A.P.R. Cooper, D.J. Herbert, and D. Christie, 2010: Glacier retreat on South Georgia and implications for the spread of rats. *Antarctic Science*, **22(3)**, 255-263.
- Cooper, W. Lee, Ashjian, J. Carin, Smith, L. Sharon, Codispoti, A. Louis, Grebmeier, M. Jacqueline, Campbell, G. Robert, Sherr, and B. Evelyn, 2006: Rapid Seasonal Sea-Ice Retreat in the Arctic Could Be Affecting Pacific Walrus (*Odobenus rosmarus divergens*) Recruitment. *Aquatic Mammals*, **32(1)**, 98-102.
- Corbett, J.J., D.A. Lack, J.J. Winebrake, S. Harder, J.A. Silberman, and M. Gold, 2010: Arctic shipping emissions inventories and future scenarios. *Atmospheric Chemistry and Physics Discussions*, **10(4)**, 10271-10311.

- Costa, D.P., L. Huckstadt, D. Crocker, B. McDonald, M. Goebel, and M. Fedak, 2010: Approaches to studying climate change and its role on the habitat selection of Antarctic pinnipeds. *Integrative and Comparative Biology*, **3**, 919–92, doi:10.1093/icb/icq054.
- Costa, D.P., L. Huckstadt, D. Crocker, B. McDonald, M. Goebel, and M. Fedak, 2010: Approaches to studying climate change and its role on the habitat selection of Antarctic pinnipeds. *Integrative and Comparative Biology*, **50(6)**, doi:10.1093/icb/icq054.
- Cowan, D.A., S.L. Chown, P. Convey, M. Tuffin, K. Hughes, S. Pointing, and W.F. Vincent, 2011: Non-indigenous microorganisms in the Antarctic: assessing the risks Elsevier Trends Journals, pp. 540-548.
- Coyle, K.J. and L.V. Susteren, 2012: *The Psychological Effects of Global Warming on the United States: Its Stresses, Traumas, and Societal Costs*. National Wildlife Federation Climate Education Program, pp. 41.
- Coyle, K.O., L.B. Eisner, F.J. Mueter, A.I. Pinchuk, M.A. Janout, K.D. Ciciel, E.V. Farley, and A.G. Andrews, 2011: Climate change in the southeastern Bering Sea: impacts on pollock stocks and implications for the oscillating control hypothesis. *Fisheries Oceanography*, **20(2)**, 139-156.
- Crate, S. A. and M. Nuttall 2009: *Anthropology and Climate Change: from encounters to actions*, Walnut Creek, CA: Left Coast Press
- Crate, S.A., B.C. Forbes, L. King, and J. Kruse, 2010: Contact with nature. In: *Arctic Social Indicators: a follow-up to the Arctic Human Development Report*. [Larsen, J.N., P. Schweitzer, and G. Fondahl(eds.)]. TemaNord, pp. 109.
- Croxall, J.P., S.H.M. Butchart, B. Lascelles, A.J. Stattersfield, B. Sullivan, A. Symes, and P. Taylor, 2012: Seabird conservation status, threats and priority actions: a global assessment. *Bird Conservation International*, **22(1)**, 1.
- Cruikshank, J., 2001: Glaciers and Climate Change: Perspectives from Oral Tradition. *Arctic*, **54(4)**.
- Cubillos, J.C., S.W. Wright, G. Nash, M.F. de Salas, B. Griffiths, B. Tilbrook, A. Poisson, and G.M. Hallegraef, 2007: Calcification morphotypes of the coccolithophorid *Emiliania huxleyi* in the Southern Ocean: changes in 2001 to 2006 compared to historical data. *Marine Ecology-Progress Series*, **348**, 47-54
- Cullen-Unsworth, L.C., R. Hill, J.R.A. Butler, and M. Wallace, 2011: A research process for integrating Indigenous and scientific knowledge in cultural landscapes: principles and determinants of success in the Wet Tropics World Heritage Area, Australia. *The Geographical Journal*, **178(4)**, 351-365.
- Curtis, T., S. Kvernmo, and P. Bjerregaard, 2005: Changing living conditions, life style and health. *International Journal of Circumpolar Health*, **64(5)**, 442.
- Dalpadado, P., R.B. Ingvaldsen, L.C. Stige, B. Bogstad, T. Knutsen, G. Ottersen and B. Ellertsen, 2012: Climate effects on Barents Sea ecosystem dynamics. *ICES Journal of Marine Science*, **69(7)**, 1303–1316.
- Danby, R.K. and D.S. Hik, 2007: Variability, contingency and rapid change in recent subarctic alpine tree line dynamics. *Journal of Ecology*, **95(2)**, 352-363.
- Danby, R.K., S. Koh, D.S. Hik, and L.W. Price, 2011: Four decades of plant community change in the alpine tundra of southwest Yukon, Canada. *Ambio*, **40(6)**, 660-671.
- Daniëls, F.J.A. and J.G. De Molenaar, 2011: Flora and vegetation of Tasiilaq, formerly Angmagssalik, southeast Greenland: A comparison of data between around 1900 and 2007. *Ambio*, **40(6)**, 650-659.
- Dankers, R. and H. Middelkoop, 2008: River discharge and freshwater runoff to the Barents Sea under present and future climate conditions. *Climatic Change*, **87(131-153)**.
- Dannevig, H., T. Rauken, and G. Hovelsrud, 2012: Implementing adaptation to climate change at the local level. *Local Environment: The International Journal of Justice and Sustainability*, **17(6-7)**, 1-15.
- Darnis, G., D. Robert, C. Pomerleau, H. Link, P. Archambault, R.J. Nelson, M. Geoffroy, J. Tremblay, C. Lovejoy, S. H. Ferguson, B. P. V. Hunt, L. Fortier, 2012: Current state and trends in Canadian Arctic marine ecosystems: II. Heterotrophic food web, pelagic-benthic coupling, and biodiversity. *Climatic Change*, **115(1)**, 179-205.
- Dawson, J., E.J. Stewart, H. Lemelin, and D. Scott, 2010: The carbon cost of polar bear viewing tourism in Churchill, Canada. *Journal of Sustainable Tourism*, **18(3)**, 319-336.
- de Neergaard E., P. Stougaard, K. Høegh, L. Munk, 2009: Climatic changes and agriculture in Greenland: Plant diseases in potatoes and grass fields. IOP Conf. Ser.: Earth Environ. Sci. **6(37)**, doi:10.1088/1755-1307/6/37/372013
- Degteva, A. and C. Nellemann. 2013. Nenets migration in the landscape: impacts of industrial development in Yamal Peninsula, Russia. *Pastoralism: Research, Policy and Practice* 2013, 3:15. SpringerOpen Journal, <http://www.pastoralismjournal.com/content/3/1/15>
- Derksen, C., S.L. Smith, M. Sharp, L. Brown, S. Howell, L. Copland, D.R. Mueller, Y. Gauthier, C.G. Fletcher, A. Tivy, M. Bernier, J. Bourgeois, R. Brown, C.R. Burn, C. Duguay, P. Kushner, A. Langlois, A.G. Lewkowicz,

- A. Royer, and A. Walker, 2012: Variability and change in the Canadian cryosphere. *Climatic Change*, **115(1)**, 59-88.
- Derocher, A.E., M. Andersen, Wiig Ø, J. Aars, E. Hansen, and M. Biuw, 2011: Sea ice and polar bear den ecology at Hopen Island, Svalbard. *Mar Ecol Prog Ser*, **441**, 273-279.
- Derocher, A.E., N.J. Lunn, and I. Stirling, 2004: Polar bears in a warming climate. *Integrative and Comparative Biology*, **44(2)**, 163-176.
- Derome, J. and N. Lukina, 2011: Interaction between environmental pollution and land-cover/land-use change in Arctic areas. In: Eurasian Arctic Land Cover and Land Use in a Changing Climate. [Gutman, G. (ed)] Springer, pp. 269-290
- Déry, S.J. and E.F. Wood, 2005: Decreasing river discharge in northern Canada. *Geophysical Research Letters*, **32(10)**, - L10401.
- Dibike, Y., T. Prowse, T. Saloranta, and R. Ahmed, 2011: Response of Northern Hemisphere lake-ice cover and lake-water thermal structure patterns to a changing climate. *Hydrological Processes*, **25(19)**, 2942-2953.
- Dinniman, M.S., J.M. Klinck, and E.E. Hofmann, 2012: Sensitivity of Circumpolar Deep Water Transport and Ice Shelf Basal Melt along the West Antarctic Peninsula to Changes in the Winds. *Journal of Climate*, **25(14)**, 4799-4816.
- Donaldson S.G. et al, 2010. Environmental contaminants and human health in the Canadian Arctic. *Science of the Total Environment* **408**, (2010).
- Donelson, J.M., P.L. Munday, M.I. McCormick, and G.E. Nilsson, 2011: Acclimation to predicted ocean warming through developmental plasticity in a tropical reef fish. *Global Change Biology*, **17(4)**, 1712-1719.
- Drinkwater, K.F., 2011: The influence of climate variability and change on the ecosystems of the Barents Sea and adjacent waters: Review and synthesis of recent studies from the NESSAS project. *Progress in Oceanography*, **90(2011)**, 47-61.
- Drinkwater, K.F., G. Beaugrand, M. Kaeriyama, S. Kim, G. Ottersen, R.I. Perry, H. Pörtner, J.J. Polovina, and A. Takasuka, 2010: On the processes linking climate to ecosystem changes. *Journal of Marine Systems*, **79(3-4)**, 374-388.
- Duc, N., P. Crill, and D. Bastviken, 2010: Implications of temperature and sediment characteristics on methane formation and oxidation in lake sediments. *Biogeochemistry*, **100(1)**, 185-196.
- Ducklow, H., A. Clarke, R. Dickhut, S. C. Doney, H. Geisz, K. Huang, D. G. Martinson, M. P. Meredith, H. V. Moeller, M. Montes-Hugo, O. Schofield, S. E. Stammerjohn, D. Steinberg, and W. Fraser. 2012: The marine system of the western Antarctic Peninsula. Pp121-159 in [A. Rogers, N. Johnston, E. Murphy, and A. Clarke, (editors)]. Antarctic Ecosystems. Blackwell Publishing, Ltd.
- Duhaime, G., A. Lemelin, V. Didyk, O. Goldsmith, G. Winther, A. Caron, N. Bernard, and A. Goodmaire, 2004: Economic systems. In: *The Arctic Human Development Report*. Stefansson Arctic Institute, Akureyri, Iceland, pp. 69.
- Dumas, J.A., G.M. Flato, and R.D. Brown, 2006: Future Projections of Landfast Ice Thickness and Duration in the Canadian Arctic. *Journal of Climate*, **19(20)**, 5175-5189.
- Durner, G., J. Whiteman, H. Harlow, S. Amstrup, E. Regehr, and M. Ben-David, 2011: Consequences of long-distance swimming and travel over deep-water pack ice for a female polar bear during a year of extreme sea ice retreat. *Polar Biology*, **34(7)**, 975-984.
- Durner, G.M., D.C. Douglas, R.M. Nielson, S.C. Amstrup, T.L. McDonald, I. Stirling, M. Mauritzen, E.W. Born, O. Wiig, E. DeWeaver, M.C. Serreze, S.E. Belikov, M.M. Holland, J. Maslanik, J. Aars, D.A. Bailey, and A.E. Derocher, 2009: Predicting 21st-century polar bear habitat distribution from global climate models. *Ecological Monographs*, **79(1)**, 25-58.
- Dyck, M. and S. Romberg, 2007: Observations of a wild polar bear (*Ursus maritimus*) successfully fishing Arctic charr (*Salvelinus alpinus*) and Fourhorn sculpin (*Myoxocephalus quadricornis*). *Polar Biology*, **30(12)**, 1625-1628.
- Dyck, M.G. and E. Kebreab, 2009: Estimating the Energetic Contribution of Polar Bear (*Ursus Maritimus*) Summer Diets to the Total Energy Budget. *Journal of Mammalogy*, **90(3)**, 585-593.
- Dyck, M.G., W. Soon, R.K. Baydack, D.R. Legates, S. Baliunas, T.F. Ball, and L.O. Hancock, 2007: Polar bears of western Hudson Bay and climate change: Are warming spring air temperatures the “ultimate” survival control factor? *Ecological Complexity*, **4(3)**, 73-84.

- Dyck, M.G., W. Soon, R.K. Baydack, D.R. Legates, S. Baliunas, T.F. Ball, and L.O. Hancock, 2008: Reply to response to Dyck et al. (2007) on polar bears and climate change in western Hudson Bay by Stirling et al. (2008). *Ecological Complexity*, **5(4)**, 289-302.
- Eicken, H., Lovcraft, A., Druckenmiller, M., 2009: Sea-Ice System Services: A Framework to Help Identify and Meet Information Needs Relevant for Observing Networks. *Arctic*, **62(2)**.
- Eicken, H., Lovcraft, A., Druckenmiller, M., 2009: Sea-Ice System Services: A Framework to Help Identify and Meet Information Needs Relevant for Observing Networks. *Arctic*, **62(2)**.
- Eira, I.M.G., C. Jaedicke, O.H. Magga, N. Maynard, D. Vikhamar-Schuler, and S.D. Mathiesen, 2012: *Traditional Sami Snow Terminology and Physical Snow Classification – Two Ways of Knowing*. *Cold Regions Science and Technology*, **85 (117-130)**, doi: 10.1016/j.coldregions.2012.09.004
- Ellingsen, I. H., P. Dalpadado, D. Slagstad, and H. Loeng. 2008. Impact of climatic change on the biological production in the Barents Sea. *Climatic Change*, **87**, 155-175.
- Elmendorf, S.C., G.H. Henry, R.D. Hollister, R.G. Bjork, A.D. Bjorkman, T.V. Callaghan, L.S. Collier, E.J. Cooper, J.H. Cornelissen, T.A. Day, A.M. Fosaa, W.A. Gould, J. Gretarsdottir, J. Harte, L. Hermanutz, D.S. Hik, A. Hofgaard, F. Jarrad, I.S. Jonsdottir, F. Keuper, K. Klanderud, J.A. Klein, S. Koh, G. Kudo, S.I. Lang, V. Loewen, J.L. May, J. Mercado, A. Michelsen, U. Molau, I.H. Myers-Smith, S.F. Oberbauer, S. Pieper, E. Post, C. Rixen, C.H. Robinson, N.M. Schmidt, G.R. Shaver, A. Stenstrom, A. Tolvanen, O. Totland, T. Troxler, C.H. Wahren, P.J. Webber, J.M. Welker, and P.A. Wookey, 2012a: Global assessment of experimental climate
- Elmendorf, S.C., G.H.R. Henry, and R.D. Hollister, R.G. Björk, N. Boulanger-Lapointe, E.J. Cooper, J.H.C. Cornelissen, T.A. Day, E. Dorrepaal, T.G. Elumeeva, M. Gill, W.A. Gould, J. Harte, D.S. Hik, A. Hofgaard, D.R. Johnson, J.F. Johnstone, I.S. Jónsdóttir, J.C. Jorgenson, K. Klanderud, J.A. Klein, S. Koh, G. Kudo, M. Lara, E. Lévesque, B. Magnússon, J.L. May, J.A. Mercado-Dí'az, A. Michelsen, U. Molau, I.H. Myers-Smith, S.F. Oberbauer, V.G. Onipchenko, C. Rixen, N. Martin Schmidt, G.R. Shaver, M.J. Spasojevic, P. Ellen Þórhallsdóttir, A. Tolvanen, T. Troxler, C.E. Tweedie, S. Villareal, C-H. Wahren, X. Walker, P. J. Webber, J.M. Welker, and S. Wipf, 2012b: Plot-scale evidence of tundra vegetation change and links to recent summer warming. *Nature Climate Change*, **2**, 453-457
- Emmert, C.A., L.F.W. Lesack, and W.F. Vincent, 2008: Mackenzie River nutrient delivery to the Arctic Ocean and effects of the Mackenzie Delta during open water conditions. *Global Biogeochemical Cycles*, **22(1)**, GB1024.
- Epstein, H.E., M.K. Reynolds, D.A. Walker, U.S. Bhatt, C.J. Tucker, and J.E. Pinzon, 2012: Dynamics of aboveground phytomass of the circumpolar Arctic tundra during the past three decades. *Environmental Research Letters*, **7(1)**, 015506
- Epstein, P.R. and D. Ferber, 2011: *Changing Planet, Changing Health: How the Climate Crisis Threatens Our Health and What We Can Do about It*. University of California Press Books, pp. 368.
- Eskeland, G. S. and L. S. Flottorp, 2006. Climate change in the Arctic: A discussion of the impact on economic activity. In: Glomsrød, Solveig and Iulie Aslaksen (eds.), *The Economy of the North*. Statistics Norway. oslo, Norway.
- Eskeland, G. S. and L. S. Flottorp, 2006: Climate change in the Arctic: A discussion of the impact on economic activity. In: Glomsrød, Solveig and Iulie Aslaksen (eds.), *The Economy of the North*. Statistics Norway. Oslo, Norway.
- Lindholdt, L., & Glomsrød, S. (2012). The Arctic: No big bonanza for the global petroleum industry. *Energy Economics*, **34**, 1465-1474.
- Fabry, V.J., J.B. McClintock, J.T. Mathis, and J.M. Grebmeier, 2009: Ocean Acidification at High Latitudes: The Bellwether. *Oceanography*, **22(4)**, 160-171.
- Falk-Petersen, S., P. Mayzaud, G. Kattner, and J. R. Sargent. 2009. Lipids and life strategy of Arctic Calanus. *Marine Biology Research*, **5**:18-39.
- Favero-Longo, S.E., N. Cannone, M.R. Worland, P. Convey, R. Piervittori, and M. Guglielmin, 2011: Changes in lichen diversity and community structure with fur seal population increase on Signy Island, South Orkney Islands. *Antarctic Science*, **23**, 65.
- Fechhelm, R.G., B. Streever, and B.J. Gallaway, 2007: The arctic cisco (*Coregonus autumnalis*) subsistence and commercial fisheries, Colville River, Alaska: a conceptual model. *Arctic*, **60**, 421-429.
- Ferguson, S., M. Kingsley, and J. Higdon, 2012: Killer whale (*Orcinus orca*) predation in a multi-prey system. *Population Ecology*, **54(1)**, 31-41.

- Ferguson, S.H., I. Stirling, and P. McLoughlin, 2005: Climate change and ringed seal (*Phoca hispida*) recruitment in western Hudson Bay. *Marine Mammal Science*, **21(1)**, 121-135.
- Fischbach, A.S., S.C. Amstrup, and D.C. Douglas, 2007: Landward and eastward shift of Alaskan polar bear denning associated with recent sea ice changes. *Polar Biology*, **30**, 1395-1405.
- Flanagan, K.M., E. Mccauley, and F. Wrona, 2006: Freshwater food webs control carbon dioxide saturation through sedimentation. *Global Change Biology*, **12(4)**, 644-651.
- Flores, H., A. Atkinson, S. Kawaguchi, B.A. Krafft, G. Milinevsky, S. Nicol, C. Reiss, G.A. Tarling, R. Werner, E. Bravo Rebolledo, V. Cirelli, J. Cuzin-Roudy, S. Fielding, J.A. van Franeker, J.J. Groeneveld, M. Haraldsson, A. Lombana, E. Marschoff, B. Meyer, E.A. Pakhomov, A.P. Van de Putte, E. Rombol, K. Schmidt, V. Siegel, M. Teschke, H. Tonkes, J.Y. Toullec, P.N. Trathan, N. Tremblay, and T. Werner, 2012: Impact of climate change on Antarctic krill. *Marine Ecology Progress Series*, 458, 1-19.
- Forbes, B.C. and F. Stammer, 2009: Arctic climate change discourse: the contrasting politics of research agendas in the West and Russia. *Polar Research*, **28(1)**, 28-42.
- Forbes, B.C. and T. Kumpula, 2009: The ecological role and geography of reindeer (*Rangifer tarandus*) in Northern Eurasia. *Geography Compass*, **3/4**, 1356-1380.
- Forbes, B.C., 2006: *Reindeer Management in Northernmost Europe: Linking Practical And Scientific Knowledge in Social-Ecological Systems*. Springer.
- Forbes, B.C., 2008: Equity, vulnerability and resilience in social-ecological systems: a contemporary example from the Russian Arctic. *Research in Social Problems and Public Policy*, **15**, 203.
- Forbes, B.C., F. Stammer, T. Kumpula, N. Meschytyb, A. Pajunen, and E. Kaarlejärvi, 2009: High resilience in the Yamal-Nenets social-ecological system, West Siberian Arctic, Russia. *Proceedings of the National Academy of Sciences of the United States of America*, **106(52)**, 22041-22048.
- Forbes, B.C., M. Bølter, L. Müller-Wille, J. Hukkinen, F. Müller, N. Gunslay, Y. Konstantinov (eds.) 2006: Reindeer management in northernmost Europe: linking practical and scientific knowledge in social-ecological systems. *Ecological Studies*, **184**. Springer-Verlag, Berlin. pp. 11-25.
- Forbes, B.C., M.M. Fauria, and P. Zetterberg, 2010: Russian Arctic warming and 'greening' are closely tracked by tundra shrub willows. *Global Change Biology*, **16(5)**, 1542-1554.
- Forbes, D.L. (ed.), 2011: *State of the Arctic Coast 2010 - Scientific Review and Outlook*. International Permafrost Association. HelmholtzZentrum, Geesthacht, Germany, pp. 178.
- Forcada, J., P.N. Trathan, P.L. Boveng, I.L. Boyd, J.M. Burns, D.P. Costa, M. Fedak, T.L. Rogers, and C.J. Southwell, 2012: Responses of Antarctic pack-ice seals to environmental change and increasing krill fishing. *Biological Conservation*, **149(1)**, 40-50.
- Forchhammer, M.C., N.M. Schmidt, T.T. Høye, T.B. Berg, D.K. Hendrichsen, and E. Post, 2008: Population Dynamical Responses to Climate Change. *Advances in Ecological Research*, **40**, 391-419.
- Ford, J., and Goldhar, C., Climate change vulnerability and adaptation in resource dependent communities: a case study from West Greenland. *Climate Research* **54**, 181-196. (2012).
- Ford, J., B. Pearce, B. Smit, J. Wandel, M. Allurut, K. Shappa, H. Ittusujurat, and K. Qrunnut, 2007: Reducing vulnerability to climate change in the Arctic: The case of Nunavut, Canada. *Arctic*, **60 (2)**, 150-166.
- Ford, J., B. Smit, and J. Wandel, 2006: Vulnerability to climate change in the Arctic: A case study from Arctic Bay, Nunavut. *Global Environmental Change*, **16 (2)**, 145-160.
- Ford, J., T. Pearce, J. Prno, F. Duerden, L. Berrang Ford, M. Beaumier, and T. Smith, 2010: Perceptions of climate change risks in primary resource use industries: a survey of the Canadian mining sector. *Regional Environmental Change*, **10(1)**, 65-81.
- Ford, J.D. and C. Furgal, 2009: Foreword to the special issue: climate change impacts, adaptation and vulnerability in the Arctic. *Polar Research*, **28(1)**, 1-9.
- Ford, J.D. and L. Berrang-Ford, 2009: Food security in Igloodik, Nunavut: an exploratory study. *Polar Record*, **45(03)**, 225.
- Ford, J.D. and T. Pearce, 2010: What we know, do not know, and need to know about climate change vulnerability in the western Canadian Arctic: a systematic literature review. *Environmental Research Letters*, **5(1)**, 014008.
- Ford, J.D., 2009: Dangerous climate change and the importance of adaptation for the Arctic's Inuit population. *Environmental Research Letters*, **4(2)**.
- Frederiksen, M., Anker-Nilssen, T., Beaugrand, G., Wanless, S. 2013: Climate, copepods and seabirds in the boreal Northeast Atlantic - current state and future outlook. *Global Change Biology*, **19**, 364-372

- Freitas, C., K.M. Kovacs, R.A. Ims, and C. Lydersen, 2008: Predicting habitat use by ringed seals (*Phoca hispida*) in a warming Arctic. *Ecological Modelling*, **217(1-2)**, 19-32.
- Frenot, Y., S.L. Chown, J. Whinam, P.M. Selkirk, P. Convey, M. Skotnicki, and D.M. Bergstrom, 2005: Biological invasions in the Antarctic: extent, impacts and implications. *Biological Reviews*, **80(01)**, 45.
- Frey, K.E. and J.W. McClelland, 2009: Impacts of permafrost degradation on arctic river biogeochemistry. *Hydrological Processes*, **23(1)**, 169-182.
- Froese, R. and A. Proelß, 2010: Rebuilding fish stocks no later than 2015: will Europe meet the deadline? *Fish and Fisheries*, **11(2)**, 194-202.
- Fuglei, E. and R.A. Ims, 2008: Global warming and effects on the arctic fox. *Science Progress*, **91(2)**, 175-191.
- Furgal C., T.D. Prowse, 2008: Northern Canada. In: *From Impacts to Adaptation: Canada in a Changing Climate*. [Lemmen,D.S., Warren, F.J., Lacroix, J. (ed.)].
- Furgal, C. and J. Seguin, 2006: Climate Change, Health, and Vulnerability in Canadian Northern Aboriginal Communities. *Environ Health Perspect*, **114(12)**.
- Galand, P.E., C. Lovejoy, J. Pouliot, M.E. Garneau, and W.F. Vincent, 2008: Microbial community diversity and heterotrophic production in a coastal Arctic ecosystem: A stamukhi lake and its source waters. *Limnology and Oceanography*, **53(2)**, 813.
- Galloway-McLean, K., 2010: *Advance Guard: Climate Change Impacts, Adaptation, Mitigation and Indigenous Peoples – A Compendium of Case Studies*. Nations University – Traditional Knowledge Initiative, Darwin, Australia, pp. 124.
- Gamon et al. 2013. Spatial and temporal variation in primary productivity (NDVI) of coastal Alaskan tundra: Decreased vegetation growth following earlier snowmelt. *Remote Sensing of Environment*, **129**, 144-153.
- GAO, 2009: *U. S. General Accounting Office Report: Alaska Native Villages. Limited Progress has been Made on Relocating Villages Threatened by Flooding and Erosion*. In: Report to Congressional Requesters, pp. 29-40.
- Gaston, A. J. 2011: Arctic seabirds: Diversity, populations, trends, and causes. in R. T. Watson, T. J. Cade, M. Fuller, G. Hunt, and E. Potapov (Eds.). *Gyrfalcons and Ptarmigan in a Changing World, Volume I. The Peregrine Fund*, Boise, Idaho, USA. Pp. 147-160, <http://dx.doi.org/10.4080/gpcw.2011.0201>
- Gaston, A.J. and K. Woo, 2008: Razorbills (*Alca Torda*) Follow Subarctic Prey into the Canadian Arctic: Colonization Results from Climate Change? *Auk*, **125(4)**, 939-942.
- Gaston, A.J., H.G. Gilchrist, and J.M. Hipfner, 2005: Climate change, ice conditions and reproduction in an Arctic nesting marine bird: Brunnich's guillemot (*Uria lomvia* L.). *Journal of Animal Ecology*, **74(5)**, 832-841.
- Gaston, A.T., H.G. Gilchrist, M.L. Mallory, and P.A. Smith, 2009: Changes in Seasonal Events, Peak Food Availability, and Consequent Breeding Adjustment in a Marine Bird: a Case of Progressive Mismatching. *Condor*, **111(1)**, 111-119.
- Gautier, D.L., K.J. Bird, R.R. Charpentier, A. Grantz, D.W. Houseknecht, T.R. Klett, T.E. Moore, J.K. Pitman, C.J. Schenk, J.H. Schuenemeyer, K. Sørensen, M.E. Tennyson, Z.C. Valin, and C.J. Wandrey, 2009: Assessment of Undiscovered Oil and Gas in the Arctic. *Science*, **324(5931)**, 1175-1179.
- Gearheard, S., C. Aporta, G. Aipellee, and K. O'Keefe, 2011: The Igliniit project: Inuit hunters document life on the trail to map and monitor arctic change. *Canadian Geographer / Le Géographe Canadien*, **55(1)**, 42-55.
- Gearheard, S., G. Aipellee, and K. O'Keefe, 2010: The Igliniit project: Combining Inuit knowledge and geomatics engineering to develop a new observation tool for hunters. In: *SIKU: Knowing our ice, documenting Inuit sea-ice knowledge and use*. [Krupnik, I., C. Aporta, S. Gearheard, G.J. Laidler, and L. Kielsen-Holm(eds.)]. Springer, pp. 181.
- Gearheard, S., W. Matumeak, I. Angutikjuaq, J. Maslanik, H.P. Huntington, J. Leavitt, D.M. Kagak, G. Tigullaraq, and R.G. Barry, 2006: "It's not that simple": A collaborative comparison of sea ice environments, their uses, observed changes, and adaptations in Barrow, Alaska, USA, and Clyde River, Nunavut, Canada. *Ambio*, **35(4)**, 203-211.
- Geffen, A.J., H. Høie, A. Folkvord, A.K. Hufthammer, C. Andersson, U. Ninnemann, R.B. Pedersen, and K. Nedreaas, 2011: High-latitude climate variability and its effect on fisheries resources as revealed by fossil cod otoliths. *ICES Journal of Marine Science*: **68(6)**, 1081-1089.
- Gilg, O., B. Sittler, and I. Hanski, 2009: Climate change and cyclic predator-prey population dynamics in the high Arctic. *Global Change Biology*, **15(11)**, 2634-2652.
- Gilg, O., Kovacs, K.M., Aars,J., Fort,J., Gauthier,G., Grémillet,D., Ims,R.A., Meltote,H., Moreau,J., Post,E., Schmidt,N.M., Yannic,G., and Bollache, L. 2012. Climate change and the ecology and evolution of Arctic vertebrates. *Annals of the New York Academy of Sciences*, **1249**, 160-190

- Gjosaeter, H., B. Bogstad, S. Tjelmeland, 2009: Ecosystem effects of the three capelin stock collapses in the Barent Sea. *Marine Biology Research*, **5**, 40-53.
- Gleason, J.S. and K.D. Rode, 2009: Polar Bear Distribution and Habitat Association Reflect Long-term Changes in Fall Sea Ice Conditions in the Alaskan Beaufort Sea. *Arctic*, **62(4)**, 405-417.
- Glomsrød, S., and Aslaksen I. 2009. *The Economy of the North 2008*. Statistics Norway. Oslo, Norway
- Goetz, S.J., A.G. Bunn, G.J. Fiske, and R.A. Houghton, 2005: Satellite-observed photosynthetic trends across boreal North America associated with climate and fire disturbance. *Proceedings of the National Academy of Sciences of the United States of America*, **102(38)**, 13521-13525.
- Grant, S., A. Constable, B. Raymond, and S. Doust, 2006: *Bioregionalisation of the Southern Ocean: Report of Experts Workshop, WWF- Australia and ACE CRC.*, Hobart, September 2006.
- Grebmeier, J.M., 2012: Shifting Patterns of Life in the Pacific Arctic and Sub-Arctic Seas. *Annual Review of Marine Science*, **4(1)**, 63-78.
- Grebmeier, J.M., J.E. Overland, S.E. Moore, E.V. Farley, E.C. Carmack, L.W. Cooper, K.E. Frey, J.H. Helle, F.A. McLaughlin, and S.L. McNutt, 2006: A Major Ecosystem Shift in the Northern Bering Sea. *Science*, **311(5766)**, 1461-1464.
- Green, D. and G. Raygorodetsky, 2010: Indigenous knowledge of a changing climate. *Climatic Change*, **100(2)**, 239-242.
- Greenwald, M.J., W.B. Bowden, M.N. Gooseff, J.P. Zarnetske, J.P. McNamara, J.H. Bradford, and T.R. Brosten, 2008: Hyporheic exchange and water chemistry of two arctic tundra streams of contrasting geomorphology. **113**, G02029.
- Gremillet D. and Boulinier T., 2009: Spatial ecology and conservation of seabirds facing global climate change: A review. *Mar.Ecol.Prog.Ser.Marine Ecology Progress Series*, **391**, 121-137.
- Grémillet, D., Welcker, J., Karnovsky, N.J., Walkusz, W., Hall, M.E., Fort8, J., Brown, Z.W., Speakman, J.R., Harding, A.M. A. 2012. Little auks buffer impacts of current Arctic climate change. *Marine Ecology Progress Series*, **45**, 197-206
- Grenfell, T.C., and J. Putkonen (2008) A method for the detection of the severe rain-on-snow event on Banks Island, October 2003, using passive microwave remote sensing. *Water Resources Research* 44: W03425.
- Griffith, B., L.G. Adams, D.C. Douglas, C. Cuyler, R.G. White, A. Gunn, D.E. Russell, and R.D. Cameron, 2010: Science: E-Letters. In: *No Evidence of Trophic Mismatch for Caribou in Greenland* .
- Grønlund, A., 2009: Virkning av klimaendring på arealbruk i norsk arktis. *Bioforsk Rapport*, **4(109)**.
- Gudasz, C., D. Bastviken, K. Steger, K. Premke, S. Sobek, and L.J. Tranvik, 2010: Temperature-controlled organic carbon mineralization in lake sediments. *Nature*, **466(7305)**, 478-481.
- Gunn, A., Miller, F., Barry, S., Buchan, A., 2009: A Near-Total Decline in Caribou on Prince of Wales, Somerset, and Russell Islands, Canadian Arctic. *Arctic*, **59(1)**.
- Gutt, J., I. Barratt, E. Domack, C.D. d'Acoz, W. Dimmler, A. Gremare, O. Heilmayer, E. Isla, D. Janussen, E. Jorgensen, K.H. Kock, L.S. Lehnert, P. Lopez-Gonzales, S. Langner, K. Linse, M.E. Manjon-Cabeza, M. Meissner, A. Montiel, M. Raes, H. Robert, A. Rose, E.S. Schepisi, T. Saucedo, M. Scheidat, H.W. Schenke, J. Seiler, and C. Smith, 2011: Biodiversity change after climate-induced ice-shelf collapse in the Antarctic. *Deep-Sea Research Part II-Topical Studies in Oceanography*, **58(1-2)**, 74-83.
- Halldórsson G., B.D. Sigurdsson, O.E.S. Hrafnkelsdóttir, Ó. Eggertsson, E. Ólafsson, 2013: New anthropod herbivores on trees and shrubs in Iceland and changes in pest dynamics: A review. *ICEL. AGRIC. SCI.* **26 (2013)**, 69-84
- Hallinger, M., M. Manthey, and M. Wilmking, 2010: Establishing a missing link: warm summers and winter snow cover promote shrub expansion into alpine tundra in Scandinavia. *New Phytologist*, **186(4)**, 890-899.
- Hansen et al. 2013. Climate events synchronize the dynamics of a resident vertebrate community in the high arctic. *Science* **339**, 313-315.
- Hansen, B.B., R. Aanes, I. Herfindal, J. Kohler, B.E. Sæther, and M.K. Oli, 2011: Climate, icing, and wild arctic reindeer: Past relationships and future prospects. *Ecology*, **92(10)**, 1917-1923.
- Harsch, M.A., P.E. Hulme, M.S. McGlone, and R.P. Duncan, 2009: Are treelines advancing? A global meta-analysis of treeline response to climate warming. *Ecology Letters*, **12(10)**, 1040-1049.
- Harsem, Ø., Eide, A. & Heen, K. (2011). Factors Influencing Future Oil and Gas Prospects in the Arctic. *Energy Policy*. 39: 8037-8045.

- Hátún, H., M. R. Payne, and J. A. Jacobsen. 2009: The North Atlantic subpolar gyre regulates the spawning distribution of blue whiting (*Micromesistius poutassou*). *Canadian Journal of Fisheries and Aquatic Sciences*, **66(5)**:759-770.
- Hay, L. and G. McCabe, 2010: Hydrologic effects of climate change in the Yukon River Basin. *Climatic Change*, **100(3-4)**, 509-523.
- Haynie, A.C. and L. Pfeiffer, 2012: Why economics matters for understanding the effects of climate change on fisheries. *ICES Journal of Marine Science*, **69(7)**, 1160-1167.
- Hedenås, H., H. Olsson, C. Jonasson, J. Bergstedt, U. Dahlberg, and T.V. Callaghan, 2011: Changes in tree growth, biomass and vegetation over a 13-year period in the Swedish sub-Arctic. *Ambio*, **40(6)**, 672-682.
- Hedensted L., D., K. Sehested, T. Hellesen, *et al.*, 2012. Climate change adaptation in Denmark: enhancement through collaboration and meta-governance? *Local Environment*. 17 (6-7), 613-28.
- Heggberget, T, M., Gaare E. and Ball, J.P. 2010. Reindeer (*Rangifer tarandus*) and climate change: Importance of winter forage. *Rangifer* 22, 13-31.
- Heino, J., R. Virkkala, and H. Toivonen, 2009: Climate change and freshwater biodiversity: detected patterns, future trends and adaptations in northern regions. *Biological Reviews*, **84(1)**, 39-54.
- Heliasz, M., T. Johansson, A. Lindroth, M. Mölder, M. Mastepanov, T. Friborg, T.V. Callaghan, and T.R. Christensen, 2011: Quantification of C uptake in subarctic birch forest after setback by an extreme insect outbreak. *Geophysical Research Letters*, **38(1)**, 5.
- Helle, T.P. and L.M. Jaakkola, 2008: *Transitions in herd management of semi-domesticated reindeer in northern Finland* Akateeminen kirjakauppa, Helsinki, Finlande, pp. 81; 21-101.
- Henden, J.-., R.A. Ims, N.G. Yoccoz, P. Hellström, and A. Angerbjörn, 2010: Strength of asymmetric competition between predators in food webs ruled by fluctuating prey: The case of foxes in tundra. *Oikos*, **119(1)**, 27-34.
- Hezel, P.J., X. Zhang, C.M. Bitz, B.P. Kelly, and F. Massonnet, 2012: Projected decline in spring snow depth on Arctic sea ice caused by progressively later autumn open ocean freeze-up this century. *Geophysical Research Letters*, **39(17)**, - L17505.
- Higdon, J.W. and S.H. Ferguson, 2009: Loss of Arctic sea ice causing punctuated change in sightings of killer whales (*Orcinus orca*) over the past century. *Ecological Applications*, **19(5)**, 1365-1375.
- Hill, G.B. and G.H.R. Henry, 2011: Responses of High Arctic wet sedge tundra to climate warming since 1980. *Global Change Biology*, **17(1)**, 276-287.
- Hill, S. L., T. Phillips, and A. Atkinson, 2013: Potential Climate Change Effects on the Habitat of Antarctic Krill in the Weddell Quadrant of the Southern Ocean. *PloS one* **8.8**: e72246.
- Hill, Simeon L., Tony Phillips, and Angus Atkinson, 2013: Potential Climate Change Effects on the Habitat of Antarctic Krill in the Weddell Quadrant of the Southern Ocean. *PloS one* 8.8 (2013): e72246.
- Hinkel, K.M., B.M. Jones, W.R. Eisner, C.J. Cuomo, R.A. Beck, and R. Frohn, 2007: Methods to assess natural and anthropogenic thaw lake drainage on the western Arctic coastal plain of northern Alaska. **112**, F02S16
- Hinz, D.J., A.J. Poulton, M.C. Nielsdottir, S. Steigenberger, R.E. Korb, E.P. Achterberg, and T.S. Bibby, 2012: Comparative seasonal biogeography of mineralising nanoplankton in the Scotia Sea: *Emiliania huxleyi*, *Fragilariopsis* spp. and *Tetraparma pelagica*. *Deep-Sea Research Part II-Topical Studies in Oceanography*, **59**, 57-66.
- Hinzman, LD, Bettez N, Bolton WR, Chapin III FS, Dyurgerov M, Fastie C, Griffith B, Hollister RD, Hope A, Huntington HP *et al.*. 2005. **Evidence and implications of recent climate change in terrestrial regions of the Arctic**. *Climate Change*. 72:251-298.
- Hodgson, D.A. and J.P. Smol, 2008: High latitude paleolimnology. In: *Polar Lakes and Rivers - Limnology of Arctic and Antarctic Aquatic Ecosystems*. [Vincent, W.F. and J. Laybourn-Parry(eds.)]. Oxford University Press, Oxford, UK, pp. 43-64; 3.
- Hodgson, D.A., 2011: First synchronous retreat of ice shelves marks a new phase of polar deglaciation. *Proceedings of the National Academy of Sciences*, **108(47)**, 18859-18860.
- Hodgson, D.A., E. Verleyen, A.H. Squier, K. Sabbe, B.J. Keely, K.M. Saunders, and W. Vyverman, 2006b: Interglacial environments of coastal east Antarctica: comparison of MIS 1 (Holocene) and MIS 5e (Last Interglacial) lake-sediment records. *Quaternary Science Reviews*, **25**, 179-197.
- Hodgson, D.A., E. Verleyen, K. Sabbe, A.H. Squier, B.J. Keely, M.J. Leng, K.M. Saunders, and W. Vyverman, 2005: Late Quaternary climate-driven environmental change in the Larsemann Hills, East Antarctica, multi-proxy evidence from a lake sediment core. *Quaternary Research*, **64**, 83-99.

- Hofgaard, A., J.O. Lokken, L. Dalen, and H. Hytteborn, 2010: Comparing warming and grazing effects on birch growth in an alpine environment - a 10-year experiment. *Plant Ecol. Diversity Plant Ecology and Diversity*, **3(1)**, 19-27.
- Hollowed, A.B., B. Planque, and H. Loeng, 2013: Potential movement of fish and shellfish stocks from the Sub-Arctic to the Arctic Ocean. *Fisheries Oceanography*, **22(5)**, 355-370.
- Hovelsrud, G.K. and B. Smit (eds.), 2010: *Community Adaptation and Vulnerability in Arctic Regions*. Springer, pp. 353.
- Hovelsrud, G.K., B. Poppel, B.E.H. van Oort, and J.D. Reist, 2011: Arctic Societies, Cultures, and Peoples. In: *Snow, Water, Ice and Permafrost in the Arctic*. AMAP, Oslo, pp. 445-483.
- Høye, T.T., E. Post, H. Meltofte, N.M. Schmidt, and M.C. Forchhammer, 2007: Rapid advancement of spring in the High Arctic. *Current Biology*, **17(12)**, R449-R451.
- Hudson, J.M. and G.H. Henry, 2009: Increased plant biomass in a High Arctic heath community from 1981 to 2008. *Ecology*, **90(10)**, 2657-63.
- Hueffler, K., Parkinson, A.J., Gerlach, R. and Berner, J. , 2013. Zoonotic infections in Alaska: Disease prevalence, potential impact of climate change and recommended actions for earlier disease detection, research, prevention and control. *Int. J. Circumpolar Health*. **72: 19562**, <http://dx.doi.org/10.3402/ijch.v72i10.19562>
- Hughes, K.A. and P. Convey, 2010: The protection of Antarctic terrestrial ecosystems from inter- and intra-continental transfer of non-indigenous species by human activities: A review of current systems and practices. *Global Environmental Change*, **20(1)**, 96-112.
- Hughes-Hanks, J.M., L.G. Rickard, C. Panuska, J.R. Saucier, T.M. O'Hara, L. Dehn, and R.M. Rolland, 2005: Prevalence of *Cryptosporidium* spp. and *Giardia* spp. in five marine mammal species. *The Journal of Parasitology*, **91(5)**, 1225-1228.
- Hung, H., Kallenborn R., Breivik K., Su Y., Brorström-Lundén E., Olafsdottir K., Thorlacius J.M., Leppänen S., Bossi R., Skov H., Manø S., Patton G.W., Stern G., Sverko E., Fellin P. 2010. Atmospheric monitoring of organic pollutants in the Arctic under the Arctic Monitoring and Assessment Programme (AMAP): 1993-2006. *Science of the Total Environment*. **408**, 2854-2873.
- Hunt Jr., G.L., A.L. Blanchard, P. Boveng, P. Dalpadado, K.F. Drinkwater, L. Eisner, R.R. Hopcroft, K.M. Kovacs, B.L. Norcross, P. Renaud, M. Reigstad, M. Renner, H.R. Skjoldal, A. Whitehouse, and R.A. Woodgate, 2013: The Barents and Chukchi Seas: Comparison of two Arctic shelf ecosystems. *Journal of Marine Systems*, **109-110**, 43-68.
- Hunt, G.L., K.O. Coyle, L.B. Eisner, E.V. Farley, R.A. Heintz, F. Mueter, J.M. Napp, J.E. Overland, P.H. Ressler, S. Salo, and P.J. Stabeno, 2011: Climate impacts on eastern Bering Sea foodwebs: a synthesis of new data and an assessment of the Oscillating Control Hypothesis. *ICES Journal of Marine Science*, **68(6)**, 1230-1243.
- Hunter, C.M., H. Caswell, M.C. Runge, E.V. Regehr, S.C. Amstrup, and I. Stirling, 2010: Climate change threatens polar bear populations: a stochastic demographic analysis. *Ecology*, **91(10)**, 2883-2897.
- Huntington, H., L. Hamilton, C. Nicolson, R. Brunner, A. Lynch, A. Ogilvie, and A. Voinov, 2007: Toward understanding the human dimensions of the rapidly changing arctic system: insights and approaches from five HARC projects. *Regional Environmental Change*, **7(4)**, 173-186.
- Huntington, O.H. and A. Watson, 2012: Interdisciplinarity, Native Resilience, and How the Riddles Can Teach Wildlife Law in an Era of Rapid Climate Change. *Wicazo Sa Review*, **(2)**, 49.
- Hurst, T. P. E. R. Fernandez, J. T. Mathis, J. A. Miller, C. M. Stinson, E. F. Ahgeak, 2012: Resiliency of juvenile walleye pollock to projected levels of ocean acidification. *Aquatic Biology* **17**: 247-259
- Huse, G., I.Ellingsen, 2008: Capelin migrations and climate change - a modelling analysis. *Climatic Change*, **87**, 177-197.
- IAATO, 2012: International Association of Antarctic Tour Operators - Overview of Antarctic Tourism: 2011-12 Season and Preliminary Estimates for 2012-13 Season, ATCM XXXV Information Paper 39.
- Ignatowski, J. and J. Rosales. 2013. Identifying the exposure of two subsistence villages in Alaska to climate change using traditional ecological knowledge. *Climatic Change*. Online: DOI: 10.1007/s10584-013-0883-4.
- Ignatowski, J. and J. Rosales, 2012: *Aksik: Stories about Adaptation and subsistence: Native voices from the Frontlines of climate change* St. Lawrence University, Canton, N.Y.
- Ims, R.A., N.G. Yoccoz, and S.T. Killengreen, 2011: Determinants of lemming outbreaks. *Proceedings of the National Academy of Sciences of the United States of America*, **108(5)**, 1970-1974.
- Ims, R.A., N.G. Yoccoz, K.A. Bråthen, P. Fauchald, T. Tveraa, and V. Hausner, 2007: Can reindeer overabundance cause a trophic cascade? *Ecosystems*, **10(4)**, 607-622.

- Ingram, K.T., et al.: Opportunities and constraints for farmers of west Africa to use seasonal precipitation forecasts with Burkina Faso as a case study. *Agricultural Systems*, **74(3)**, (2002)
- IPCC, 2012: Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation. A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G.-K. Plattner, S.K. Allen, M. Tignor, and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, UK, and New York, NY, USA, 582 pp.
- Irons, D.B., T. Anker-Nilssen, A.J. Gaston, G.V. Byrd, K. Falk, G. Gilchrist, M. Hario, M. Hjernquist, Y.V. Krasnov, A. Mosbech, B. Olsen, A. Petersen, J.B. Reid, G.J. Robertson, H. Strom, and K.D. Wohl, 2008: Fluctuations in circumpolar seabird populations linked to climate oscillations. *Global Change Biology*, **14(7)**, 1455-1463.
- Iverson, S.J., I. Stirling, and S.L.C. Lang, 2006: Spatial and temporal variation in the diets of polar bears across the Canadian arctic: indicators of changes in prey populations and environment. In: *Top Predators in Marine Ecosystems*. [I. L. Boyd, S. Wanless and C.J. Camphuysen (ed.)]. Cambridge University Press, Cambridge, pp. 98.
- Jansson, M., A. Jonsson, A. Andersson, and J. Karlsson, 2010: Biomass and structure of planktonic communities along an air temperature gradient in subarctic Sweden. *Freshwater Biology*, **55(3)**, 691-700.
- Jarvis, T., N. Kelly, et al. (2010). "Acoustic characterisation of the broad-scale distribution and abundance of Antarctic krill (*Euphausia superba*) off East Antarctica (30-80 degrees E) in January-March 2006." *Deep-Sea Research Part II-Topical Studies in Oceanography* **57(9-10)**: 916-933.
- Jenouvrier, S., C. Barbraud, B. Cazelles, and H. Weimerskirch, 2005a: Modelling population dynamics of seabirds: importance of the effects of climate fluctuations on breeding proportions. *Oikos*, **108(3)**, 511-522.
- Jenouvrier, S., H. Caswell, C. Barbraud, M. Holland, J. Stroeve, and H. Weimerskirch, 2009: Demographic models and IPCC climate projections predict the decline of an emperor penguin population. *Proceedings of the National Academy of Sciences of the United States of America*, **106(6)**, 1844-1847.
- Jenouvrier, S., H. Weimerskirch, C. Barbraud, Y.H. Park, and B. Cazelles, 2005b: Evidence of a shift in the cyclicity of Antarctic seabird dynamics linked to climate. *Proceedings of the Royal Society B-Biological Sciences*, **272(1566)**, 887-895.
- Jenouvrier, S., M. Holland, J. Stroeve, C. Barbraud, H. Weimerskirch, M. Serreze, and H. Caswell, 2012: Effects of climate change on an emperor penguin population: analysis of coupled demographic and climate models. *Global Change Biology*, **18(9)**, 2756-2770.
- Jepsen, J.U., L. Kapari, S.B. Hagen, T. Schott, O.P.L. Vindstad, A.C. Nilssen, and R.A. Ims, 2011: Rapid northwards expansion of a forest insect pest attributed to spring phenology matching with sub-Arctic birch. *Global Change Biology*, **17(6)**, 2071-2083.
- Jepsen, J.U., S.B. Hagen, R.A. Ims, and N.G. Yoccoz, 2008: Climate change and outbreaks of the geometrids *Operophtera brumata* and *Epirrita autumnata* in subarctic birch forest: Evidence of a recent outbreak range expansion. *Journal of Animal Ecology*, **77(2)**, 257-264.
- Jernsletten, J.-. and K. Klokov, 2002: *Sustainable reindeer husbandry*. Arctic Council/Centre for Saami Studies, Tromsø, Norway.
- Ji, R., C.J. Ashjian, R.G. Campbell, C. Chen, G. Gao, C.S. Davis, G.W. Cowles, and R.C. Beardsley, 2012: Life history and biogeography of *Calanus* copepods in the Arctic Ocean: An individual-based modeling study. *Progress in Oceanography*, **96(1)**, 40-56.
- Johansson, C., V.A. Pohjola, C. Jonasson, and T.V. Callaghan, 2011: Multi-decadal changes in snow characteristics in sub-Arctic Sweden. *Ambio*, **40(6)**, 566-574.
- Joly, K., D.R. Klein, D.L. Verbyla, T.S. Rupp, and F.S. Chapin, 2011: Linkages between large-scale climate patterns and the dynamics of Arctic caribou populations. *Ecography*, **34(2)**, 345-352.
- Juday, G.P., 2009: Boreal Forests and Climate Change. In: *Oxford Companion to Global Change*. [Goudie, A. and D. Cuff (eds.)]. Oxford University Press, Oxford, UK, pp. 75-84.
- Kahru, M., V. Brotas, M. Manzano-Sarabia, and B.G. Mitchell, 2011: Are phytoplankton blooms occurring earlier in the Arctic? *Global Change Biology*, **17(4)**, 1733-1739.
- Kaplan, J.O. and M. New, 2006: Arctic climate change with a 2°C global warming: Timing, climate patterns and vegetation change. *Climatic Change*, **79(3-4)**, 213-241.
- Karnovsky, N., A. Harding, W. Walkusz, S. Kwasniewski, I. Goszczko, J. Wiktor, H. Routti, A. Bailey, L. McFadden, Z. Brown, G. Beaugrand, and D. Gremillet, 2010: Foraging distributions of little auks *Alle alle*

- across the Greenland Sea: implications of present and future Arctic climate change. *Marine Ecology-Progress Series*, **415**, 283-293.
- Kausrud, K.L., Atle Mysterud, Harald Steen, Jon Olav Vik, Eivind Østbye, Bernard Cazelles, Erik Framstad, Anne Maria Eikeset, Ivar Mysterud, Torstein Solhøy and & Nils Chr. Stenseth 2008. Linking climate change to lemming cycles. *Nature* 456, 93-97.
- Kawaguchi, S., A. Ishida, R. King, B. Raymond, N. Waller, A. Constable, S. Nicol, M. Wakita, and A. Ishimatsu. 2013. Risk maps for Antarctic krill under projected Southern Ocean acidification. *Nature Climate Change*. doi:10.1038/nclimate1937
- Kawaguchi, S., H. Kurihara, R. King, L. Hale, T. Berli, J.P. Robinson, A. Ishida, M. Wakita, P. Virtue, S. Nicol, and A. Ishimatsu, 2011: Will krill fare well under Southern Ocean acidification? *Biology Letters*, **7(2)**, 288-291.
- Kawaguchi, S., S. Nicol, and A.J. Press, 2009: Direct effects of climate change on the Antarctic krill fishery. *Fisheries Management and Ecology*, **16(5)**, 424-427.
- Keskitalo, E., 2008: Vulnerability and adaptive capacity in forestry in northern Europe: a Swedish case study. *Climatic Change*, **87(1)**, 219-234.
- Keskitalo, E., 2009: Governance in vulnerability assessment: the role of globalising decision-making networks in determining local vulnerability and adaptive capacity. *Mitigation and Adaptation Strategies for Global Change*, **14(2)**, 185-201.
- Kharuk, V.I., K.J. Ranson, S.T. Im, and M.M. Naurzbaev, 2006: Forest-tundra larch forests and climatic trends. *Russian Journal of Ecology*, **37(5)**, 291-298.
- Khon, V., I. Mokhov, M. Latif, V. Semenov, and W. Park, 2010: Perspectives of Northern Sea Route and Northwest Passage in the twenty-first century. *Climatic Change*, **100(3)**, 757-768.
- Kilpelainen, A., H. Gregow, H. Strandman, S. Kellomaki, A. Venalainen and H. Peltola, 2010. Impacts of climate change on the risk of snow-induced forest damage in Finland. *Climate Change*, 99: 193-209.
- Kitti, H., B.C. Forbes, and J. Oksanen, 2009: Long- and short-term effects of reindeer grazing on tundra wetland vegetation. *Polar Biology*, **32(2)**, 253-261.
- Kokelj, S.V., B. Zajdlik, and M.S. Thompson, 2009: The impacts of thawing permafrost on the chemistry of lakes across the subarctic boreal-tundra transition, Mackenzie Delta region, Canada. *Permafrost and Periglacial Processes*, **20(2)**, 185-199.
- Kotwicki, S. and R.R. Lauth. 2013: Detecting temporal trends and environmentally-driven changes in the spatial distribution of groundfishes and crabs on the eastern Bering Sea shelf. *Deep-Sea Research Part II: Topical Studies in Oceanography*, **94**, 155-175.
- Kovacs, K.M. and C. Lydersen, 2008: Climate change impacts on seals and whales in the North Atlantic Arctic and adjacent shelf seas. *Science Progress*, **91(Pt 2)**, 117-150.
- Kovacs, K.M., C. Lydersen, J.E. Overland and S.E. Moore. 2010. Impacts of changing sea-ice conditions on Arctic marine mammals. *Mar. Biodivers.*, **41**, 181-194.
- Krebs, C.J., 2011: Of lemmings and snowshoe hares: The ecology of northern Canada. *Proceedings of the Royal Society B: Biological Sciences*, **278(1705)**, 481-489.
- Kristiansen, T., K.F. Drinkwater, R.G. Lough, and S. Sundby, 2011: Recruitment variability in North Atlantic cod and match-mismatch dynamics. *PLoS ONE*, **6(3)**, e17456.
- Krupnik, I. and G.C. Ray, 2007: Pacific walruses, indigenous hunters, and climate change: Bridging scientific and indigenous knowledge. *Deep Sea Research Part II: Topical Studies in Oceanography*, **54(23-26)**, 2946-2957.
- Krupnik, I., Allison, I., Bell, R., Cutler, P., Hik, D., López-Martínez, J., Rachold, V., Sarukhanian, E., Summerhayes, C, eds. 2011: Understanding Earth's Polar Challenges: International Polar Year 2007-2008. Summary by the IPY Joint Committee. University of the Arctic publications series (4). University of the Arctic and ICSU/WMO Joint Committee for International Polar Year 2007-2008, Rovaniemi, Finland. ISBN 978-1-896445-55-7
- Krupnik, I., Aporta, C., Gearheard, S., Kielsen Holm, L., and Laidler, G. 2010. SIKU: Knowing our ice – Documenting Inuit sea ice knowledge and use. New York: Springer, 501pp
- Krupnik, I., Ed. and D. Jolly Ed. (eds.), 2002: *The Earth Is Faster Now: Indigenous Observations of Arctic Environmental Change. Frontiers in Polar Social Science*. Arctic Research Consortium of the United States, 3535 College Road, Suite 101, Fairbanks, AK 99709, pp. 383.
- Kullman, L. and L. Öberg, 2009: Post-Little Ice Age tree line rise and climate warming in the Swedish Scandes: A landscape ecological perspective. *Journal of Ecology*, **97(3)**, 415-429.

- Kumpula, T., A. Pajunen, E. Kaarlejärvi, B.C. Forbes, and F. Stammler, 2011: Land use and land cover change in Arctic Russia: Ecological and social implications of industrial development. *Global Environmental Change*, **21(2)**, 550-562.
- Kumpula, T., B.C. Forbes, F. Stammler, and N. Meschtyb, 2012: Dynamics of a coupled system: multi-resolution remote sensing in assessing social-ecological responses during 25 years of gas field development in Arctic Russia. *Remote Sensing*, **4**, 1046-1068.
- Lack, D. A. and Corbett, J. J. 2012: Black carbon from ships: a review of the effects of ship speed, fuel quality and exhaust gas scrubbing. *Atmos. Chem. Phys.*, **12**, 3985-4000, doi:10.5194/acp-12-3985-2012, 2012.
- Laidler, G.J., T. Hirose, M. Kapfer, T. Ikummaq, E. Joamie, and P. Elee, 2011: Evaluating the Floe Edge Service: how well can SAR imagery address Inuit community concerns around sea ice change and travel safety? *Canadian Geographer / Le Géographe Canadien*, **55(1)**, 91-107.
- Laidre, K.L., I. Stirling, L.F. Lowry, Å Wiig, M.P. Heide-JÃ,rgensen, and S.H. Ferguson, 2008: quantifying the sensitivity of arctic marine mammals to climate-induced habitat change. *Ecological Applications*, **18**, S97-S125.
- Landerer, F.W., J.O. Dickey, and A. Guntner, 2010: Terrestrial water budget of the Eurasian pan-Arctic from GRACE satellite measurements during 2003–2009. **115**, D23115.
- Lantz, T.C. and S.V. Kokelj, 2008: Increasing rates of retrogressive thaw slump activity in the Mackenzie Delta region, N.W.T., Canada. *Geophysical Research Letters*, **35(6)**, L06502.
- Larsen, J.N. and L. Huskey, 2010: Material wellbeing in the Arctic. In: *Arctic Social Indicators*. [Larsen, J.N., P. Schweitzer, and G. Fondahl(eds.)]. TemaNord, pp. 47-66.
- Larsen, J.N., P. Schweitzer, and G. Fondahl, 2010: **Arctic Social Indicators**. *TemaNord*, **519**, 160.
- Laurion, I., W.F. Vincent, S. MacIntyre, L. Retamal, C. Dupont, P. Francus, and R. Pienitz, 2010: Variability in greenhouse gas emissions from permafrost thaw ponds. *Limonology and Oceanography*, **55(1)**, 115.
- Lee, S., M. Jin, and T. Whitledge, 2010: Comparison of bottom sea-ice algal characteristics from coastal and offshore regions in the Arctic Ocean. *Polar Biology*, **33(10)**, 1331-1337.
- Lesack, L.F.W. and P. Marsh, 2007: Lengthening plus shortening of river-to-lake connection times in the Mackenzie River Delta respectively via two global change mechanisms along the arctic coast. *Geophysical Research Letters*, **34(23)**, L23404.
- Lesack, L.F.W. and P. Marsh, 2010: River-to-lake connectivities, water renewal, and aquatic habitat diversity in the Mackenzie River Delta. *Water Resources Research*, **46(12)**, W12504.
- Levintova, M, W.I. Zapol, N. Engmann (Eds.) (2010) Behavioral and Mental Health Research in the Arctic: A Strategy Setting Meeting. *Inter J. Circumpolar Health Suppl.* **5**. [http://www.ijch.fi/CHS/CHS_2010\(5\).pdf](http://www.ijch.fi/CHS/CHS_2010(5).pdf)
- Lewis, T. and S.F. Lamoureux, 2010: Twenty-first century discharge and sediment yield predictions in a small high Arctic watershed. *Global and Planetary Change*, **71(1-2)**, 27-41.
- Li, W.K.W., F.A. McLaughlin, C. Lovejoy, and E.C. Carmack, 2009: Smallest algae thrive as the Arctic Ocean freshens. *Science*, **326(5952)**, 539.
- Lima, M., S.A. Estay, 2013: Warming effects in the western Antarctic Peninsula ecosystem: the role of population dynamic models for explaining and predicting penguin trends. *Population Ecology*, DOI 10.1007/s10144-013-0386-1
- Lind, S. and R.B. Ingvaldsen, 2012: Variability and impacts of Atlantic Water entering the Barents Sea from the north. *Deep Sea Research Part I: Oceanographic Research Papers*, **62**, 70-88.
- Lindgren, E. and R. Gustafson, 2001: Tick-borne encephalitis in Sweden and climate change. *The Lancet*. **358**. Issue 9275. p 16-18
- Lindholt, Lars. (2006). "Arctic Natural Resources in a Global Perspective." S. Glomsrød and Iulie Aslaksen (eds.), *The Economy of the North*, pp. 27-39. Oslo: Statistics Norway. www.ssb.no
- Livingston, P., K. Aydin, J.L. Boldt, A.B. Hollowed, and J.M. Napp, 2011: Alaska Marine Fisheries Management: Advancements and Linkages to Ecosystem Research. In: *Ecosystem Based Management: An Evolving Perspective*. [Belgrano, A. and C. Fowler(eds.)]. Cambridge University Press, Cambridge, pp. 113-152.
- Lloyd, A.H., A.G. Bunn, and L. Berner, 2011: A latitudinal gradient in tree growth response to climate warming in the Siberian taiga. *Global Change Biology*, **17(5)**, 1935-1945.
- Long, W. C., K. M. Swiney, C. Harris, H. N. Page, and R. J. Foy. 2013. Effects of ocean acidification on juvenile red king crab (*Paralithodes camtschaticus*) and Tanner crab (*Chionoecetes bairdi*) growth, condition, calcification, and survival. *PLoS ONE* **8**:e60959.
- Lorenzen, E.D., D. Nogués-Bravo, L. Orlando, J. Weinstock, J. Binladen, K.A. Marske, A. Ugan, M.K. Borregaard, M.T.P. Gilbert, R. Nielsen, S.Y.W. Ho, T. Goebel, K.E. Graf, D. Byers, J.T. Stenderup, M. Rasmussen, P.F.

- Campos, J.A. Leonard, K.-. Koepfli, D. Froese, G. Zazula, T.W. Stafford Jr., K. Aaris-Sørensen, P. Batra, A.M. Haywood, J.S. Singarayer, P.J. Valdes, G. Boeskorov, J.A. Burns, S.P. Davydov, J. Haile, D.L. Jenkins, P. Kosintsev, T. Kuznetsova, X. Lai, L.D. Martin, H.G. McDonald, D. Mol, M. Meldgaard, K. Munch, E. Stephan, M. Sablin, R.S. Sommer, T. Sipko, E. Scott, M.A. Suchard, A. Tikhonov, R. Willerslev, R.K. Wayne, A. Cooper, M. Hofreiter, A. Sher, B. Shapiro, C. Rahbek, and E. Willerslev, 2011: Species-specific responses of Late Quaternary megafauna to climate and humans. *Nature*, **479(7373)**, 359-364.
- Lund, D.H., K. Sehested, T. Helleisen and V. Nellemann, 2012: Climate change adaptation in Denmark: enhancement through collaboration and meta-governance? *Local Environment*. 17 (6-7), 613-28, DOI:10.1080/13549839.2012.678318
- Lynch, H.J., R. Naveen, P.N. Trathan, and W.F. Fagan, 2012: Spatially integrated assessment reveals widespread changes in penguin populations on the Antarctic Peninsula. *Ecology*, **93(6)**, 1367-1377.
- Lynn, K., J. Daigle, J. Hoffman, F. Lake, N. Michelle, D. Ranco, D. C. Viles C., G. Voggesser, P. Williams, 2013. The impacts of climate change on tribal traditional foods. *Climatic Change*. doi:10.1007/s10584-013-0736-1.
- Lyons, W.B., J. Laybourn-Parry, K.A. Welch, and J.C. Priscu, 2006: Antarctic lake systems and climate change. In: *Trends in Antarctic Terrestrial and Limnetic Ecosystems. Antarctica as a Global Indicator*. [Bergstrom, D.M., P. Convey, and A.H.L. Huiskes(eds.)]. Springer, Netherlands, pp. 273-295.
- Ma, J., H. Hung, C. Tian, and R. Kallenborn. 2001. Revolatilization of persistent organic pollutants in the Arctic induced by climate change. *Nature Climate Change*. 1. 255-260. Doi:1038/nclimate1167.
- Ma, J., Hung, H., Tian, C and Kallenborn, R. 2011. Revolatilization of persistent organic pollutants in
- Maccracken, J.G., 2012: Pacific Walrus and climate change: observations and predictions. *Ecology and Evolution*, **2(8)**, 2072-2090.
- MacDonald, G.M., K.V. Kremenetski, and D.W. Beilman, 2008a: Climate change and the northern Russian treeline zone. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **363(1501)**, 2285-2299.
- Macias-Fauria, M., B.C. Forbes, P. Zetterberg, and T. Kumpula, 2012: Eurasian Arctic greening reveals teleconnections and the potential for novel ecosystems. *Nature Climate Change*, **2**, 613-618, doi:10.1038/nclimate1558.
- Mack, M.C., M.S. Bret-Harte, T.N. Hollingsworth, R.R. Jandt, E.A.G. Schuur, G.R. Shaver, and D.L. Verbyla, 2011: Carbon loss from an unprecedented Arctic tundra wildfire. *Nature*, **475(7357)**, 489-492.
- Mackey, A.P., A. Atkinson, S.L. Hill, P. Ward, N.J. Cunningham, N.M. Johnston, and E.J. Murphy, 2012: Antarctic macrozooplankton of the southwest Atlantic sector and Bellingshausen Sea: Baseline historical distributions (Discovery Investigations, 1928-1935) related to temperature and food, with projections for subsequent ocean warming. *Deep-Sea Research Part Ii-Topical Studies in Oceanography*, 59, 130-146.
- Magga, O.H., S.D. Mathiesen, R.W. Corell, and A. Oskal, 2011: *Reindeer Herding, Traditional Knowledge And Adaptation To Climate Change And Loss Of Grazing Land. A project led by Norway and Association of World Reindeer Herders (WRH) in Arctic Council, Sustainable Development Working Group (SDWG)*. In: International Centre for Reindeer Husbandry Report 1. International Centre for Reindeer Husbandry and Association of World Reindeer Herders., Fagtrykk Idé AS, Alta, Norway.
- Mahoney, A., S. Gearheard, T. Oshima, and T. Qillaq, 2009: Sea Ice Thickness Measurements from a Community-Based Observing Network. *Bulletin of the American Meteorological Society*, **90(3)**, 370-377.
- Maldonado, J.K., C. Shearer, R. Bronen, K. Peterson and H. Lazrus. 2013. The impact of climate change on tribal communities in the US: displacement, relocation, and human rights. *Climatic Change*. doi: 10.1007/s10584-013-0746-z.
- Mamet, S.D. and G.P. Kershaw, 2012: Subarctic and alpine tree line dynamics during the last 400 years in north-western and central Canada. *Journal of Biogeography*, **39(5)**, 855-868.
- Marsh, P., M. Russell, S. Pohl, H. Haywood, and C. Onclin, 2009: Changes in thaw lake drainage in the Western Canadian Arctic from 1950 to 2000. *Hydrological Processes*, **23(1)**, 145-158.
- Massom, R.A. and S.E. Stammerjohn, 2010: Antarctic sea ice change and variability – Physical and ecological implications. *Polar Science*, **4(2)**, 149-186.
- Matrai, P.A., E. Olson, S. Suttles, V. Hill, L.A. Codispoti, B. Light, and M. Steele, 2013: Synthesis of primary production in the Arctic Ocean: I. Surface waters, 1954–2007. *Progress in Oceanography*, **110**, 93–106.
- Matthews, C., S. Luque, S. Petersen, R. Andrews, and S. Ferguson, 2011: Satellite tracking of a killer whale (*Orcinus orca*) in the eastern Canadian Arctic documents ice avoidance and rapid, long-distance movement into the North Atlantic. *Polar Biology*, **34(7)**, 1091-1096.

- Maynard, N.G. 2006: Satellites, settlements, and human health. In: Remote Sensing of Human Settlements (Ridd, M and J.D. Hipple, Eds) Manual of Remote Sensing. American Society of Photogrammetry and Remote Sensing. 3rd Edition, **5**. Bethesda, MD. Pp. 379-399.
- Maynard, N.G. and G.A. Conway, 2007: A view from above: use of satellite imagery to enhance our understanding of potential impacts of climate change on human health in the Arctic. *Alaska Medicine*, **49(2 Suppl)**, 38-43.
- Maynard, N.G., A. Oskal, J.M. Turi, S.D. Mathiesen, I.M.G. Eira, B. Yurchak, V. Etylin, and J. Gebelein, 2011: Impacts of Arctic Climate and Land Use Changes on Reindeer Pastoralism: Indigenous Knowledge & Remote Sensing. In: Eurasian Arctic Land Cover and Land Use in a Changing Climate. [Gutman, G. (ed.)]. Springer, pp. 177-205.
- McLaughlin, J.B., A. DePaola, C.A. Bopp, K.A. Martinek, N.P. Napolilli, C.G. Allison, S.L. Murray, E.C. Thompson, M.M. Bird, and J.P. Middaugh, 2005: Outbreak of *Vibrio parahaemolyticus* Gastroenteritis Associated with Alaskan Oysters. *N Engl J Med*, **353(14)**, 1463-1470.
- McLeod, D., G. Hallegraeff, G. Hosie, and A. Richardson. 2012. Climate-driven range expansion of the red-tide dinoflagellate *Noctiluca scintillans* into the Southern Ocean. *Journal of Plankton Research* 34:332-337.
- McNamara, J.P. and D.L. Kane, 2009: The impact of a shrinking cryosphere on the form of arctic alluvial channels. *Hydrological Processes*, **23(1)**, 159-168.
- McNeeley, S.M., 2012: Examining barriers and opportunities for sustainable adaptation to climate change in Interior Alaska. *Climatic Change*, **111(3-4)**, 835-857.
- Mehlum, F. 2012: Effects of sea ice on breeding numbers and clutch size of a high arctic population of the common eider *Somateria mollissima*. *Polar Science*, **6**, 143-153.
- Melbourne-Thomas, J., A. Constable, S. Wotherspoon, and B. Raymond, 2013: Testing paradigms of ecosystem change under climate warming in Antarctica. *Plos One*, **8(2)**, e55093.
- Melles, M., J. Brigham-Grette, O. Glushkova, P. Minyuk, N. Nowaczyk, and H. Hubberten, 2007: Sedimentary geochemistry of core PG1351 from Lake El'gygytyn—a sensitive record of climate variability in the East Siberian Arctic during the past three glacial–interglacial cycles. *Journal of Paleolimnology*, **37(1)**, 89-104.
- Merino, G., M. Barange, J.L. Blanchard, J. Harle, R. Holmes, I. Allen, E.H. Allison, M.C. Badjeck, N.K. Dulvy, J. Holt, S. Jennings, C. Mullon, and L.D. Rodwell, 2012: Can marine fisheries and aquaculture meet fish demand from a growing human population in a changing climate? *Global Environmental Change*, **22(4)**, 795–806.
- Meschtyb, N., Forbes, B., Kankaanpää, P., 2010: Social Impact Assessment along Russia's Northern Sea Route: Petroleum Transport and the Arctic Operational Platform (ARCOP). *Arctic*, **58(3)**.
- Mesquita, P.S., F.J. Wrona, and T.D. Prowse, 2010: Effects of retrogressive permafrost thaw slumping on sediment chemistry and submerged macrophytes in Arctic tundra lakes. *Freshwater Biology*, **55(11)**, 2347-2358.
- Metje, M. and P. Frenzel, 2007: Methanogenesis and methanogenic pathways in a peat from subarctic permafrost. *Environmental Microbiology*, **9(4)**, 954-964.
- Miller, F.L. and S.J. Barry, 2009: Long-Term Control of Peary Caribou Numbers by Unpredictable, Exceptionally Severe Snow or Ice Conditions in a Non-equilibrium Grazing System. *Arctic*, **62(2)**, 175-189.
- Miller, W., S.C. Schuster, A.J. Welch, A. Ratan, O.C. Bedoya-Reina, F. Zhao, H.L. Kim, R.C. Burhans, D.I. Drautz, N.E. Wittekindt, L.P. Tomsho, E. Ibarra-Laclette, L. Herrera-Estrella, E. Peacock, S. Farley, G.K. Sage, K. Rode, M. Obbard, R. Montiel, L. Bachmann, Ó Ingólfsson, J. Aars, T. Mailund, Ø Wiig, S.L. Talbot, and C. Lindqvist, 2012: Polar and brown bear genomes reveal ancient admixture and demographic footprints of past climate change. *Proceedings of the National Academy of Sciences*, **109 (36)**, E2382-E2390
- Milliman, J.D., K.L. Farnsworth, P.D. Jones, K.H. Xu, and L.C. Smith, 2008: Climatic and anthropogenic factors affecting river discharge to the global ocean, 1951–2000. *Global and Planetary Change*, **62(3-4)**, 187-194.
- Milner, A.M., L.E. Brown, and D.M. Hannah, 2009: Hydroecological response of river systems to shrinking glaciers. *Hydrological Processes*, **23(1)**, 62-77.
- Moe, B., L. Stempniewicz, D. Jakubas, F. Angelier, O. Chastel, F. Dinessen, G.W. Gabrielsen, F. Hanssen, N.J. Karnovsky, B. Ronning, J. Welcker, K. Wojczulanis-Jakubas, and C. Bech, 2009: Climate change and phenological responses of two seabird species breeding in the high-Arctic. *Marine Ecology Progress Series*, **393**.
- Moen, J., K. Aune, L. Edenius, and A. Angerbjörn, 2004: Potential effects of climate change on treeline position in the Swedish mountains. *Ecology and Society*, **9(1)**.
- Mokhov, I.I. and V.C. Khon, 2008: Assessment of the Northern Sea Route perspectives under climate changes on the basis of simulations with the climate models ensemble. In: Natural processes in Polar regions, Part 2. [V.M. Kotlijakov, (ed.)] IG RAS: Moscow. p. 20-27.

- Molenaar, E.J., 2009: Climate Change and Arctic Fisheries. In: *Climate Governance in the Arctic*. [Koivurova, T., E.C.H. Keskkitalo, and N. Bankes(eds.)]. Springer Netherlands, pp. 145-169.
- Moline, M.A., N.J. Karnovsky, Z. Brown, G.J. Divoky, T.K. Frazer, C.A. Jacoby, J.J. Torrese, and W.R. Fraser, 2008: High latitude changes in ice dynamics and their impact on polar marine ecosystems. *Year in Ecology and Conservation Biology* 2008, **1134**, 267-319.
- Molnar, P.K., A.E. Derocher, T. Klanjscek, and M.A. Lewis, 2011: Predicting climate change impacts on polar bear litter size. **2**, 186.
- Monnett, C. and J. Gleason, 2006: Observations of mortality associated with extended open-water swimming by polar bears in the Alaskan Beaufort Sea. *Polar Biology*, **29(8)**, 681-687.
- Montes-Hugo, M., S.C. Doney, H.W. Ducklow, W. Fraser, D. Martinson, S.E. Stammerjohn, and O. Schofield, 2009: Recent Changes in Phytoplankton Communities Associated with Rapid Regional Climate Change Along the Western Antarctic Peninsula. *Science*, **323(5920)**, 1470-1473.
- Moore, R.D., S.W. Fleming, B. Menounos, R. Wheate, A. Fountain, K. Stahl, K. Holm, and M. Jakob, 2009: Glacier change in western North America: influences on hydrology, geomorphic hazards and water quality. *Hydrological Processes*, **23(25)**, 42-61.
- Moore, S.E. and H.P. Huntington, 2008: Arctic marine mammals and climate change: impacts and resilience. *Ecological Applications : A Publication of the Ecological Society of America*, **18(2 Suppl)**, s157-165.
- Moore, S.E., 2008: Marine mammals as ecosystem sentinels. *Journal of Mammology*, **89(3)**, 534-540.
- Morán, X.A.G., Á López-Urrutia, A. Calvo-Díaz, and W.K.W. Li, 2010: Increasing importance of small phytoplankton in a warmer ocean. *Global Change Biology*, **16(3)**, 1137-1144.
- Moy, A.D., W.R. Howard, S.G. Bray, and T.W. Trull, 2009: Reduced calcification in modern Southern Ocean planktonic foraminifera. *Nature Geoscience*, **2(4)**, 276-280.
- Mueter, F.J. and M.A. Litzow, 2008: Sea ice retreat alters the biogeography of the Bering Sea continental shelf. *Ecological Applications*, **18(2)**, 309-320.
- Mueter, F.J., J.L. Boldt, B.A. Megrey, and R.M. Peterman, 2007: Recruitment and survival of northeast Pacific Ocean fish stocks: temporal trends, covariation, and regime shifts. *Canadian Journal of Fisheries and Aquatic Sciences*, **64(6)**, 911-927.
- Mueter, F.J., N.A. Bond, J.N. Ianelli, A.B. Hollowed. 2011: Expected declines in recruitment of walleye pollock (*Theragra chacoogramma*) in the eastern Bering Sea under future climate change. *ICES Journal of Marine Science*, **68(6)**, 1284-1296.
- Mundy, P.R., and D.F. Evenson, 2011: Environmental controls of phenology of high-latitude Chinook salmon populations of the Yukon River, North America, with application to fishery management. *ICES Journal of Marine Science*, **68(6)**, 1155-1164.
- Murphy, E. J., E. E. Hofmann, J. L. Watkins, N. M. Johnston, A. Pinones, T. Ballerini, S. L. Hill, P. N. Trathan, G. A. Tarling, R. A. Cavanagh, E. F. Young, S. E. Thorpe, and P. Fretwell. 2013: Comparison of the structure and function of Southern Ocean regional ecosystems: The Antarctic Peninsula and South Georgia. *Journal of Marine Systems* **109:22-42**.
- Murphy, E., J. Watkins, P. Trathan, K. Reid, M. Meredith, S. Thorpe, N. Johnston, A. Clarke, G. Tarling, M. Collins, J. Forcada, R. Shreeve, A. Atkinson, R. Korb, M. Whitehouse, P. Ward, P. Rodhouse, P. Enderlein, A. Hirst, A. Martin, S. Hill, I. Staniland, D. Pond, D. Briggs, N. Cunningham, and A. Fleming. 2007. Spatial and temporal operation of the Scotia Sea ecosystem: a review of large-scale links in a krill centred food web. *Philosophical Transactions of the Royal Society B-Biological Sciences* **362**:113-148.
- Murphy, E.J., J.L. Watkins, P.N. Trathan, K. Reid, M.P. Meredith, S.L. Hill, S.E. Thorpe, N.M. Johnston, A. Clarke, G.A. Tarling, M.A. Collins, J. Forcada, A. Atkinson, P. Ward, I.J. Staniland, D.W. Pond, R.A. Cavanagh, R.S. Shreeve, R.E. Korb, M.J. Whitehouse, P.G. Rodhouse, P. Enderlein, A.G. Hirst, A.R. Martin, D.R. Briggs, N.J. Cunningham, and A.H. Fleming, 2012b: Spatial and Temporal Operation of the Scotia Sea Ecosystem. In: *Antarctic Ecosystems*. [Rogers, A., N. Johnston, E. Murphy, and A. Clarke (eds.)]. Blackwell Publishing, Ltd, pp. 160-212.
- Murphy, E.J., R.D. Cavanagh, E.E. Hofmann, S.L. Hill, A.J. Constable, D.P. Costa, M.H. Pinkerton, N.M. Johnston, P.N. Trathan, J.M. Klinck, D.A. Wolf-Gladrow, K.L. Daly, O. Maury, and S.C. Doney, 2012a: Developing integrated models of Southern Ocean food webs: Including ecological complexity, accounting for uncertainty and the importance of scale. *Progress in Oceanography*, **102**, 74-92.
- Myers-Smith, I.H., B.C. Forbes, M. Wilmking, M. Hallinger, T. Lantz, D. Blok, K.D. Tape, M. MacIas-Fauria, U. Sass-Klaassen, E. Lévesque, S. Boudreau, P. Ropars, L. Hermanutz, A. Trant, L.S. Collier, S. Weijers, J.

- Rozema, S.A. Rayback, N.M. Schmidt, G. Schaepman-Strub, S. Wipf, C. Rixen, C.B. Ménard, S. Venn, S. Goetz, L. Andreu-Hayles, S. Elmendorf, V. Ravolainen, J. Welker, P. Grogan, H.E. Epstein, and D.S. Hik, 2011a: Shrub expansion in tundra ecosystems: Dynamics, impacts and research priorities. *Environmental Research Letters*, **6(4)**.
- Myers-Smith, I.H., D.S. Hik, C. Kennedy, D. Cooley, J.F. Johnstone, A.J. Kenney, and C.J. Krebs, 2011b: Expansion of canopy-forming willows over the twentieth century on Herschel Island, Yukon Territory, Canada. *Ambio*, **40(6)**, 610-623.
- Nakashima, D.J., McLean, K.G., Thulstrup, H., Ameyali, R.C. and Rubis, J. 2012. Indigenous Knowledge, Marginalized Peoples and Climate Change: Foundations for Assessment and Adaptation. Technical Report prepared for the Intergovernmental Panel on Climate Change – Working Group II. UNESCO/UNU: Paris/Darwin. 82 pp.
- Nayha, S., 2005: Environmental temperature and mortality. *International Journal of Circumpolar Health*, **64(5)**, 451-458.
- Nicol, S, Pauly, T, Bindoff, NL, Wright, S, Thiele, D, Hosie, GW, Strutton, PG, Woehler, E (2000a) Ocean circulation off east Antarctica affects ecosystem structure and sea-ice extent. *Nature* **406**:504-507
- Nicol, S. and A. Brierley, 2010: Through a glass less darkly—new approaches for studying the distribution, abundance and biology of euphausiids. *Deep-Sea Research II*, **57**, 496–507.
- Nicol, S., A. Worby, and R. Leaper, 2008: Changes in the Antarctic sea ice ecosystem: potential effects on krill and baleen whales. *Marine and Freshwater Research*, **59**, 361-382.
- Nicol, S., Constable, A.J., Pauly, T. (2000b) Estimates of circumpolar abundance of Antarctic krill based on recent acoustic density measurements. *CCAMLR Science* **7:87-99**
- Nicol, S., J. Foster, and S. Kawaguchi, 2011: The fishery for Antarctic krill – recent developments. *Fish and Fisheries*, **13(1)**, DOI: 10.1111/j.1467-2979.2011.00406.x.
- Nikanorov, A., V. Bryzgalov, and G. Chernogaeva, 2007: Anthropogenically modified natural background and its formation in the Russian freshwater ecosystems. *Russian Meteorology and Hydrology*, **32(11)**, 698-710.
- NorAcia. 2010. Klimaendringer i Norsk Arktis – Konsekvenser for Livet i Nord. Norsk Polarinstitutt Rapportserie 136
- NRTEE. Canada. National Round Table on the Environment and the Economy. 2009. True North: Adapting Infrastructure to Climate Change in Northern Canada. Ottawa. NRTEE
- Nuttall, M., F. Berkes, B. Forbes, G. Kofinas, T. Vlassova, and G. Wenzel, 2005: Hunting, herding, fishing and gathering: indigenous peoples and renewable resource use in the Arctic. *Arctic Climate Impact Assessment*, (January), 649-690.
- Ogden, N.H., C. Bouchard, K. Kurtenbach, G. Margos, L.R. Lindsay, L. Trudel, S. Nguon, and F. Milord, 2010: Active and Passive Surveillance and Phylogenetic Analysis of *Borrelia burgdorferi* Elucidate the Process of Lyme Disease Risk Emergence in Canada. *Environ Health Perspect*, **118(7)**.
- Olech, M. and K.J. Chwedorzewska, 2011: Short Note: The first appearance and establishment of an alien vascular plant in natural habitats on the forefield of a retreating glacier in Antarctica. *Antarctic Science*, **23(02)**, 153.
- Olivier Gilg, Kit M. Kovacs, Jon Aars, Jerome Fort, Gilles Gauthier, David Grémillet, Rolf A. Ims, Hans Meltofte, Jérôme Moreau, Eric Post, Niels Martin Schmidt, Glenn Yannic, and Loïc Bollache. 2012. Climate change and the ecology and evolution of Arctic vertebrates. *Annals of the New York Academy of Sciences* 1249, 166-190
- Olli, K., P. Wassmann, M. Reigstad, T.N. Ratkova, E. Arashkevich, A. Pasternak, P.A. Matrai, J. Knulst, L. Tranvik, R. Klais, and A. Jacobsen, 2007: The fate of production in the central Arctic Ocean – top-down regulation by zooplankton expatriates? *Progress in Oceanography*, **72(1)**, 84-113.
- Olofsson, A., O. Danell, P. Forslund, and B. Hman, 2011: Monitoring changes in lichen resources for range management purposes in reindeer husbandry. *Ecological Indicators*, **11(5)**, 1149-1159.
- Olofsson, J., H. Tømmervik, and T.V. Callaghan, 2012: Vole and lemming activity observed from space: vegetation cycles at a regional scale. *Nature Climate Change*, published online April 2012.
- Olofsson, J., L. Oksanen, T. Callaghan, P.E. Hulme, T. Oksanen, and O. Suominen, 2009: Herbivores inhibit climate-driven shrub expansion on the tundra. *Global Change Biology*, **15(11)**, 2681-2693.
- Ormseth, O.A. and B.L. Norcross, 2009: Causes and consequences of life-history variation in North American stocks of Pacific cod. *ICES Journal of Marine Science*, **66(2)**, 349-357.
- Orr, J.C., K. Caldeira, V. Fabry, J.P. Gattuso, P. Haugan, P. Lethodey, S. Pantoja, H.O. Pörtner, U. Riebesell, T. Trull, M. Hood, E. Urban, and W. Broadgate, 2009: Research priorities for understanding ocean acidification: Summary from the second symposium on the ocean in a high-CO2 world. *Oceanography*, **22(4)**, 182.

- Oskal, A., 2008: Old livelihoods in new weather: Arctic indigenous reindeer herders face the challenges of climate change. In: *Development Outreach, World Bank Institute Special Paper*. World Bank, pp. 22- 25.
- Østreng, W., 2006: The International Northern Sea Route Programme (INSROP): Applicable lessons learned, in *Polar Record*, Cambridge University Press **42(220)**, January, pp. 71-82
- Outridge, H. Sanei, Stern, Hamilton, and F. Goodarzi, 2007: Evidence for Control of Mercury Accumulation Rates in Canadian High Arctic Lake Sediments by Variations of Aquatic Primary Productivity. *Environmental Science & Technology*, **41(15)**, 5259-5265.
- Overeem, I. and J.P.M. Syvitski, 2010: Shifting discharge peaks in arctic rivers, 1977-2007. *Geografiska Annaler: Series A, Physical Geography*, **92(2)**, 285-296.
- Overland, J.E., J.A. Francis, E. Hanna, and M. Wang. 2012. The recent shift in early summer Arctic atmospheric circulation. *Geophysical Research Letters*. 39. L19804, doi: 10.1029/2012g1053268.
- Pagano, A.M., G.M. Durner, S.C. Amstrup, K.S. Simac, and G.S. York, 2012: Long-distance swimming by polar bears (*Ursus maritimus*) of the southern Beaufort Sea during years of extensive open water. *Canadian Journal of Zoology*, **90(5)**, 663-676.
- Parada, C., D.A. Armstrong, B. Ernst, S. Hinckley, and J.M. Orensanz, 2010: Spatial dynamics of snow crab (*Chionoecetes opilio*) in the eastern Bering Sea—putting together the pieces of the puzzle. *Bulletin of Marine Science*, **86(2)**, 413-437.
- Parkinson, A.J. and B. Evengard. 2009. Climate change, its impact on human health in the Arctic and the public health response to threats of emerging infectious diseases. *Global Health Action*. **2**, doi:10.3402/gha.v2i0.2075.
- Parkinson, A.J., 2009: Sustainable Development, Climate Change and Human Health in the Arctic. In: *UNESCO. Climate Change and Arctic Sustainable Development: Scientific, Social Cultural, and Educational Challenges*. UNESCO, Paris, pp. 156-163.
- Parkinson, A.J., M.G. Bruce, and T. Zulz, 2008: *International Circumpolar Surveillance, an Arctic network for the surveillance of infectious diseases*. **14(1)**.
- Parnikoza, I., P. Convey, I. Dykyy, V. Trokhymets, G. Milinevsky, O. Tyschenko, D. Inozemtseva, and I. Kozeretska, 2009: Current status of the Antarctic herb tundra formation in the Central Argentine Islands. *Global Change Biology*, **15(7)**, 1685-1693.
- Paxian, A., V. Eyring, W. Beer, R. Sausen, and C. Wright, 2010: Present-Day and Future Global Bottom-Up Ship Emission Inventories Including Polar Routes. *Environmental Science & Technology*, **44(4)**, 1333-1339.
- Payette, S., 2007: Contrasted dynamics of Northern Labrador tree lines caused by climate change and migrational lag. *Ecology*, **88(3)**, 770-780.
- Pearson, R.G., Phillips, S.J., Lorant, M.M., Beck, P.S.A., Damoulas, T., Knight, S.J. and Goetz, S.J. 2013. Shifts in Arctic vegetation and associated feedbacks under climate change. *Nature Climate Change* **3**, 673-677.
- Peck, L.S., D.K.A. Barnes, A.J. Cook, A.H. Fleming, and A. Clarke, 2009: Negative feedback in the cold: ice retreat produces new carbon sinks in Antarctica. *Global Change Biology*, **16(9)**, 2614-2623.
- Péron, C., H. Weimerskirch, and C.A. Bost, 2012: Projected poleward shift of king penguins' (*Aptenodytes patagonicus*) foraging range at the Crozet Islands, southern Indian Ocean. *Proceedings of the Royal Society B*, **279**, 2515-2523.
- Péron, C., M. Authier, C. Barbraud, K. Delord, D. Besson, and H. Weimerskirch, 2010: Interdecadal changes in at-sea distribution and abundance of subantarctic seabirds along a latitudinal gradient in the Southern Indian Ocean. *Global Change Biology*, **16(7)**, 1895-1909.
- Perrette, M., A. Yool, G.D. Quartly, and E.E. Popova, 2011: Near-ubiquity of ice-edge blooms in the Arctic. *Biogeosciences*, **8(2)**, 515-524.
- Peters G.P., Nilssen T.B., Lindholdt L., Eide m.S., Glomsrød S., Eide L.I., and Fuglested J.S., 2011 : Future emissions from shipping and petroleum activities in the Arctic. *Atmos. Chem. Phys.*, **11**, 5305-5320.
- Peters G.P., T.B. Nilssen, L. Lindholdt, M.S. Eide, S. Glomsrød, L.I. Eide, and Fuglested J.S., 2011: Future emissions from shipping and petroleum activities in the Arctic. *Atmos. Chem. Phys.*, **11**, 5305-5320.
- Peterson, B.J., R.M. Holmes, J.W. McClelland, C.J. Vörösmarty, R.B. Lammers, A.I. Shiklomanov, I.A. Shiklomanov, and S. Rahmstorf, 2002: Increasing River Discharge to the Arctic Ocean. *Science*, **298(5601)**, 2171-2173.
- Pohl, S., P. Marsh, and B. Bonsal, 2007: Modeling the Impact of Climate Change on Runoff and Annual Water Balance of an Arctic Headwater Basin. *Arctic*, **60(2)**, 173-186.
- Poloczanska, E.S., C.J. Brown, W.J. Sydeman, W. Kiessling, D. S. Schoeman, P.J. Moore, K. Brander, J.F. Bruno, L.B. Buckley, M.T. Burrows, C.M. Duarte, B.S. Halpern, J. Holding, C.V. Kappel, M.I. O'Connor, J.M.

- Pandolfi, C. Parmesan, F. Schwing, S. A. Thompson, and A.J. Richardson. 2013. Global imprint of climate change on marine life. *Nature Climate Change*, DOI: 10.1038/NCLIMATE1958
- Poppel, B. and J. Kruse, 2009: The importance of a mixed cash and harvest herding based economy to living the Arctic: an analysis based on Survey of Living Conditions in the Arctic (SLICA). In: *Quality of Life in the New Millennium: Advances in the Quality-of-Life Studies. Theory and Research*. Springer, Verlag, pp. 27-42.
- Portier, C.J., T.K. Thigpen, S.R. Carter, C.H. Dilworth, A.E. Grambsch, J. Gohlke, J. Hess, S.N. Howard, G. Luber, J.T. Lutz, T. Maslak, N. Prudent, M. Radtke, J.P. Rosenthal, T. Rowles, P.A. Sandifer, J. Scheraga, P.J. Schramm, D. Strickman, J.M. Trtanj, and P. Whung, 2010: *A Human Health Perspective On Climate Change: A Report Outlining the Research Needs on the Human Health Effects of Climate Change*. Research Triangle Park, NC:Environmental Health Perspectives/National Institute of Environmental Health Sciences.
- Pörtner, H.O. and M.A. Peck, 2010: Climate change effects on fishes and fisheries: towards a cause-and-effect understanding. *Journal of Fish Biology*, **77(8)**, 1745-1779.
- Post, E. and M.C. Forchhammer, 2008: Climate change reduces reproductive success of an Arctic herbivore through trophic mismatch. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **363(1501)**, 2369-2375.
- Post, E., C. Pedersen, C.C. Wilmers, and M.C. Forchhammer, 2008: Warming, plant phenology and the spatial dimension of trophic mismatch for large herbivores. *Proceedings of the Royal Society B: Biological Sciences*, **275(1646)**, 2005-2013.
- Post, E., J. Brodie, M. Hebblewhite, A.D. Anders, J.A.K. Maier, and C.C. Wilmers, 2009a: Global population dynamics and hot spots of response to climate change. *Bioscience*, **59(6)**, 489-497.
- Post, E., M.C. Forchhammer, M.S. Bret-Harte, T.V. Callaghan, T.R. Christensen, B. Elberling, A.D. Fox, O. Gilg, D.S. Hik, T.T. Høye, R.A. Ims, E. Jeppesen, D.R. Klein, J. Madsen, A.D. McGuire, S. Rysgaard, D.E. Schindler, I. Stirling, M.P. Tamstorf, N.J.C. Tyler, R. Van Der Wal, J. Welker, P.A. Wookey, N.M. Schmidt, and P. Aastrup, 2009b: Ecological dynamics across the arctic associated with recent climate change. *Science*, **325(5946)**, 1355-1358.
- Pouliot, J., P.E. Galand, C. Lovejoy and W.F. Vincent, 2009. Vertical structure of archaeal communities and the distribution of ammonia monooxygenase A gene variants in two meromictic High Arctic lakes. *Environmental Microbiology*. **11:687-699**.
- Prach, K., J. Košnar, J. Klimešová, and M. Hais, 2010: High Arctic vegetation after 70 years: A repeated analysis from Svalbard. *Polar Biology*, **33(5)**, 635-639.
- Prowse, T., K. Alfredsen, S. Beltaos, B. Bonsal, C. Duguay, A. Korhola, J. McNamara, W. Vincent, V. Vuglinsky, G. Weyhenmeyer, K. Walter Anthony, B. Bowden, V. Buzin, Y. Dibike, N. Gantner, L. Hinzman, L. Lia, T. Ouarda, R. Pientiz, J.D. Reist, M. Stickler, J. Weckström, and F. Wrona, 2011. Changing lake and river ice regimes: trends, effects, and implications. In: SWIPA, Snow, Water, Ice Permafrost in the Arctic. Scientific Assessment of the Arctic Monitoring and Assessment Program, Oslo, Norway, p. 6.1-6.52.
- Prowse, T.D. and Brown, K. 2010b. Hydro-ecological effects of changing Arctic river and lake ice covers: a review. *Hydrology Research* **41.6**: 454-461.
- Prowse, T.D. and K. Brown, 2010a: Appearing and disappearing lakes in the Arctic and their impacts on biodiversity. In: *Arctic Biodiversity Trends 2010 – Selected indicators of change*. [Kurvits, T., B. Alfthan, and E. Mork(eds.)]. CAFF International Secretariat, Akureyri, Iceland, pp. 68-70.
- Prowse, T.D., B.R. Bonsal, C.R. Duguay, D.O. Hessen, and V.S. Vuglinsky, 2007: River and Lake Ice. In: *Global Outlook for Ice & Snow*. United Nations Environment Programme, pp. 201.
- Prowse, T.D., C. Furgal, R. Chouinard, H. Melling, D. Milburn, and S.L. Smith, 2009: Implications of Climate Change for Economic Development in Northern Canada: Energy, Resource, and Transportation Sectors. *AMBIO: A Journal of the Human Environment*, **38(5)**, 272-281.
- Quayle, W. C., P. Convey, L. S. Peck, C. J. Ellis-Evans, H. G. Butler and H. J. Peat, 2013: Ecological Responses of Maritime Antarctic Lakes to Regional Climate Change. In: *Antarctic Peninsula Climate Variability: Historical and Paleoenvironmental Perspectives*. [E. Domack, A. Leventer, A. Burnett, R. Bindshadler, P. Convey, and M. Kirby (eds.)], American Geophysical Union: 159-170.
- Quesada, A. and Velazquez, D., 2012: Global Change Effects on Antarctic Lakes. In: *Effects of Global Warming on Freshwater Ecosystems of the World: what can be done to reduce negative impacts?* (Kumagai, M., Goldman, C.R. and Robarts, R.D. eds.). Wiley-Blackwell Ltd .pp 367-382
- Quesada, A. and Velazquez, D., 2013: Global Change Effects on Antarctic Freshwater Ecosystems: The Case of Maritime Antarctic Lakes. *Climatic Change and Global Warming of Inland Waters: Impacts and Mitigation for*

- Ecosystems and Societies, First Edition. Edited by Charles R. Goldman, Michio Kumagai and Richard D. Robarts. John Wiley & Sons. 17pp.
- Quesada, A., W.F. Vincent, E. Kaup, J.E. Hobbie, I. Laurion, R. Pienitz, J. López-Martínez, and J.J. Durán, 2006: Landscape control of high latitude lakes in a changing climate. In: *Trends in Antarctic Terrestrial and Limnetic Ecosystems. Antarctica as a Global Indicator*. [Bergstrom, D.M., P. Convey, and A.H.L. Huiskes(eds.)]. Springer, Dordrecht, pp. 221-252.
- Ragen, T.J., H.P. Huntington, and G.K. Hovelsrud, 2008: Conservation of arctic marine mammals faced with climate change. *Ecological Applications*, **18**(2), S166-S174.
- Rawlins, M.A., H. Ye, D. Yang, A. Shiklomanov, and K.C. McDonald, 2009a: Divergence in seasonal hydrology across northern Eurasia: Emerging trends and water cycle linkages. **114**, D18119.
- Rawlins, M.A., M.C. Serreze, R. Schroeder, X. Zhang, and K.C. McDonald, 2009b: Diagnosis of the record discharge of Arctic-draining Eurasian rivers in 2007. *Environmental Research Letters*, **4**(4), 045011.
- Regehr, E.V., C.M. Hunter, H. Caswell, S.C. Amstrup, and I. Stirling, 2010: Survival and breeding of polar bears in the southern Beaufort Sea in relation to sea ice. *The Journal of Animal Ecology*, **79**(1), 117-127.
- Regehr, E.V., N.J. Lunn, S.C. Amstrup, and I. Stirling, 2007: Effects of earlier sea ice breakup on survival and population size of polar bears in western Hudson Bay. *Journal of Wildlife Management*, **71**(8), 2673-2683.
- Regular, P.M., G.J. Robertson, W.A. Montevecchi, F. Shuhood, T. Power, D. Ballam, and J.F. Piatt, 2010: Relative importance of human activities and climate driving common murre population trends in the Northwest Atlantic. *Polar Biology*, **33**(9), 1215-1226.
- Reinert, E.S., I. Aslaksen, I.M.G. Eira, S.D. Mathiesen, H. Reindert and E.I. Turi. 2009. Adapting to climate change in Saami reindeer herding: the nation-state as problem and solution. In: W.N. Adger et al (2009): Adapting to climate change: Thresholds, Values, Governance, Pp. 417-432. Cambridge University Press. New York
- Reinert, H., S. Mathiesen and E. Reinert, 2008: Climate Change and Pastoral Flexibility: A Norwegian Saami Case. In: The Political Economy of Northern Regional Development – Yearbook [Winther, G., G. Duhaime, J. Kruse, C. Southcott, H. Aage (eds.)], TemaNord, ISBN:978-9289320160
- Reist, J.D., F.J. Wrona, T.D. Prowse, J.B. Dempson, M. Power, G. Kock, T.J. Carmichael, C.D. Sawatzky, H. Lehtonen, and R.F. Tallman, 2006: Effects of climate change and UV radiation on fisheries for Arctic freshwater and anadromous species. *AMBIO: A Journal of the Human Environment*, **35**(7), 402-410.
- Rennermalm, A., E. Wood, and T. Troy, 2010: Observed changes in pan-arctic cold-season minimum monthly river discharge. *Climate Dynamics*, **35**(6), 923-939.
- Revich, B. and M. Podolnaya, 2011: Thawing of permafrost may disturb historic cattle burial grounds in East Siberia. *Global Health Action*, **4**(0).
- Revich, B., 2008: Climate change alters human health in Russia. *Studies in Russian Economic Development*. 19(3) pp. 311-317, doi:10.1134/S1075700708030106.
- Revich, B., N. Tokarevich, and A.J. Parkinson, 2012: Climate change and zoonotic infections in the Russian Arctic. *International Journal of Circumpolar Health*, **71**, 18792.
- Revich, B.A. 2008. Climate change alters human health in Russia. *Studies in Russian Economic Development*. **19**(3), 311-317. doi:10.1134/S1075700708030106.
- Revich, B.A. and D.A. Shaposhnikov, 2010: Extreme temperature episodes and mortality in Yakutsk, East Siberia. *Rural and Remote Health*, **10**(2), 1338.
- Revich, B.A. and D.A. Shaposhnikov. 2012. Climate change, heat waves, and cold spells as risk factors for increased mortality in some regions of Russia. *Studies on Russian Economic Development*. **23**(2). Pp 195-207. Doi: 10.1134/S1075700712020116
- Rice, J.C. and S.M. Garcia, 2011: Fisheries, food security, climate change, and biodiversity: characteristics of the sector and perspectives on emerging issues. *ICES Journal of Marine Science*, **68**(6), 1343-1353.
- Roberts, D., W. R. Howard, A. D. Moy, J. L. Roberts, T. W. Trull, S. G. Bray, and R. R. Hopcroft. 2011. Interannual pteropod variability in sediment traps deployed above and below the aragonite saturation horizon in the Sub-Antarctic Southern Ocean. *Polar Biology* 34:1739-1750.
- Robertson, C., T.A. Nelson, D.E. Jelinski, M.A. Wulder, and B. Boots, 2009: Spatial-temporal analysis of species range expansion: The case of the mountain pine beetle, *Dendroctonus ponderosae*. *Journal of Biogeography*, **36**(8), 1446-1458.
- Robinson, R., T. Smith, B.J. Kirschhoffer, and C. Rosa, 2011: Polar bear (*Ursus maritimus*) cub mortality at a den site in northern Alaska. *Polar Biology*, **34**(1), 139-142.

- Rockwell, R.F. and L.J. Gormezano, 2009: The early bear gets the goose: climate change, polar bears and lesser snow geese in western Hudson Bay. *Polar Biology*, **32(4)**, 539-547.
- Rode, K., E. Peacock, M. Taylor, I. Stirling, E. Born, K. Laidre, and Ø Wiig, 2012: A tale of two polar bear populations: ice habitat, harvest, and body condition. *Population Ecology*, **54(1)**, 3-18.
- Rode, K.D., J.D. Reist, E. Peacock, and I. Stirling, 2010b: Comments in response to "Estimating the energetic contribution of polar bear (*Ursus maritimus*) summer diets to the total energy budget" by Dyck and Kebreab (2009). *Journal of Mammalogy*, **91(6)**, 1517-1523.
- Rode, K.D., S.C. Amstrup, and E.V. Regehr, 2010a: Reduced body size and cub recruitment in polar bears associated with sea ice decline. *Ecological Applications*, **20(3)**, 768-782.
- Rogers, A., N. Johnston, E. Murphy, and A. Clarke (eds.), 2012: *Antarctic Ecosystems: An Extreme Environment in a Changing World*. John Wiley & Sons, Ltd.
- Roturier, S. and M. Roué, 2009: Of forest, snow and lichen: Sámi reindeer herders' knowledge of winter pastures in northern Sweden. *Forest Ecology and Management*, **258(9)**, 1960-1967.
- Royles, J., M.J. A., Convey, P., Griffiths, H., Hodgson, D.A., Leng, M.J. and Charman, D.J., 2013. Plants and Soil Microbes Respond to Recent Warming on the Antarctic Peninsula. *Current Biology*, **23(17)**, <http://dx.doi.org/10.1016/j.cub.2013.07.011>.
- Rudberg, P.M., O. Wallgren and A.G. Swartling, 2012. Beyond generic adaptive capacity: exploring the adaptation space of the water supply and wastewater sector of the Stockholm region, Sweden. *Climatic Change*. **114 (3-4)**, 707-21.
- Rundqvist, S., H. Hedenås, A. Sandström, U. Emanuelsson, H. Eriksson, C. Jonasson, and T.V. Callaghan, 2011: Tree and shrub expansion over the past 34 years at the tree-line near Abisko, Sweden. *Ambio*, **40(6)**, 683-692.
- Russell, D. and A. Gunn, 2010: Reindeer herding. In: Arctic Biodiversity Trends 2010 – Selected indicators of change. [Kurvits, T., B. Alfthan, and E. Mork(eds.)]. CAFF International Secretariat, Akureyri, Iceland.
- Rybråten, S. and G. Hovelsrud, 2010: Local Effects of Global Climate Change: Differential Experiences of Sheep Farmers and Reindeer Herders in Unjárga/Nesseby, a Coastal Sámi Community in Northern Norway. In: *Community Adaptation and Vulnerability in Arctic Regions*. [Hovelsrud, G.K. and B. Smit(eds.)]. Springer Netherlands, pp. 313-333.
- Rylander, R. and R.S.F. Schilling, 2011: Diseases caused by organic dusts. In: 10. Respiratory System. Encyclopedia of occupational health and Safety. J.M. Stellman, Editor in Chief. International Labor Organization Geneva. 7 pp.
- Saba, G. K., O. Schofield, J. J. Torres, E. H. Ombres, and D. K. Steinberg. 2012: Increased feeding and nutrient excretion of adult Antarctic krill, *Euphausia superba*, exposed to enhanced carbon dioxide (CO₂). PLoS ONE 7:e52224.
- Sahanatien, V. and A.E. Derocher, 2012: Monitoring sea ice habitat fragmentation for polar bear conservation. *Animal Conservation*, **15: 397–406**. doi: 10.1111/j.1469-1795.2012.00529.x
- Salick, J. and N. Ross, 2009: Traditional peoples and climate change. *Global Environmental Change*, **19(2)**, 137-139.
- Salonen, J.S., H. Seppä, M. Väliranta, V.J. Jones, A. Self, M. Heikkilä, S. Kultti, and H. Yang, 2011: The Holocene thermal maximum and late-Holocene cooling in the tundra of NE European Russia. *Quaternary Research*, **75(3)**, 501-511.
- Schliebe, S., K.D. Rode, J.S. Gleason, J. Wilder, K. Proffitt, T.J. Evans, and S. Miller, 2008: Effects of sea ice extent and food availability on spatial and temporal distribution of polar bears during the fall open-water period in the Southern Beaufort Sea. *Polar Biology*, **31(8)**, 999-1010.
- Schmidt, K., A. Atkinson, S. Steigenberger, S. Fielding, M. Lindsay, D. Pond, G. Tarling, T. Klevjer, C. Allen, S. Nicol, and E. Achterberg, 2011: Seabed foraging by Antarctic krill: Implications for stock assessment, benthopelagic coupling, and the vertical transfer of iron. *Limnology and Oceanography*, **56(4)**, 1411–1428.
- Schneider, P. and S.J. Hook, 2010: Space observations of inland water bodies show rapid surface warming since 1985. *Geophysical Research Letters*, **37(22)**, L22405.
- Shaposhnikov, D., B. Revich, V. Meleshko, V. Govorkova, and T. Pavlova, 2010: Climate Change May Reduce Annual Temperature-Dependent Mortality in Subarctic: A Case Study of Archangelsk, Russian Federation. *Environment and Natural Resources Research*, **1(1)**, 75.
- Shearer, C. 2012. The political ecology of climate adaptation assistance: Alaska Natives, displacement, and relocation. *Journal of Political Ecology*. **19**. 174-183. [http://jpe.library.arizona.edu/volume_19/Shearer.pdf]
- Shearer, C., 2011: Kivalina: a climate change story. Haymarket Books, Chicago.

- Sherman, K., I.M. Belkin, K.D. Friedland, J. O'Reilly and K. Hyde, 2009. Accelerated warming and emergent trends in fisheries biomass yields of the world's large marine ecosystems. *Ambio*, **38**:215-224.
- Sherman, K., I.M. Belkin, K.D. Friedland, J. O'Reilly and K. Hyde, 2009: Accelerated warming and emergent trends in fisheries biomass yields of the world's large marine ecosystems. *Ambio*, **38**:215-224.
- Shiklomanov, A.I., R.B. Lammers, M.A. Rawlins, L.C. Smith, and T.M. Pavelsky, 2007: Temporal and spatial variations in maximum river discharge from a new Russian data set. **112**, G04S53.
- Shiyatov, S.G., M.M. Terent'ev, V.V. Fomin, and N.E. Zimmermann, 2007: Altitudinal and horizontal shifts of the upper boundaries of open and closed forests in the Polar Urals in the 20th century. *Russian Journal of Ecology*, **38**(4), 223-227.
- Shreeve, R.S., M.A. Collins, G.A. Tarling, C.E. Main, P. Ward, and N.M. Johnston, 2009: Feeding ecology of myctophid fishes in the northern Scotia Sea. *Marine Ecology-Progress Series*, **386**, 221-236.
- Sigler, M.F., M. Renner, S.L. Danielson, L.B. Eisner, R.R. Lauth, K.J. Kuletz, E.A. Logerwell, and Hunt George L. Jr., 2011: Fluxes, fins, and feathers: Relationships among the Bering, Chukchi, and Beaufort Seas in a time of climate change. *Oceanography*, **24**(3), 250-265.
- Sigurdsson, B.D., A. Snorrason, B. Kjartansson and J.A. Jonsson, 2007: Total area of planted forests in Iceland and their carbon stocks and fluxes. In: Effects of afforestation on ecosystems, land-scape and rural development. [Halldórsson G., Oddsdóttir E.S. and Eggertsson O. (eds.)], TemaNord 508, pp. 211-217.
- Simpson, S., S. Jennings, M. Johnson, J. Blanchard, P. Schun, D. Sims, and M. Genner, 2011: *Continental Shelf-Wide Response of a Fish Assemblage to Rapid Warming of the Sea*. Cell Press, pp. 1565-1570.
- Siniff, D. B., R. A. Garrott, J. J. Rotella, W. R. Fraser, and D. G. Ainley. 2008: Opinion: Projecting the effects of environmental change on Antarctic seals. *Antarctic Science* **20**:425-435.
- Slagstad, D., I.H. Ellingsen, and P. Wassmann, 2011: Evaluating primary and secondary production in an Arctic Ocean void of summer sea ice: An experimental simulation approach. *Progress in Oceanography*, **90**(1-4), 117-131.
- Slater, G.J., B. Figueirido, L. Louis, P. Yang, and B. van Valkenburgh, 2010: Biomechanical Consequences of Rapid Evolution in the Polar Bear Lineage. *PLoS ONE*, **5**(11), e13870, 1-7.
- Smith Jr, W., D. Ainley, R. Cattaneo-Vietti, and E. Hofmann, 2012: The Ross Sea Continental Shelf: regional biogeochemical cycles, trophic interactions, and potential future changes. In: *Antarctic ecosystems: an extreme environment in a changing world*. [Rogers, A., N. Johnston, E. Murphy, and A. Clarke(eds.)]. Blackwell Publishing, London, pp. 213-242; 7.
- Smith, J.A., D.A. Hodgson, M.J. Bentley, E. Verleyen, M.J. Leng, and S.J. Roberts, 2006: Limnology of two Antarctic epishelf lakes and their potential to record periods of ice shelf loss. *Journal of Paleolimnology*, **35**, 373-394.
- Smith, Jr. W., D. Ainley, R. Cattaneo-Vietti, and E. Hofmann, 2012a: The Ross Sea Continental Shelf: regional biogeochemical cycles, trophic interactions, and potential future changes. In: *Antarctic ecosystems: an extreme environment in a changing world*. [Rogers, A., N. Johnston, E. Murphy, and A. Clarke(eds.)]. Blackwell Publishing, London, pp. 213-242; 7.
- Smith, L.C., T.M. Pavelsky, G.M. MacDonald, A.I. Shiklomanov, and R.B. Lammers, 2007: Rising minimum daily flows in northern Eurasian rivers: A growing influence of groundwater in the high-latitude hydrologic cycle. **112**, G04S47.
- Smith, P.A., K.H. Elliott, A.J. Gaston, and H.G. Gilchrist, 2010: Has early ice clearance increased predation on breeding birds by polar bears? *Polar Biology*, **33**(8), 1149-1153.
- Smith, S.L. and D.W. Riseborough, 2010: Modelling the Thermal Response of Permafrost Terrain to right-of-way disturbance and climate warming. *Cold Regions Science and Technology*, **60**(1), 92-103
- Smol, J.P. and M.S.V. Douglas, 2007a: Crossing the final ecological threshold in high Arctic ponds. *Proceedings of the National Academy of Sciences of the United States of America*, **104**(30), 12395-12397.
- Smol, J.P. and M.S.V. Douglas, 2007b: From controversy to consensus: making the case for recent climate change in the Arctic using lake sediments. *Frontiers in Ecology and the Environment*, **5**(9), 466-474, doi:10.1890/060162.
- Solomon, S.D., D. Qin, M. Manning, Z. Chen, M. Marquie, K.B. Averyt, M. Tignor, and H.L. Miller (eds.), 2007: Climate change 2007: The physical science basis. Contribution of working group I to the fourth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge, United Kingdom, pp. 996.

- Søreide, J.E., E. Leu, J. Berge, M. Graeve, and S. Falk-Petersen, 2010: Timing of blooms, algal food quality and *Calanus glacialis* reproduction and growth in a changing Arctic. *Global Change Biology*, **16(11)**, 3154-3163.
- St. Jacques, J. and D.J. Sauchyn, 2009: Increasing winter baseflow and mean annual streamflow from possible permafrost thawing in the Northwest Territories, Canada. *Geophysical Research Letters*, **36(1)**, L01401.
- Stabeno, P.J., E.V. Farley Jr., N.B. Kachel, S. Moore, C.W. Mordy, J.M. Napp, J.E. Overland, A.I. Pinchuk, and M.F. Sigler., 2012b: Comparison of the physics of the northern and southern shelves of the eastern Bering Sea and some implications for the ecosystem. *Deep Sea Research Part II: Topical Studies in Oceanography*, **65-70**, 14-30.
- Stabeno, P.J., N.B. Kachel, S.E. Moore, J.M. Napp, M. Sigler, A. Yamaguchi, and A.N. Zerbini, 2012a: Comparison of warm and cold years on the southeastern Bering Sea shelf and some implications for the ecosystem. *Deep Sea Research Part II: Topical Studies in Oceanography*, **65-70**, 31-45.
- Stammler, F., 2005: *Reindeer Nomads Meet the Market: Culture, Property and Globalisation at the End of the Land*. ISBN-13: 978-3825880460
- Stephenson, S.R., L.C. Smith, and J.A. Agnew, 2011: Divergent long-term trajectories of human access to the Arctic. **1(3)**, 156-160.
- Stewart E.J., S.E.L. Howell, D. Draper, J. Yackel And A. Tivy (2007): Sea Ice in Canada's Arctic: Implications for Cruise Tourism. ARCTIC VOL. **60(4)**, P. 370-380
- Stewart, E. J., Howell, S. E. L, Dawson, J. D., Tivy, A. & Draper, D. (2010) Cruise tourism and sea ice in Canada's Hudson Bay region. ARCTIC. (**63**) 57-66.
- Stien, A., L.E. Loel, A. Mysterud, T. Severinsen, J. Kohler, and R. Langvatn, 2010: Icing events trigger range displacement in a high-arctic ungulate. *Ecology*, **91(3)**, 915-920.
- Stien, A., Rolf A. Ims, Steve D. Albon, Eva Fuglei, R. Justin Irvine, Erik Ropstad, Odd Halvorsen, Rolf Langvatn, Leif Egil Loe, Vebjørn Veiberg, Nigel G. Yoccoz . 2012. Congruent responses to weather variability in high Arctic herbivores.
- Stirling, I. and C.L. Parkinson, 2006: Possible Effects of Climate Warming on Selected Populations of Polar Bears (*Ursus maritimus*) in the Canadian Arctic. *Arctic*, **59(3)**, 261-275.
- Stirling, I., A.E. Derocher, W.A. Gough, and K. Rode, 2008b: Response to Dyck et al. (2007) on polar bears and climate change in western Hudson Bay. *Ecological Complexity*, **5(3)**, 193-201.
- Stirling, I., E. Richardson, G.W. Thiemann, and A.E. Derocher, 2008a: Unusual predation attempts of polar bears on ringed seals in the southern Beaufort Sea: Possible significance of changing spring ice conditions. *Arctic*, **61(1)**, 14-22.
- Stirling, I., T.L. McDonald, E.S. Richardson, E.V. Regehr, and S.C. Amstrup, 2011: Polar bear population status in the northern Beaufort Sea, Canada, 1971-2006. *Ecological Applications : A Publication of the Ecological Society of America JID - 9889808*, (**3**).
- Stram, D.L. and D.C.K. Evans, 2009: Fishery management responses to climate change in the North Pacific. *ICES Journal of Marine Science: Journal Du Conseil*, **66(7)**, 1633-1639.
- Stroeve, J., M.M. Holland, W. Meier, T. Scambos, and M. Serreze, 2007: Arctic sea ice decline: Faster than forecast. *Geophysical Research Letters*, **34**.
- Sundby, S. and O. Nakken, 2008: Spatial shifts in spawning habitats of Arcto-Norwegian cod related to multidecadal climate oscillations and climate change. *ICES Journal of Marine Science*, **65(6)**, 953-962.
- Swann, A. L., Fung, I. Y., Levis, S., Bonan, G., and Doney, S.2010. Changes in Arctic vegetation induce high-latitude warming through the greenhouse effect. *Proceedings of the National Academy of Sciences*, **107**, 1295-1300, doi:10.1073/pnas.0913846107.
- Sydneysmith, R., M. . Andrachuk, B. Smit, and G.K. Hovelsrud, 2010: Vulnerability and Adaptive Capacity in Arctic Communities. In: *Adaptive Capacity and Environmental Governance*. [Armitage, D. and R. Plummer(eds.)]. Springer, Berlin, pp. 133-156.
- Tan, A., J.C. Adam, and D.P. Lettenmaier, 2011: Change in spring snowmelt timing in Eurasian Arctic rivers. **116**, D03101
- Tanabe, Y., S. Kudoh, S. Imura, and M. Fukuchi, 2007: Phytoplankton blooms under dim and cold conditions in freshwater lakes of East Antarctica. *Polar Biology*, **31(2)**, 199-208.
- Tchebakova, N.M. , G.E. Rehfeldt, and E.I. Parfenova, 2009: From Vegetation Zones to Climates: Effects of Climate Warming on Siberian Ecosystems. In: *Permafrost Ecosystems: Siberian Larch Forest*, ISBN:978-1-4020-9692-1

- Teng, Y., Z. Xu, Y. Luo, and F. Reverchon, 2012: How do persistent organic pollutants be coupled with biogeochemical cycles of carbon and nutrients in terrestrial ecosystems under global climate change? *Journal of Soils and Sediments*, **12(3)**, 411-419.
- Teran, T., L. Lamon, and A. Marcomini. 2012: Climate change effects on POPs' environmental behavior: a scientific perspective for future regulatory actions. *Atmospheric Pollution Research*. **3**. 466-476.
- Terauds, A., S.L. Chown, F. Morgan, H. J. Peat, D.J. Watts, H. Keys, P. Convey, and D.M. Bergstrom, 2012: Conservation biogeography of the Antarctic. *Diversity and Distributions*, **18(7)**, 726-741.
- the Arctic induced by climate change. *Nature Climate Change* **1**, 255-260.
- Thiemann, G.W., S.J. Iverson, and I. Stirling, 2008: Polar Bear Diets and Arctic Marine Food Webs: Insights from Fatty Acid Analysis. *Ecological Monographs*, **78(4)**, 591-613.
- Tholstrup, L. and R.O. Rasmussen, 2009. Climate Change - and the North Atlantic. 130 pp. Norden, Stockholm, Sweden.
- Thompson, M., Wrona, F., Prowse, T., 2012: Shifts in Plankton, Nutrient and Light Relationships in Small Tundra Lakes Caused by Localized Permafrost Thaw. *Arctic*, **65(4)**.
- Thompson, M.S., S.V. Kokelj, T.D. Prowse and F.J. Wrona, 2008. The impact of sediments derived from thawing permafrost on tundra lake water chemistry: An experimental approach. In D.L. Kane and K.M. Hinkel (eds.), *Proceedings of the Ninth International Conference on Permafrost*, Fairbanks, Alaska. p. 1763-1768.
- Thrush, S.F. and V.J. Cummings, 2011: Massive icebergs, alteration in primary food resources and change in benthic communities at Cape Evans, Antarctica. *Marine Ecology-an Evolutionary Perspective*, **32(3)**, 289-299.
- Tokarevich, N., A. Tronin, O. Blinova, R. Buzinov, V. Boltenkov, E. Yurasova, and J. Nurse, 2011: The impact of climate change on the expansion of *Ixodes persulcatus* habitat and the incidence of tickborne encephalitis in the north of European Russia. *Global Health Action*, **4**.
- Tømmervik, H., B. Johansen, J.A. Riseth, S.R. Karlsen, B. Solberg, and K.A. Høgda, 2009: Above ground biomass changes in the mountain birch forests and mountain heaths of Finnmarksvidda, northern Norway, in the period 1957-2006. *Forest Ecology and Management*, **257(1)**, 244-257.
- Tømmervik, H., J.W. Bjerke, E. Gaare, B. Johansen, and D. Thannheiser, 2012: Rapid recovery of recently overexploited winter grazing pastures for reindeer in northern Norway. *Fungal Ecology*, **5(1)**, 3-15.
- Towns, L., A.E. Derocher, I. Stirling, and N.J. Lunn, 2010: Changes in Land Distribution of Polar Bears in Western Hudson Bay. *Arctic*, **63(2)**, 206-212.
- Towns, L., A.E. Derocher, I. Stirling, N.J. Lunn, and D. Hedman, 2009: Spatial and temporal patterns of problem polar bears in Churchill, Manitoba. *Polar Biology*, **32(10)**, 1529-1537.
- Trathan, P. and K. Reid (2009). "Exploitation of the marine ecosystem in the sub-Antarctic: historical impacts and current consequences." *Papers and Proceeding of the Royal Society of Tasmania* **143: 9-14**.
- Trathan, P., P. Fretwell, and B. Stonehouse, 2011: First recorded loss of an emperor penguin colony in the recent period of Antarctic regional warming: implications for other colonies. *PLoS ONE*, **6**, e14738.
- Trathan, P.N. and D. Agnew, 2010: Climate change and the Antarctic marine ecosystem: an essay on management implications. *Antarctic Science*, **22(4)**, 387-398.
- Trathan, P.N., J. Forcada, and E.J. Murphy, 2007: Environmental forcing and Southern Ocean marine predator populations: effects of climate change and variability. *Philosophical Transactions of the Royal Society B-Biological Sciences*, **362**, 2351-2365.
- Trathan, P.N., J. Forcada, and E.J. Murphy, 2012: Environmental forcing and Southern Ocean marine predator populations. In: *Antarctic Ecosystems*. [Rogers, A., N. Johnston, E. Murphy, and A. Clarke(eds.)]. Blackwell Publishing, Ltd, pp. 335-354.
- Tremblay, J., D. Robert, D.E. Varela, C. Lovejoy, G. Darnis, R.J. Nelson, and A.R. Sastri, 2012: Current state and trends in Canadian Arctic marine ecosystems: I. Primary production. *Climatic Change*, **115(1)**, 161-178.
- Trivelpiece, W.Z., J.T. Hinke, A.K. Miller, C.S. Reiss, S.G. Trivelpiece, and G.M. Watters, 2011: Variability in krill biomass links harvesting and climate warming to penguin population changes in Antarctica. *Proceedings of the National Academy of Sciences of the United States of America*, **108(18)**, 7625-7628.
- Turner, J. A., N. E. Barrand, T. J. Bracegirdle, P. Convey, D. A. Hodgson, M. Jarvis, A. Jenkins, G. Marshall, M. P. Meredith, H. Roscos, J. Shanklin, J. French, H. Goosse, M. Guglielmin, J. Gutt, S. Jacobs, M. C. Kennicutt, V. Masson-Delmotte, P. Mayewski, F. Navarro, S. Robinson, T. Scambos, M. Sparrow, C. Summerhayes, K. Speer, and A. Klepnikov. 2013. Antarctic climate change and the environment: An update. *Polar Record*:**1-23**.

- Turner, J., R.A. Bindschadler, P. Convey, G. Di Prisco, E. Fahrbach, J. Gutt, D.A. Hodgson, P.A. Mayewski, and C.P. Summerhayes (eds.), 2009: *Antarctic Climate Change and the Environment*. Scientific Committee on Antarctic Research, ISBN 978-0-948277-22-1, Cambridge, pp. 526.
- Turunen, M., P. Soppela, H. Kinnunen, M.-. Sutinen, and F. Martz, 2009: Does climate change influence the availability and quality of reindeer forage plants? *Polar Biology*, **32(6)**, 813-832.
- Tyler, N.J.C., 2010: Climate, snow, ice, crashes, and declines in populations of reindeer and caribou (*Rangifer tarandus* L.). *Ecological Monographs*, **80(2)**, 197-219.
- Tyler, N.J.C., J.M. Turi, M.A. Sundset, K.S. Bull, M.N. Sara, E. Reinert, N. Oskal, C. Nellemann, J.J. McCarthy, S.D. Mathiesen, M.L. Martello, O.H. Magga, G.K. Hovelsrud, I. Hanssen-Bauer, N.I. Eira, I.M.G. Eira, and R.W. Corell, 2007: Saami reindeer pastoralism under climate change: Applying a generalized framework for vulnerability studies to a sub-arctic social-ecological system. *Global Environmental Change-Human and Policy Dimensions*, **17(2)**, 191-206.
- Tyler, N.J.C., M.C. Forchhammer, and N.A. Øritsland, 2008: Nonlinear effects of climate and density in the dynamics of a fluctuating population of reindeer. *Ecology*, **89(6)**, 1675-1686.
- UNEP/AMAP, 2011: Climate Change and POPs: Predicting the Impacts. Report of the UNEP/AMAP Expert Group. Secretariat of the Stockholm Convention. Geneva 62 pp.
- USARC. 2010. Report on Goals and Objectives for Arctic Research 2009-2010. US Arctic Research Commission. Arlington, VA. 52 pp.
- Valdimarsson, H., O.S. Astthorsson, and J. Pálsson, 2012: Hydrographic variability in Icelandic waters during recent decades and related changes in distribution of some fish species. *ICES Journal of Marine Science*, **69(5)**, 815-825.
- Valdivia C, Seth et.al.: Adapting to Climate Change in Andean Ecosystems: Landscapes, Capitals, and Perceptions Shaping Rural Livelihood Strategies and Linking Knowledge Systems. *Annals of the Association of American Geographers*, **100(4)**, (2010).
- Van Bogaert, R., C. Jonasson, M. De Dapper, and T.V. Callaghan, 2010: Range expansion of thermophilic aspen (*Populus tremula* L.) in the Swedish subarctic. *Arctic, Antarctic, and Alpine Research*, **42(3)**, 362-375.
- Van Bogaert, R., K. Haneca, J. Hoogesteger, C. Jonasson, M. De Dapper, and T.V. Callaghan, 2011: A century of tree line changes in sub-Arctic Sweden shows local and regional variability and only a minor influence of 20th century climate warming. *Journal of Biogeography*, **38(5)**, 907-921.
- van Der Wal, R., S. Sjögersten, S.J. Woodin, E.J. Cooper, I.S. Jónsdóttir, D. Kuijper, T.A.D. Fox, and A.D. Huiskes, 2007: Spring feeding by pink-footed geese reduces carbon stocks and sink strength in tundra ecosystems. *Global Change Biology*, **13(2)**, 539-545.
- Veillette, J., D.R. Mueller, D. Antoniadis, and W.F. Vincent, 2008: Arctic epishelf lakes as sentinel ecosystems: Past, present and future. **113**, G04014.
- Velázquez, D., A. Frias, M.A. Lezcano and A. Quesada, 2013: Ecological relationships and stoichiometry within a maritime Antarctic watershed. *Antarctic Science* **25(2)**, 191-197
- Verbyla, D., 2008: The greening and browning of Alaska based on 1982-2003 satellite data. *Global Ecology and Biogeography*, **17(4)**, 547-555.
- Vert-pre, K.A., R.O. Amoroso, O.P. Jensen, and R. Hilborn. 2013. Frequency and intensity of productivity regime shifts in marine stocks. *Proceedings of the National Academy of Sciences*, **110(5)**, 1779-1784.
- Vieira, G., J. Bockheim, M. Guglielmin, M. Balks, A.A. Abramov, J. Boelhouwers, N. Cannone, L. Ganzert, D.A. Gilichinsky, S. Goryachkin, J. Lopez-Martinez, I. Meiklejohn, R. Raffi, M. Ramos, C. Schaefer, E. Serrano, F. Simas, R. Sletten, D. Wagner, 2010: Thermal State of permafrost and active-layer monitoring in the Antarctic: advances during the International Polar Year 2007-09. *Permafrost and Periglacial Processes*, **21(2)**: 182-197
- Vikebø, F.B., Huseø, Å., Slotte, A., Stenevik, E.K., and Lien, V.S., 2010: Effect of hatching date, vertical distribution and interannual variation in physical forcing on northward displacement and temperature conditions of Norwegian spring-spawning herring larvae. *ICES Journal of Marine Science*, **67**, 1948-1956.
- Vikhamar-Schuler, D., I. Hanssen-Bauer, T.V., Schuler, S.D. Mathiesen, and M. Lehning. 2013.: Use of a multilayer snow model to assess grazing conditions for reindeer. *Annals of Glaciology*. 54(62) pp 214-226.
- Vincent, W., J. Hobbie, J. Laybourn-Parry, 2008: Introduction to the Limnology of High-latitude Lake and River Ecosystems. In: *Polar lakes and Rivers: Limnology of Arctic and Antarctic Aquatic Ecosystems*. [Laybourn-Parry, J. and J. Laybourn-Parry(eds.)]. Oxford University Press, pp. 1-18.

- Vincent, W.F., L.G. Whyte, C. Lovejoy, C.W. Greer, I. Laurion, C.A. Suttle, J. Corbeil, and D.R. Mueller, 2009: Arctic microbial ecosystems and impacts of extreme warming during the International Polar Year. *Polar Science*, **3(3)**, 171-180.
- Virginia, R.S. and K.S. Yalowitz, 2012: *A New Paradigm for Arctic Health: Challenges and Responses to Rapid Climate, Environmental, and Social Change*. In: Workshop Report for May 23-25, Applied Circumpolar Policy [Virginia, R.S. and Yalowitz, K.S. (eds.)]. Dartmouth College, Hanover, N.H.
- Voggesser, G., K. Lynn, J. Daigle, F. Lake, D. Ranco. 2013 Cultural impacts to tribes from climate change influences on forests. Special Issue Climatic Change. Vol. 12. Issue 3. 615-626.
- Von Biela, V., R., C. E. Zimmerman, and L. L. Moulton, 2011. Long-term increases in young-of-the-year growth of Arctic cisco *Coregonus autumnalis* and environmental influences. *Journal Fish Biology* 78(1): 39-56.
- Vors, L.S. and M.S. Boyce, 2009: Global declines of caribou and reindeer. *Global Change Biology*, **15(11)**, 2626-2633.
- Vuojala-Magga, T., M. Turunen, T. Ryyppö, and M. Tennberg, 2011: Resonance Strategies of Sámi Reindeer Herders in Northernmost Finland during Climatically Extreme Years. *Arctic*, **64(2)**.
- Vyverman, W., E. Verleyen, A. Wilmotte, D.A. Hodgson, A. Willems, K. Peeters, B. Van de Vijver, A. De Wever, F. Leliaert, and K. Sabbe, 2010: Evidence for widespread endemism among Antarctic micro-organisms. *Polar Science*, **4(2)**, 103-113. 10.1016/j.polar.2010.03.006.
- Walker, D.A., M.O. Leibman, H.E. Epstein, B.C. Forbes, U.S. Bhatt, M.K. Raynolds, J.C. Comiso, A.A. Gubarkov, A.V. Khomutov, G.J. Jia, E. Kaarlejärvi, J.O. Kaplan, T. Kumpula, P. Kuss, G. Matyshak, N.G. Moskalenko, P. Orekhov, V.E. Romanovsky, N.G. Ukraintseva, and Q. Yu, 2009: Spatial and temporal patterns of greenness on the Yamal Peninsula, Russia: Interactions of ecological and social factors affecting the Arctic normalized difference vegetation index. *Environmental Research Letters*, **4(4)**, 16.
- Walker, D.A., P. Kuss, H.E. Epstein, A.N. Kade, C.M. Vonlanthen, M.K. Raynolds, and F.J. Daniëls, 2011: Vegetation of zonal patterned-ground ecosystems along the North America Arctic bioclimate gradient. *Applied Vegetation Science*, **14(4)**, 440-463.
- Walsh, J.J., D.A. Dieterle, F.R.Chen, J.M. Lenes, W. Maslowski, J.J. Cassano, T.E. Whitlege, D. Stockwell, M. Flint, R.N. Sukhanova, J. Christensen. 2011. Trophic cascades and future harmful algal blooms within ice-free Arctic seas north of Bering Strait. A simulation analysis. *Progress in Oceanography*. 91 (2011) 312-343. <http://www.sciencedirect.com/science/article/pii/S0079661111000164>
- Walter, K.M., J.P. Chanton, Chapin, F.S., III, E.A.G. Schuur, and S.A. Zimov, 2008: Methane production and bubble emissions from arctic lakes: Isotopic implications for source pathways and ages. **113**, G00A08.
- Walter, K.M., L.C. Smith, and F. Stuart Chapin, 2007a: Methane bubbling from northern lakes: present and future contributions to the global methane budget. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, **365(1856)**, 1657-1676.
- Walter, K.M., M.E. Edwards, G. Grosse, S.A. Zimov and F.S. Chapin III, 2007b. Thermokarst lakes as a source of atmospheric CH₄ during the last deglaciation. *Science*, **318:633-636**.
- Waluda, C.M., M.A. Collins, A.D. Black, I.J. Staniland, and P.N. Trathan, 2010: Linking predator and prey behaviour: contrasts between Antarctic fur seals and macaroni penguins at South Georgia. *Marine Biology*, **157(1)**, 99-112.
- Walvoord, M.A. and R.G. Striegl, 2007: Increased groundwater to stream discharge from permafrost thawing in the Yukon River basin: Potential impacts on lateral export of carbon and nitrogen. *Geophysical Research Letters*, **34(12)**, L12402.
- Wang and Overland 2011. I do not have access to the literature databasae and cannot search this ref. But it was in the *Geophysical Research Letters*
- Wasley, J., S.A. Robinson, M. Popp, and C.E. Lovelock, 2006: Some like it wet – biological characteristics underpinning tolerance of extreme water events in Antarctic bryophytes. *Functional Plant Biology*, **33**, 443-455.
- Wassman, P., C. M. Duarte, S. Agusti, and M. K. Sejr, 2011: Footprints of climate change in the Arctic marine ecosystem. *Global Change Biology*, **17**, 1235-1249.
- Wassmann, P., 2011: Arctic marine ecosystems in an era of rapid climate change. *Progress in Oceanography*, **90**, 1-17.
- Watson, S., L.S. Peck, P.A. Tyler, P.C. Southgate, K.S. Tan, R.W. Day, and S.A. Morley, 2012: Marine invertebrate skeleton size varies with latitude, temperature and carbonate saturation: implications for global change and ocean acidification. *Global Change Biology*, **18(10)**, 3026-3038.

- Weatherhead, E., S. Gearheard, and R.G. Barry, 2010: Changes in weather persistence: Insight from Inuit knowledge. *Global Environmental Change*, **20(3)**, 523-528.
- Weimerskirch, H., M. Louzao, S. de Grissac, and K. Delord, 2012: Changes in wind pattern alter albatross distribution and life-history traits. *Science*, **335**, 211-214.
- Weimerskirch, H., P. Inchausti, C. Guinet, and C. Barbraud, 2003: Trends in bird and seal populations as indicators of a system shift in the Southern Ocean. *Antarctic Science*, **15(2)**, 249-256.
- Wenzel, G.W., 2009: Canadian Inuit subsistence and ecological instability? if the climate changes, must the Inuit? *Polar Research*, **28(1)**, 89-99.
- West, J. and G. Hovelsrud, 2010: Cross-scale Adaptation Challenges in the Coastal Fisheries: Findings from Lebesby, Northern Norway. *Arctic*, **63(3)**.
- Whyte, K.P. 2013: Justice forward: tribes, climate adaptation and responsibility. Special Issue Climatic Change. Vol. 12, Issue 3. 517-530.
- Wiedenmann, J., K. Cresswell, and M. Mangel, 2008: Temperature-dependent growth of Antarctic krill: predictions for a changing climate from a cohort model. *Mar Ecol Prog Ser*, **358**, 191-202.
- Wiedenmann, J., K.A. Cresswell, and M. Mangel, 2009: Connecting recruitment of Antarctic krill and sea ice. *Limnology and Oceanography*, **54(3)**, 799-811.
- Wiig, O., J. Aars, and E.W. Born, 2008: Effects of climate change on polar bears. *Science Progress*, **91(Pt 2)**, 151-173.
- Wildcat, D. 2009. Red Alert!: Saving the Planet with Indigenous Knowledge. Golden, CO: Fulcrum Publishing. 143 pp.
- Wilderbuer, T., W. Stockhausen, N. Bond, 2013: Updated analysis of flatfish recruitment response to climate variability and ocean conditions in the Eastern Bering Sea. *Deep-Sea Research II*, **94**, 157-164.
- Williams, T. and P. Hardison. 2013. Culture, law, risk and governance: contexts of traditional knowledge in climate change adaptation. Special Issue Climatic Change. Vol. 12. Issue 3. 531d-544.
- Williams, T.M., S.R. Noren and M. Glenn. 2011. Extreme physiological adaptations as predictors of climate-change sensitivity in the narwhal, *Monodon monoceros*. *Mar. Mamm. Sci.*, **27**, 334-349.
- Wilson, W.J., and O.A. Ormseth, 2009: A new management plan for Arctic waters of the United States. *Fisheries*, **34(11)**, 555-558.
- Wolf, A., T.V. Callaghan, and K. Larson, 2008: Future changes in vegetation and ecosystem function of the Barents Region. *Climatic Change*, **87(1-2)**, 51-73.
- Wramneby, A., B. Smith, and P. Samuelsson, 2010: Hot spots of vegetation-climate feedbacks under future greenhouse forcing in Europe. *Journal of Geophysical Research D: Atmospheres*, **115(21)**, 12.
- Xu, L., R.B. Myneni, F.S. Chapin III, T.V. Callaghan, J.E. Pinzon, C.J. Tucker, Z. Zhu, J. Bi, P. Ciais, H. Tommervik, E.S. Euskirchen, B.C. Forbes, S.L. Piao, B.T. Anderson, S. Ganguly, R.R. Nemani, S.J. Goetz, P.S.A. Beck, A.G. Bunn, C. Cao, and J.C. Stroeve, 2013: Temperature and vegetation seasonality diminishment over northern lands. *Nature Climate Change*, **advance online publication**.
- Yamamoto-Kawai, M., F.A. McLaughlin, E.C. Carmack, S. Nishino, and K. Shimada, 2009: Aragonite undersaturation in the Arctic Ocean: Effects of ocean acidification and sea ice melt. *Science*, **326(5956)**, 1098-1100.
- Yarie, J., 2008: Effects of moisture limitation on tree growth in upland and floodplain forest ecosystems in interior Alaska. *Forest Ecology and Management*, **256(5)**, 1055-1063.
- Ye, B., D. Yang, Z. Zhang, and D.L. Kane, 2009: Variation of hydrological regime with permafrost coverage over Lena Basin in Siberia. **114**, D07102.
- Zarnetske, J.P., M.N. Gooseff, W.B. Bowden, M.J. Greenwald, T.R. Brosten, J.H. Bradford, and J.P. McNamara, 2008: Influence of morphology and permafrost dynamics on hyporheic exchange in arctic headwater streams under warming climate conditions. *Geophysical Research Letters*, **35(2)**, L02501.
- Zeng H., Jia G., Forbes B., 2013. Shifts in Arctic phenology in response to climate and anthropogenic factors as detected from multiple satellite time series. *Environmental Research Letters* **8**,
- Zerbini, A., P. Clapham, and P. Wade, 2010: Assessing plausible rates of population growth in humpback whales from life-history data. *Marine Biology*, **157(6)**, 1225-1236, DOI: 10.1007/s00227-010-1403-y.
- Zhang, J., Y.H. Spitz, M. Steele, C. Ashjian, R. Campbell, L. Berline, and P. Matrai, 2010: Modeling the impact of declining sea ice on the Arctic marine planktonic ecosystem. **115**, C10015.

- Zhang, K., J.S. Kimball, E.H. Hogg, M. Zhao, W.C. Oechel, J.J. Cassano, and S.W. Running, 2008: Satellite-based model detection of recent climate-driven changes in northern high-latitude vegetation productivity. *Journal of Geophysical Research G: Biogeosciences*, **113(3)**, 13.
- Zhang, X., J. He, J. Zhang, I. Polyakov, I., R. Gerdes, J. Inoue and P. Wu, 2013. Enhanced poleward moisture transport and amplified high-latitude wetting trend. *Nature Climate Change*, **3: 47-51**
- Zhukova, N.G., V.N. Nesterova, I.P. Prokopchuk, and G.B. Rudneva, 2009: Winter distribution of euphausiids (*Euphausiacea*) in the Barents Sea (2000–2005). *Deep Sea Research Part II: Topical Studies in Oceanography*, **56(21–22)**, 1959-1967.

Table 28-1: Annually averaged changes in inland and maritime transportation accessibility by mid-century (2045–2059) versus baseline (2000–2014).

	Change in winter road-accessible land area (km²) (2,000-kg GVWR vehicle)	Change in maritime-accessible ocean area (km²) (Type A vessel)—current EEZ
Canada	-13%	19%
Finland	-41%	0%
Greenland	-11%	28%
Iceland	-82%	<1%
Norway	-51%	2%
Russia	-13%	16%
Sweden	-46%	0%
USA (Alaska)	-29%	5%
High seas	n/a	406%
Total	-14%	23%

Do Not Cite, Quote, or Distribute Prior to Public Release on 31 March 2014




Figure 28-1: Location maps of the north and south polar regions.
Credit: P. Fretwell, British Antarctic Survey.

Do Not Cite, Quote, or Distribute Prior to Public Release on 31 March 2014




Figure 28-2: Temporal change in onset of flowering (plants), median date of emergence (arthropods) and clutch initiation dates (birds) in high-Arctic Greenland. Red dots are statistically significant, blue dots are not. Source: Høye *et al.*, 2007. [Illustration to be redrawn to conform to IPCC publication specifications.]

Do Not Cite, Quote, or Distribute Prior to Public Release on 31 March 2014




Figure 28-3: Significant changes ($p < 0.01$) in photosynthetically active period NDVI between 1982 and 2012. Source: Xu et al., 2013. [Illustration to be redrawn to conform to IPCC publication specifications.]




Figure 28-4: Projected duration of the navigation period (days) over the Northwest Passage and Northern Sea Route. Source: Mokhov and Khon, 2008. **[Illustration to be redrawn to conform to IPCC publication specifications.]**